

D-instantons and Indefinite Theta series

Boris Pioline

CERN & LPTHE



KIPMU, Oct 21, 2016

based on arXiv:1605.05945,1606.05495 with S. Alexandrov, S. Banerjee and J. Manschot

- 1 Physics introduction: D-instanton corrections to HM moduli spaces
- 2 Math introduction: indefinite theta series
- 3 Indefinite theta series and generalized error functions
- 4 Back to physics: S-duality in HM moduli space

- 1 Physics introduction: D-instanton corrections to HM moduli spaces
- 2 Math introduction: indefinite theta series
- 3 Indefinite theta series and generalized error functions
- 4 Back to physics: S-duality in HM moduli space

In type IIB string theory compactified on a CY threefold X , the moduli space decomposes into two factors:


$$\mathcal{M} = \mathcal{M}_V(X) \times \mathcal{M}_H(X)$$

- The **vector-multiplet moduli space** $\mathcal{M}_V(X)$ describes the **complex structure moduli** of X . It carries a **Kähler** metric derived from the Kähler potential $\mathcal{K}_V = -\log \int_X \Omega \wedge \bar{\Omega}$.
- The **hypermultiplet moduli space** $\mathcal{M}_H(X)$ describes the **Kähler moduli of X** , along with the **string coupling** $g = e^\phi$, the **RR axions** $C \in H_{\text{even}}(X, \mathbb{R})/H_{\text{even}}(X, \mathbb{Z})$ and the **NS axion** $\sigma \in \mathbb{R}/\mathbb{Z}$. It carries a **quaternion-Kähler** metric (which is NOT a Kähler, not even complex !). In the weak coupling limit $g \rightarrow 0$,

$$\mathcal{M}_H(X) \sim_{g \rightarrow 0} \text{c-map}(\mathcal{M}_V(\hat{X}))$$

Physics Introduction II

- For g finite, one expects $\mathcal{O}(e^{-1/g})$ corrections from **Euclidean D-branes**, i.e. stable objects in the derived category of stable sheaves on X , weighted by the corresponding **generalized Donaldson-Thomas invariants** $\Omega(\gamma)$; as well as $\mathcal{O}(e^{-1/g^2})$ corrections from **NS five-branes**, whose mathematical description is unknown.
- The effect of Euclidean D-branes on the QK metric on $\mathcal{M}_H(X)$ is well understood in terms of the **twistor space** \mathcal{Z} , a \mathbb{P}^1 bundle over $\mathcal{M}_H(X)$ equipped with a canonical **complex contact structure**:



Alexandrov BP Saueressig Vandoren 2008; Gaiotto Moore Neitzke 2008

Physics Introduction III

- An key constraint is that $\mathcal{M}_H(X)$ should admit a smooth QK manifold with an **isometric action of $SL(2, \mathbb{Z})$** , originating from S-duality in type IIB string theory. Equivalently, the contact structure on the twistor space must be invariant under $SL(2, \mathbb{Z})$.

Robles-Llana, Rocek, Saueressig, Theis, Vandoren 2006

- S-duality relates D1-instantons to F1-instantons (hence DT invariants to GW invariants); D5-instanton to NS5-instantons; but maps D3-instantons to themselves. Thus it gives an important constraint on **DT-invariants for pure dimension 2 sheaves supported on a divisor $\mathcal{P} \subset X$** , with $p = [\mathcal{P}] \in H_4(\mathcal{X}, \mathbb{Z}) \cong \Lambda$.

Physics Introduction IV

- Indeed, these same invariants count **D4-D2-D0 black hole microstates** in type IIA/X, and are described by given by the (modified) elliptic genus of a suitable $N = (4, 0)$ SCFT:

$$Z_p(\tau, c) = \sum_{q_a \in \Lambda^*, q_0 \in \mathbb{Z}} \Omega(0, p^a, q_a, q_0) e^{-2\pi\tau_2 |Z(\gamma)| - 2\pi i(\tau_1 q_0 + c^a q_a)}$$

is a Jacobi form of weight $(-3/2, 1/2)$.

Maldacena Strominger Witten 1997

Physics Introduction V

- Invariance under spectral flow $(q_a, q_0) \mapsto (q_a - \kappa_{abc} p^b \epsilon^c, q_0 - \epsilon^a q_a + \frac{1}{2} \kappa_{abc} p^a \epsilon^b \epsilon^c)$ implies the theta series decomposition

$$Z_p(\tau, c^a) = \sum_{\mu \in \Lambda^* / \Lambda} h_{p, \mu}(\tau) \theta_{p, \mu}(\tau, c)$$

where $h_{p, \mu}(\tau)$ is a vector-valued holomorphic modular form of weight $-\frac{b_2}{2} - 1$, and $\theta_{p, \mu}(\tau, c)$ a Siegel theta series of signature $(1, b_2 - 1)$ and weight $(\frac{b_2 - 1}{2}, \frac{1}{2})$.

Denef Moore; Gaiotto Strominger Yin; de Boer Cheng Dijkgraaf Manschot Verlinde

- Explicitly,

$$\theta_{p, \mu}(\tau, c) = \sum_{k \in \Lambda + \mu} e^{-i\pi\tau Q(k_-) - i\pi\bar{\tau} Q(k_+)}$$

where $Q(k) = \kappa_{abc} p^a k^b k^c$ is a quadratic form of signature $(1, b_2 - 1)$, k_+ is the projection of k along the timelike vector $t \in \mathbb{R}^{1, b_2 - 1}$ parametrizing the Kähler cone, and $k_- = k - k_+$.

- The indefinite theta series $\theta_{\rho,\mu}(\tau, c)$, corresponding to the sum over fluxes on D3-brane wrapping a fixed divisor ρ , is intrinsically **non-holomorphic**. Yet, it must somehow correct the **holomorphic contact structure** on \mathcal{Z} ...
- Similarly, the sum over fluxes on a pair of two-D-branes wrapping a divisor $\mathcal{P}_1 \cup \mathcal{P}_2$ will lead to an indefinite theta series of signature $(2, 2b_2 - 2)$. Yet it must somehow correct the holomorphic contact structure on \mathcal{Z} ...
- In the rest of this talk, I will describe **a general procedure for constructing holomorphic theta series of arbitrary signature $(r, n - r)$ and finding their modular properties**. Time permitting, I will explain how to apply this procedure to show that D3-instanton corrections are consistent with S-duality.

- 1 Physics introduction: D-instanton corrections to HM moduli spaces
- 2 Math introduction: indefinite theta series**
- 3 Indefinite theta series and generalized error functions
- 4 Back to physics: S-duality in HM moduli space

- Theta series for Euclidean lattices are an important source of holomorphic modular forms: for $Q(x) = B(x, x)$ a positive definite form on \mathbb{R}^n ,

$$\Theta_Q(\tau, \nu) = \sum_{k \in \mathbb{Z}^n} q^{\frac{1}{2}Q(k)} e^{2\pi i B(\nu, k)}, \quad q = e^{2\pi i \tau}$$

is a holomorphic Jacobi form of weight $n/2$ for a suitable congruence subgroup of $SL(2, \mathbb{Z})$.

- Let $Q(x)$ is a signature $(r, n - r)$ quadratic form and \mathcal{C} an open cone in $\mathbb{R}^{n-r, r}$ such that $x \in \mathcal{C} \Rightarrow Q(x) > 0$, then

$$\Theta_{Q, \mathcal{C}}(\tau, \nu) = \sum_{k \in \mathbb{Z}^n \cap \mathcal{C}} q^{\frac{1}{2}Q(k)} e^{2\pi i B(\nu, k)}$$

defines a holomorphic q -series, but is it modular ?

- Such indefinite theta series occur in many branches of maths and physics:
 - Partition functions of coset models / branching functions for affine Lie algebras: *Kac Peterson 1984*
 - Characters of superconformal field theories / super-Lie algebras: *Eguchi Taormina 1988, Kac Wakimoto 2000, Semikhatov Taormina Tipunin 2003*
 - Donaldson and Vafa-Witten invariants of 4-manifolds: (*Zagier 1991*), *Vafa Witten 1994, Goettsche 1996, Goettsche-Zagier 1998, Manschot 2010*
 - Quantum invariants of knots and 3-manifolds: *Lawrence Zagier 1999, Hikami 2007, Hikami Lovejoy 2014*
 - Combinatorics of partitions: *Bringmann Ono 2005*
 - Mirror symmetry for elliptic curves and Abelian varieties: *Polischchuk 1998*
 - Gromov-Witten invariants of Landau-Ginzburg orbifolds: *Lau Zhou 2014, Bringmann Rolin Zwegers 2015*

- Examples of modular theta series of signature $(1, 1)$ were studied by Kronecker and Hecke in 1925.
- Non-modular examples were studied by Appell (1886) and Lerch (1892): the sum can be written as a signature $(1, 1)$ theta series, e.g. for $|q| < |y| < 1$, $y = e^{2\pi i v}$,

$$\begin{aligned}\theta_1(\tau, \nu) \mu(\tau, \nu) &\equiv \sum_{n \in \mathbb{Z}} (-1)^n \frac{q^{\frac{1}{2}n(n+1)} y^{n+\frac{1}{2}}}{1 - yq^n} \\ &= \left(\sum_{n, m \geq 0} - \sum_{n < 0, m \leq 0} \right) (-1)^n q^{\frac{1}{2}n(n+1) + mn} y^{n+m+\frac{1}{2}}\end{aligned}$$

This Appell-Lerch sum and generalizations thereof appear in many of examples mentioned earlier.

- In his ground-breaking PhD thesis (2002), Zwegers showed how to correct $\mu(\tau, \nu)$ into a **non-holomorphic, real-analytic, Jacobi form** of weight 1. Schematically, one replaces

$$\frac{1}{2} [\text{sign}(m) + \text{sign}(n)] \mapsto \frac{1}{2} \left[E_1(m\sqrt{2\tau_2}) + \text{sign}(n) \right]$$

in the summand, where $E_1(u) \equiv \text{Erf}(u\sqrt{\pi})$ is the error function.

- The difference is a theta series with an insertion of $M_1(m\sqrt{2\tau_2})$, where M_1 is the complementary error function,

$$M_1(u) = E_1(u) - \text{sign}(u) = -\text{sign}(u)\text{Erfc}(|u|\sqrt{\pi}),$$

which can be written as an **Eichler integral** of an ordinary unary theta series. From this the modular behavior of $\mu(\tau, \nu)$ follows.

- More generally, Zagier showed that the modular completion of an indefinite theta series of signature $(n - 1, 1)$ of the form

$$\Theta_{C,C'}(\tau, v) = \frac{1}{2} \sum_{k \in \mathbb{Z}^{n+b}} [\text{sign}B(k, C) - \text{sign}B(k, C')] q^{\frac{1}{2}Q(k)} e^{-2\pi i B(c, k - \frac{1}{2}b)}$$

where C, C' are a pair of vectors with $Q(C), Q(C'), B(C, C') < 0$ and $v = b\tau - c$, is obtained by replacing

$$\text{sign}B(k, C) \mapsto E_1 \left(B(k, C) \sqrt{\frac{2\tau_2}{-Q(C)}} \right)$$

- Our goal will be to generalize Zagier's construction to arbitrary signature $(n - r, r)$. Just to confuse you, I will henceforth flip the sign of Q : $(n - r, r) \rightarrow (r, n - r)$.

- 1 Physics introduction: D-instanton corrections to HM moduli spaces
- 2 Math introduction: indefinite theta series
- 3 Indefinite theta series and generalized error functions**
- 4 Back to physics: S-duality in HM moduli space

Vignéras' Theorem (1977) I

- Let $\Lambda \subset \mathbb{R}^n$ be an n -dimensional lattice with a signature $(r, n - r)$ quadratic form $Q(x) = B(x, x)$, such that $Q(k) \in 2\mathbb{Z}$ for $k \in \Lambda$. For any $\mu \in \Lambda^*/\Lambda$, $\lambda \in \mathbb{Z}$, $\Phi : \mathbb{R}^n \rightarrow \mathbb{C}$ such that $\Phi(x)e^{\frac{\pi}{2}Q(x)} \in L_1(\mathbb{R}^n)$,

$$\vartheta_\mu[\Phi, \lambda](\tau, b, c) = \tau_2^{-\lambda/2} \sum_{k \in \Lambda + \mu + b} \Phi(\sqrt{2\tau_2}k) q^{-\frac{1}{2}Q(k)} e^{2\pi i B(c, k - \frac{1}{2}b)}$$

satisfies the quasi-periodicity conditions

$$\begin{aligned}\vartheta_\mu[\Phi, \lambda](\tau, b, c) &= e^{i\pi B(c, k)} \vartheta_\mu[\Phi, \lambda](\tau, b + k, c) \\ &= e^{i\pi B(b, k)} \vartheta_\mu[\Phi, \lambda](\tau, b, c + k) \\ &= e^{i\pi Q(\mu)} \vartheta_\mu[\Phi, \lambda](\tau + 1, b, c + b)\end{aligned}$$

Vignéras' Theorem (1977) II

- If in addition Φ satisfies

$$\left[B^{-1}(\partial_x, \partial_x) + 2\pi x \partial_x \right] \Phi(x) = 2\pi \lambda \Phi(x),$$

$$R(x)\Phi(x)e^{\frac{\pi}{2}Q(x)} \in L^2(\mathbb{R}^n) \cap L^1(\mathbb{R}^n)$$

$$R(\partial)\Phi(x)e^{\frac{\pi}{2}Q(x)} \in L^2(\mathbb{R}^n) \cap L^1(\mathbb{R}^n)$$

for any quadratic polynomial R , then $\vartheta_\mu[\Phi, \lambda](\tau, b, c)$ transforms as a vector-valued Jacobi form of weight $(\lambda + \frac{n}{2}, 0)$. Namely,

$$\vartheta_\mu[\Phi, \lambda] \left(-\frac{1}{\tau}, c, -b \right) = \frac{(-i\tau)^{\lambda + \frac{n}{2}}}{\sqrt{|\Lambda^*/\Lambda|}} \sum_{\nu \in \Lambda^*/\Lambda} e^{2\pi i B(\mu, \nu)} \vartheta_\nu[\Phi, \lambda](\tau, b, c)$$

Vignéras' Theorem (1977) III

- Remark 1: The transformations are those of a Jacobi theta series with $\nu = 0$ and characteristics (b, c) and $c \in \Lambda \otimes \mathbb{R}$. To obtain the usual Jacobi form, set $\vartheta_\mu(\tau, b, c) = e^{i\pi B(b, b\tau - c)} \tilde{\vartheta}_\mu(\tau, \nu = b\tau - c)$.
- Remark 2: Under the Maass raising and lowering operators,

$$\tau_2^2 \partial_{\bar{\tau}} \vartheta_\mu[\Phi, \lambda] = \vartheta_\mu \left[\frac{i}{4} (x \partial_x \Phi - \lambda \Phi), \lambda - 2 \right],$$
$$\left(\partial_\tau - \frac{i(\lambda + \frac{n}{2})}{2\tau_2} \right) \vartheta_\mu[\Phi, \lambda] = \vartheta_\mu \left[-\frac{i}{4} (x \partial_x \Phi + [\lambda + n + 2\pi Q(x)] \Phi), \lambda + 2 \right].$$

We call $\tau_2^2 \partial_{\bar{\tau}}$ and $x \partial_x - \lambda$ the 'shadow' operators.

- Remark 3: For any r -dimensional time-like plane $\mathcal{P} \subset \mathbb{R}^n$ and $x \in \mathbb{R}^n$, let $x = x_+ + x_-$ with $x_+ \in \mathcal{P}$, $x_- \in \mathcal{P}^\perp$. The function $\Phi(x) = e^{-\pi Q(x_+)}$ satisfies the assumptions of the theorem with $\lambda = -r$, and leads to the usual Siegel-Narain theta series, also known as Narain lattice partition function $\Gamma_{r,n-r}$.
- Remark 4: In order to get a holomorphic theta series, one needs $x\partial_x\Phi = \lambda\Phi$, but a homogeneous polynomial of degree λ will not satisfy the assumptions of the theorem. Thus, there is a clear tension between holomorphy and modularity.
- To achieve mock modularity, we shall take Φ to be *locally* a polynomial of degree λ . For simplicity, take a locally constant homogeneous function ($\lambda = 0$).

Indefinite theta series of Lorentzian signature I

Let $Q(x)$ a signature $(1, n - 1)$ quadratic form, and C, C' linearly independent vectors with $Q(C) = Q(C') = 1$, $B(C, C') > 0$,

$$\Phi_1(x) = \frac{1}{2}(\text{sign}[B(C, x)] - \text{sign}[B(C', x)]) ,$$

$$\widehat{\Phi}_1(x) = \frac{1}{2}(E_1[B(C, x)] - E_1[B(C', x)])$$

$$\Psi_1(x) = \frac{i}{4}(B(C, x)e^{-\pi[B(C, x)]^2} - B(C', x)e^{-\pi[B(C', x)]^2})$$

- 1 $\Theta_\mu[\Phi_1, 0], \Theta_\mu[\widehat{\Phi}_1, 0], \Theta_\mu[\widehat{\Phi}_1, 0]$ are all convergent;
- 2 $\Theta_\mu[\widehat{\Phi}_1, 0]$ is real-analytic vector-valued Jacobi form of weight $n/2$;
- 3 $\Theta_\mu[\Phi_1, 0]$ is a holomorphic in τ and in z , but not modular;
- 4 Their difference is proportional to the Eichler integral of Ψ_1 ,


$$\Theta_\mu[\widehat{\Phi}_1 - \Phi_1, 0](\tau, b, c) = -4 \int_{-i\infty}^{\bar{\tau}} \frac{d\bar{w}}{(\tau - \bar{w})^2} \Theta_\mu[\Psi_1, -2](\tau, \bar{w}, b, c).$$

Indefinite theta series of Lorentzian signature II

Sketch of proof:

- 1 $\text{sign}[B(C, x)] = \text{sign}[B[(C, x)]$ unless $Q(x) < 0$;
- 2 $\Phi_1(x)$ is locally constant so $\Theta_\mu[\Phi_1, 0]$ is holomorphic;
- 3 $E_1(u)$ satisfies the 1D Vignéras equation,

$$(\partial_u^2 + 2\pi u \partial_u)E_1(u) = 0$$



so $\widehat{\Phi}_1$ satisfies the assumptions of Vignéras theorem;


- 4 Ψ_1 is proportional to the shadow of $\widehat{\Phi}_1$.

Remark: In the limit where C becomes null, then $E_1[B(C, x)] \rightarrow \text{sign}[B(C, x)]$. If both C and C' are null and $B(C, C') > 0$, then $\Theta_\mu[\Phi_1]$ is a **holomorphic** vector-valued Jacobi form of weight $n/2$.


Indefinite theta series of Lorentzian signature III

To prepare the ground for higher signature case, note the following integral representations:

$$E_1(u) = \int_{\mathbb{R}} du' \operatorname{sign}(u') e^{-\pi(u-u')^2}$$



$$M_1(u) = \frac{i}{\pi} \int_{\mathbb{R}-iu} \frac{dz}{z} e^{-\pi z^2 - 2\pi izu}$$



which make it clear that M_1, E_1 are solutions to 1D Vignéras equation,

$$[\partial_u^2 + 2\pi u \partial_u] M_1 = \frac{i}{\pi} \int_{\mathbb{R}-iu} \frac{dz}{z} 2\pi z \partial_z [e^{-\pi z^2 - 2\pi izu}] = 0$$

$$[\partial_u^2 + 2\pi u \partial_u] E_1 = \int_{\mathbb{R}} du' \operatorname{sign}(u') [\partial_u \partial_{u'} + 2\pi u \partial_{u'}] e^{-\pi(u-u')^2} = 0$$

Indefinite theta series of Lorentzian signature IV

Indefinite theta series of signature $(r, n - r)$ I

- In signature $(r, n - r)$, consider the locally constant function

$$\Phi_r(x) = \frac{1}{2^r} \prod_{i=1}^r (\text{sign}[B(C_i, x)] - \text{sign}[B(C'_i, x)])$$

where C_i, C'_i are chosen such that $Q(x) < 0$ whenever $\Phi_r(x) \neq 0$. To find its modular completion, we need a C^∞ solution $\widehat{\Phi}_r$ of Vignéras equation which asymptotes to Φ_r as $|x| \rightarrow \infty$.

- For $r = 2$, Alexandrov Banerjee Manschot BP (2016) found sufficient conditions for the convergence of $\vartheta[\Phi_r, 0]$. Kudla (2016) gave weaker conditions which work for arbitrary r .

Indefinite theta series of signature $(r, n - r)$ II

- To state Kudla's conditions, note that the space \mathcal{D} of positive oriented r planes in $\mathbb{R}^{r, n-r}$ has two disconnected components \mathcal{D}^\pm . For any $I \subset \{1, \dots, r\}$, let $C_I = (C_1'', \dots, C_r'')$ where $C_i'' = C_i$ if $i \in I$, and $C_i'' = C_i'$ if $i \notin I$.

Theorem (Kudla 2016): Assume that all such collections C_I span **distinct positive oriented r planes in the same component**, say \mathcal{D}^\pm . Then $\vartheta[\Phi_r, 0]$ is absolutely convergent.

- For $r = 2$, this reduces to

$$\begin{aligned} Q(C_1), Q(C_1'), Q(C_2), Q(C_2') &> 0 \\ \Delta_{12}, \Delta_{12'}, \Delta_{1'2}, \Delta_{1'2'} &> 0 \\ B(C_{1\perp 2}, C_{1'\perp 2}), B(C_{1\perp 2'}, C_{1'\perp 2'}) &> 0 \\ B(C_{2\perp 1}, C_{2'\perp 1}), B(C_{2\perp 1'}, C_{2'\perp 1'}) &> 0 \end{aligned}$$

where $\Delta_{12} = Q(C_1)Q(C_2) - B(C_1, C_2)^2$, $C_{1\perp 2} = C_1 - \frac{B(C_1, C_2)}{Q(C_2)}C_2$,
etc

Indefinite theta series of signature $(r, n - r)$ III



- To construct $\widehat{\Phi}_r$, consider the natural generalizations of the error functions $M_1(u)$ and $E_1(u)$,

$$E_r(\{C_i\}; x) = \int_{\langle C_1, \dots, C_r \rangle} d^r y \prod_{i=1}^r \text{sign} B(C_i, y) e^{-\pi Q(y-x_+)},$$

$$M_r(\{C_i\}; x) = \left(\frac{i}{\pi}\right)^r \int_{\langle C_1, \dots, C_r \rangle - iX} d^r z \frac{\sqrt{\Delta(\{C_i\})} e^{-\pi Q(z) - 2\pi i B(z, x)}}{\prod_{i=1}^r B(C_i, z)}$$

where $d^r y$ is the uniform measure on the plane $\langle C_1, \dots, C_r \rangle$, normalized such that $\int_{\langle C_1, \dots, C_r \rangle} d^r y e^{-\pi Q(y)} = 1$, and x_+ is the orthogonal projection of x on the same plane.

Indefinite theta series of signature $(r, n - r)$ IV



Indefinite theta series of signature $(r, n - r)$ V

Proposition: (ABMP 2016; Nazaroglu 2016)

- Prop: $E_r(\{C_i\}; x)$ is a C^∞ solution of Vignéras' equation with $\lambda = 0$, which asymptotes to $\prod_{i=1}^r \text{sign} B(C_i, x)$ as $|x| \rightarrow \infty$.
- $M_r(\{C_i\}; x)$ is a C^∞ solution of Vignéras' equation with $\lambda = 0$, away from the walls $B(C_i, x) = 0$, exponentially suppressed in all directions
- The difference $E_r(\{C_i\}) - \prod_{i=1}^r \text{sign} B(C_i, x)$ is a linear combination of $M_{r'}$ functions with $1 \leq r' \leq r$, with locally constant coefficients.
- The shadow of $E_r(\{C_i\})$ is a linear combination of $E_{r'}$ functions with $0 \leq r' < r$, with Gaussian coefficients.

Indefinite theta series of signature $(r, n - r)$ VI

Theorem:

- The modular completion of $\vartheta[\Phi_r, 0]$ is the non-holomorphic theta series $\vartheta[\widehat{\Phi}_r, 0]$ with kernel

$$\widehat{\Phi}_r(x) = \frac{1}{2^r} \sum_{I \subset \{1, \dots, r\}} (-1)^{r-|I|} E_r(C_I; x)$$

- Its shadow Ψ_r is a linear combination of indefinite theta series of signature $(r - 1, n - r + 1)$.
- The difference $\vartheta[\widehat{\Phi}_r - \Phi_r, 0]$ is an Eichler integral of Ψ_r , giving access to the modular properties of the holomorphic theta series $\vartheta[\Phi_r, 0]$.

Other approaches to indefinite theta series I

- An alternative approach to indefinite theta series was developed in the 80s by Kudla and Millson, who constructed a closed (r, r) -form $\theta_r^{KM}(z, \tau)$ on $\mathcal{D} \times \mathcal{H}$, which is invariant under a (finite index subgroup) of $\text{Aut}(\Lambda)$ and has modular weight $n/2$ in τ .
- Kudla (2016) proposes (and shows for $r = 1$ and $r = 2$) that

$$\vartheta[\widehat{\Phi}_r, 0] \propto \int_S \theta_r^{KM}(z, \tau)$$

where S is a geodesic hypercube in \mathcal{D}^+ , parametrized by

$$\phi : [0, 1]^r \rightarrow S \subset \mathcal{D}, [s_1, \dots, s_r] \mapsto \langle B_1(s_1), \dots, B_r(s_r) \rangle$$

where $B_i(s_i) = (1 - s_i)C_i + s_iC_i$.

Other approaches to indefinite theta series II

- The shadow is proportional to the integral of $\theta_{r-1}^{KM}(z, \tau)$ on the faces of the cube, hence a linear combination of indefinite theta series of signature $(r-1, n-r+1)$.
- Any compact geodesic polyhedron S in \mathcal{D}^+ similarly leads to a non-holomorphic theta series, which is the modular completion of a holomorphic indefinite theta series.
- Any such S can be decomposed as a sum of geodesic simplexes. Each term corresponds to a theta series $\Theta_{Q, \mathcal{C}}$ where \mathcal{C} is a tetrahedral cone.

Westerholt-Raum

- 1 Physics introduction: D-instanton corrections to HM moduli spaces
- 2 Math introduction: indefinite theta series
- 3 Indefinite theta series and generalized error functions
- 4 Back to physics: S-duality in HM moduli space**

S-duality in HM moduli space I

- A canonical system of Darboux coordinates Ξ on \mathcal{Z} is obtained by solving the integral equations

$$\Xi = \Xi_{\text{sf}} + \frac{1}{8\pi^2} \sum_{\gamma} \gamma \Omega(\gamma, z^a) \int_{\ell_{\gamma}} \frac{dt'}{t'} \frac{t+t'}{t-t'} \log \left[1 - e^{-2\pi i \langle \Xi(t'), \gamma \rangle} \right],$$

where Ξ_{sf} are the semi-flat coordinates and

$\ell(\gamma) = \{t : Z(\gamma, z^a)/t \in i\mathbb{R}^+\}$ are the BPS rays and $\Omega(\gamma, z^a)$ the generalized DT-invariants.

- Using the wall-crossing formula, DT invariants can be expressed in terms of MSW invariants, defined as the DT invariants in the ‘large volume attractor chamber’:

$$\Omega^{\text{MSW}}(p^a, q_a, q_0) = \lim_{\lambda \rightarrow +\infty} \Omega \left(0, p^a, q_a, q_0; -\kappa^{ab} q_b + i\lambda p^a \right)$$

S-duality in HM moduli space II

- Solving the integral system iteratively and expressing Ω in terms of Ω^{MSW} leads to an infinite 'multi-instanton sum'.
- In the large volume limit, zooming near $t = 0$, one finds that at one-instanton order, the Darboux coordinates Ξ can be expressed in terms of contour integrals of the form

$$\mathcal{J}_\rho(t) = \sum_{\substack{q \in \Lambda \\ q_0 \in \mathbb{Z}}} \int_{\mathbb{R}} \frac{dt'}{t' - t} \Omega^{\text{MSW}}(\gamma) e^{-2\pi i \langle \Xi_{\text{sf}}(t'), \gamma \rangle},$$

- Using spectral flow invariance of $\Omega^{\text{MSW}}(\gamma)$, and restricting to $t = 0$ for simplicity, this can be rewritten as

$$\mathcal{J}_\rho(0) = \sum_{\mu \in \Lambda^* / \Lambda} h_{\rho, \mu} \left[\sum_{k \in \Lambda + \mu} M_1(k + \sqrt{\tau_2}) q^{\frac{1}{2}(k, k)} \right]$$

S-duality in HM moduli space III

- The series $\sum_k M_1(k_+ \sqrt{\tau_2}) q^{\frac{1}{2}(k,k)}$ is an Eichler integral of the Gaussian theta series $\sum_k k_+ e^{-\pi \tau_2 k_+^2} q^{\frac{1}{2}(k,k)}$, therefore it transforms non-homogeneously under $SL(2, \mathbb{Z})$.
- However, since it appears in the modular completion of the Zagier-type indefinite theta series $\sum_k [\text{sign}(k_+) - \text{sign}(k'_+)] q^{\frac{1}{2}(k,k)}$ (where $k'_+ = k_a t'^a$ with $Q(t') = 0$), its modular anomaly is holomorphic, and therefore Ξ transforms by a holomorphic contact transformation !
- This shows that at one-instanton level, D3-instanton corrections are consistent with S-duality, provided the generating function $h_{\rho, \mu}$ of MSW invariants is a vector-valued modular form of fixed weight and multiplier system, as predicted by MSW.

Alexandrov Manschot BP, 2012

S-duality in HM moduli space IV

- At two-instanton level, one has to deal with indefinite theta series of signature $(2, 2b_2 - 2)$, corresponding to sums over fluxes on pairs of D3-branes wrapped on a divisor $\mathcal{P} = \mathcal{P}_1 + \mathcal{P}_2$.
- After a lot of work, one finds that D3-instanton corrections are consistent with S-duality, provided $h_{p,\mu}$ is a vector-valued **mock** modular form of fixed weight, multiplier system and shadow. More precisely, $\widehat{h}_{p,\mu} \equiv h_{p,\mu} - R_{p,\mu}$ is a vector-valued modular form of weight $-\frac{b_2}{2} - 1$, where $R_{p,\mu}$ is a non-holomorphic function of τ constructed from $h_{p_1,\mu_1}, h_{p_2,\mu_2}$.

Alexandrov Banerjee Manschot BP, 2016

- Such mock modularity is known to appear in the context of rank 2 sheaves on complex surfaces, here we see it arise more generally in the context of DT invariants of pure dimension 2 sheaves on Calabi-Yau threefolds.