

Rankin-Selberg methods for String Amplitudes

Boris Pioline

CERN & LPTHE



Math/Phys seminar, Köln
May 10, 2012

*based on work with C. Angelantonj and I. Florakis,
arXiv:1110.5318,1203.0566*

- In string theory, an interesting class of terms (often known as BPS-saturated coupling, topological amplitude or F-term) in the low energy effective action are given by a modular integral

$$\mathcal{A} = \int_{\mathcal{F}} d\mu \Gamma_{(d+k,d)} \Phi(\tau)$$


- $\mathcal{F} = \Gamma \backslash \mathcal{H}$: fundamental domain of the modular group $\Gamma = SL(2, \mathbb{Z})$ on the Poincaré UHP \mathcal{H} ;
- $d\mu = d\tau_1 d\tau_2 / \tau_2^2$ is the Γ -invariant measure;
- $\Gamma_{(d+k,d)} = \tau_2^{d/2} \sum q^{\frac{1}{2}p_L^2} \bar{q}^{\frac{1}{2}p_R^2}$: a theta series for an even self-dual lattice of signature $(d+k, d)$, known as the Narain lattice partition function;
- $\Phi(\tau)$: an (almost, weak) holomorphic modular form of weight $w = -k/2$, known as the elliptic genus

Modular integrals and BPS amplitudes II

- Such amplitudes arise in a variety of examples:
 - Gauge thresholds, $R^2 F^{2h-2}$ in $\text{Het}/K3 \times T^2$ at one-loop
Dixon Kaplunovsky Louis; Harvey Moore
 - F^4 couplings in Het/T^d at one-loop
Bachas Fabre Kiritsis Obers Vanhove
 - R^4 couplings in type II/ T^d at one-loop ($\Phi = 1$)
Green Vanhove; Kiritsis BP
 - R^2 couplings in type II/ $K3 \times T^2$ at one-loop ("")
Harvey Moore; Gregori Kiritsis Kounnas Obers Petropoulos BP
 - F^4 couplings in type II/ T^4/\mathbb{Z}_N at tree-level ("")
Obers BP
 - $\nabla^4 R^4$ couplings in M/T^d at two-loops ("")
Green Vanhove Russo
- These amplitudes are strongly constrained by supersymmetry, and offer precise tests of string dualities.

Modular integrals and BPS amplitudes III

- When \mathcal{A} arises at one-loop, and upon choosing \mathcal{F} as the standard 'keyhole' domain, τ_2 can be interpreted as the **Schwinger parameter**, while τ_1 is a **Lagrange multiplier** enforcing the level-matching constraint $p_L^2 - p_R^2 = N$.



Theta correspondances

- From the mathematical point of view, modular integrals give a **theta correspondence**

$$\Phi : \Gamma \backslash \mathcal{H} \rightarrow \mathbb{C} \quad \leftrightarrow \quad \mathcal{A} : O(\Gamma_{d+k,d}) \backslash G_{d+k,d} \rightarrow \mathbb{C}$$

between modular forms on \mathcal{H} and automorphic forms on the Grassmannian $G_{d+k,d}$, or Narain moduli space

$$G_{d+k,d} = \frac{O(d+k, d)}{O(d+k) \times O(d)} \ni (g_{ij}, B_{ij}, Y_i^a)$$

Theta correspondances

- From the mathematical point of view, modular integrals give a **theta correspondence**

$$\Phi : \Gamma \backslash \mathcal{H} \rightarrow \mathbb{C} \quad \leftrightarrow \quad \mathcal{A} : O(\Gamma_{d+k,d}) \backslash G_{d+k,d} \rightarrow \mathbb{C}$$

between modular forms on \mathcal{H} and automorphic forms on the Grassmannian $G_{d+k,d}$, or Narain moduli space

$$G_{d+k,d} = \frac{O(d+k, d)}{O(d+k) \times O(d)} \ni (g_{ij}, B_{ij}, Y_i^a)$$

- Indeed, $SL(2) \times O(d+k, d)$ forms a dual pair in $Sp(d+k, d)$, and the lattice partition function is invariant under $\Gamma \times O(\Gamma_{d+k,d})$.

Theta correspondances

- From the mathematical point of view, modular integrals give a **theta correspondence**

$$\Phi : \Gamma \backslash \mathcal{H} \rightarrow \mathbb{C} \quad \leftrightarrow \quad \mathcal{A} : O(\Gamma_{d+k,d}) \backslash G_{d+k,d} \rightarrow \mathbb{C}$$

between modular forms on \mathcal{H} and automorphic forms on the Grassmannian $G_{d+k,d}$, or Narain moduli space

$$G_{d+k,d} = \frac{O(d+k, d)}{O(d+k) \times O(d)} \ni (g_{ij}, B_{ij}, Y_i^a)$$

- Indeed, $SL(2) \times O(d+k, d)$ forms a dual pair in $Sp(d+k, d)$, and the lattice partition function is invariant under $\Gamma \times O(\Gamma_{d+k,d})$.
- Theta correspondences are one of the few general ways (together with Langlands-Eisenstein series) to construct automorphic forms, and are central in the Langlands programme

Unfolding trick

- In the physics literature, the time-honored way to evaluate such integrals has been the **unfolding trick** or **orbit method** where the domain of integration \mathcal{F} is unfolded by grouping the terms in the theta series into orbits.

Unfolding trick

- In the physics literature, the time-honored way to evaluate such integrals has been the **unfolding trick** or **orbit method** where the domain of integration \mathcal{F} is unfolded by grouping the terms in the theta series into orbits.
- E.g for $d = 1$, representing $\Gamma_{(1,1)} = R \sum_{m,n} e^{-\pi R^2 |m-n\tau|^2 / \tau_2}$,

$$\int_{\mathcal{F}} \Gamma_{(1,1)} = \frac{\pi}{3} R + \frac{\pi}{3} R^{-1}$$

Unfolding trick

- In the physics literature, the time-honored way to evaluate such integrals has been the **unfolding trick** or **orbit method** where the domain of integration \mathcal{F} is unfolded by grouping the terms in the theta series into orbits.
- E.g for $d = 1$, representing $\Gamma_{(1,1)} = R \sum_{m,n} e^{-\pi R^2 |m - n\tau|^2 / \tau_2}$,

$$\int_{\mathcal{F}} \Gamma_{(1,1)} = \frac{\pi}{3} R + \frac{\pi}{3} R^{-1}$$

- For $d = 2$, a landmark (lengthy) computation shows

$$\int_{\mathcal{F}} (\Gamma_{(2,2)}(T, U) - \tau_2) d\mu = -\log \left(\frac{8\pi e^{1-\gamma}}{3\sqrt{3}} T_2 U_2 |\eta(T)\eta(U)|^4 \right)$$

Dixon Kaplunovsky Louis

where T, U parametrize the Grassmannian $G_{2,2} = \mathcal{H}_T \times \mathcal{H}_U / \mathbb{Z}_2$.

Rankin-Selberg method I

- The unfolding trick is also at the basis of the **Rankin-Selberg method** in analytic number theory: let

$$\begin{aligned} E^*(\tau; s) &\equiv \zeta^*(2s) \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} [\operatorname{Im}(\gamma \cdot \tau)]^s \\ &= \frac{1}{2} \zeta^*(2s) \sum_{(c,d)=1} \frac{\tau_2^s}{|c\tau + d|^{2s}} \end{aligned}$$

be the completed **non-holomorphic Eisenstein series**, where $\zeta^*(s) \equiv \pi^{-s/2} \Gamma(s/2) \zeta(s) = \zeta^*(1-s)$ is the completed zeta function with simple poles at $s = 1, 0$

- $E^*(\tau; s) = E^*(\tau; 1-s)$ is analytic in s away from $s = 0, 1$,

$$E^*(\tau; s) = \frac{1}{2(s-1)} + \frac{1}{2} \left(\gamma - \log(4\pi \tau_2 |\eta(\tau)|^4) \right) + \mathcal{O}(s-1),$$

- Let $F(\tau)$ a modular function of rapid decay at the cusp and consider the Rankin-Selberg transform

$$\mathcal{R}^*(F, s) \equiv \int_{\mathcal{F}} d\mu E^*(\tau; s) F(\tau)$$

- By the unfolding trick, $\mathcal{R}^*(F, s)$ is proportional to the **Mellin transform of the constant term** $F_0(\tau_2) = \int_{-1/2}^{1/2} d\tau_1 F(\tau)$,

$$\begin{aligned} \mathcal{R}^*(F; s) &= \zeta^*(2s) \int_{\mathcal{S}} \frac{d\tau_1 d\tau_2}{\tau_2^{2-s}} F(\tau) \\ &= \zeta^*(2s) \int_0^\infty d\tau_2 \tau_2^{s-2} F_0(\tau_2), \end{aligned}$$

- The analyticity and functional relation for E^* implies similar properties for $\mathcal{R}^*(F; s)$. For $F = f.g$ product of two cusp forms, this is used e.g. to show the analyticity and functional relation of the L-function $L(s) = \sum_n a_n b_n n^{-s} \propto \mathcal{R}^*(F; s)$.
- For us, the main point is that, since the residue of E^* at $s = 0, 1$ is constant, the residue of $\mathcal{R}^*(F; s)$ at $s = 0$ is proportional to the modular integral of F ,

$$\text{Res } \mathcal{R}^*(F; s)|_{s=1} = \frac{1}{2} \int_{\mathcal{F}} d\mu F = -\text{Res } \mathcal{R}^*(F; s)|_{s=0} .$$

Rankin-Selberg-Zagier method I

- This was extended by Zagier to the case where F is of moderate growth $F(\tau) \sim \phi(\tau_2)$ at the cusp ($\phi(\tau_2)$ at most a power): the renormalized modular integral

$$\text{R.N.} \int_{\mathcal{F}} d\mu F(\tau) \equiv \lim_{\mathcal{T} \rightarrow \infty} \left[\int_{\mathcal{F}_{\mathcal{T}}} d\mu F(\tau) - \hat{\varphi}(\mathcal{T}) \right]$$

is related to the Mellin transform of the (regularized) constant term

$$\mathcal{R}^*(F; s) = \zeta^*(2s) \int_0^{\infty} d\tau_2 \tau_2^{s-2} (F_0 - \varphi) ,$$

via

$$\text{R.N.} \int_{\mathcal{F}} d\mu F(\tau) = 2 \text{Res} [\mathcal{R}^*(F; s)]_{s=1} + \delta$$

- The correction δ depends only of the leading behavior $\phi(\tau_2)$, and is given by

$$\delta = 2 \operatorname{Res} [\zeta^*(2s) h_{\mathcal{T}}(s) + \zeta^*(2s - 1) h_{\mathcal{T}}(1 - s)]_{s=1} - \hat{\phi}(\mathcal{T}),$$

where

$$h_{\mathcal{T}}(s) = \int_0^{\mathcal{T}} d\tau_2 \varphi(\tau_2) \tau_2^{s-2}, \quad \hat{\phi}(\mathcal{T}) = \operatorname{Res} \left[\frac{h_{\mathcal{T}}(s)}{s-1} \right]_{s=1}$$

- Other renormalization schemes may give a different constant δ

Epstein series from modular integrals

- The RSZ method applies immediately to modular integrals with $\Phi = 1$:

$$\begin{aligned}\mathcal{R}^*(\Gamma_{(d,d)}; \mathbf{s}) &= \text{R.N.} \int_{\mathcal{F}} d\mu \tau_2^{d/2} \sum_{m_i, n^i} q^{\frac{1}{2}p_L^2} \bar{q}^{\frac{1}{2}p_R^2} E^*(\mathbf{s}, \tau), \\ &= \zeta^*(2\mathbf{s}) \int_0^\infty d\tau_2 \tau_2^{s+d/2-2} \sum_{m_i n^i=0} e^{-\pi\tau_2 \mathcal{M}^2} \\ &= \zeta^*(2\mathbf{s}) \frac{\Gamma(\mathbf{s} + \frac{d}{2} - 1)}{\pi^{s+\frac{d}{2}-1}} \mathcal{E}_V^d(\mathbf{g}, \mathbf{B}; \mathbf{s} + \frac{d}{2} - 1) \\ &\equiv \mathcal{E}_V^{d*}(\mathbf{g}, \mathbf{B}; \mathbf{s} + \frac{d}{2} - 1)\end{aligned}$$

where $\mathcal{E}_V^d(\mathbf{g}, \mathbf{B}; \mathbf{s})$ is the **constrained Epstein Zeta series**

$$\mathcal{E}_V^d(\mathbf{g}, \mathbf{B}; \mathbf{s}) \equiv \sum_{\substack{(m_i, n^i) \in \mathbb{Z}^{2d} \setminus (0,0) \\ m_i n^i = 0}} \mathcal{M}^{-2s}, \quad \mathcal{M}^2 = p_L^2 + p_R^2$$

Epstein series and BPS state sums I

- This is identified as a **sum over all BPS states** of momentum m_j and winding n^j along the torus, $O(\Gamma_{d,d})$ -invariant mass

$$\mathcal{M}^2 = (m_j + B_{ik}n^k)g^{ij}(m_j + B_{jl}n^l) + n^j g_{ij}n^i$$

subject to the $O(\Gamma_{d,d})$ -invariant **BPS condition** $m_j n^j = 0$.

- The constrained Epstein Zeta series $\mathcal{E}_V^d(g, B; s)$ converges absolutely for $s + \frac{d}{2} - 1 > 1$. The RSZ method shows that it admits a meromorphic continuation in the s -plane satisfying

$$\mathcal{E}_V^{d*}(g, B; s) = \mathcal{E}_V^{d*}(g, B; d - 1 - s),$$

with a simple pole at $s = 0, \frac{d}{2} - 1, \frac{d}{2}, 1$ (assume $d > 2$).

- The residue at $s = \frac{d}{2}$ produces the modular integral of interest:

$$\begin{aligned} \text{R.N. } \int_{\mathcal{F}} d\mu \Gamma_{(d,d)}(g, B) &= \frac{\pi}{3} \frac{\Gamma(d/2)}{\pi^{d/2}} \text{Res } \mathcal{E}_V^d(g, B; s) \Big|_{s=d/2} \\ &= \frac{\Gamma(d/2 - 1)}{\pi^{d/2-1}} \mathcal{E}_V^d(g, B; \frac{1}{2} d - 1) \end{aligned}$$

rigorously proving an old conjecture of Obers and myself.

- For $d = 1$ or $d = 2$:

$$\begin{aligned} \mathcal{E}_V^{1,*}(g, B; s - \frac{1}{2}) &= 2 \zeta^*(2s) \zeta^*(2s - 1) \left(R^{1-2s} + R^{2s-1} \right) \\ \mathcal{E}_V^{2,*}(T, U; s) &= 2 E^*(T; s) E^*(U; s) \end{aligned}$$

leading immediately to advertised results.

- The differential equations

$$0 = \left[\Delta_{\text{SO}(d,d)} - 2 \Delta_{\text{SL}(2)} + \frac{1}{4} d(d-2) \right] \Gamma_{(d,d)}(g, B)$$

$$0 = \left[\Delta_{\text{SL}(2)} - \frac{1}{2} s(s-1) \right] E^*(\tau; s),$$

imply that $\mathcal{E}_V^{d*}(s)$ is an eigenmode of the Laplace-Beltrami operator on the Grassmannian $G_{d,d}$ with eigenvalue $s(s-d+1)$, and more generally, of any $O(d, d)$ invariant differential operator.

Relation with other constructions

- The differential equations

$$0 = \left[\Delta_{\mathrm{SO}(d,d)} - 2 \Delta_{\mathrm{SL}(2)} + \frac{1}{4} d(d-2) \right] \Gamma_{(d,d)}(g, B)$$

$$0 = \left[\Delta_{\mathrm{SL}(2)} - \frac{1}{2} s(s-1) \right] E^*(\tau; s),$$

imply that $\mathcal{E}_V^{d*}(s)$ is an eigenmode of the Laplace-Beltrami operator on the Grassmannian $G_{d,d}$ with eigenvalue $s(s-d+1)$, and more generally, of any $O(d, d)$ invariant differential operator.

- $\mathcal{E}_V^{d*}(g, B; s)$ must be equal to the Langlands-Eisenstein series of $O(d, d)$ with infinitesimal character $\rho - 2s\alpha_1$, according to the Siegel-Weil formula.

Relation with other constructions

- The differential equations

$$0 = \left[\Delta_{SO(d,d)} - 2 \Delta_{SL(2)} + \frac{1}{4} d(d-2) \right] \Gamma_{(d,d)}(g, B)$$
$$0 = \left[\Delta_{SL(2)} - \frac{1}{2} s(s-1) \right] E^*(\tau; s),$$

imply that $\mathcal{E}_V^{d*}(s)$ is an eigenmode of the Laplace-Beltrami operator on the Grassmannian $G_{d,d}$ with eigenvalue $s(s-d+1)$, and more generally, of any $O(d, d)$ invariant differential operator.

- $\mathcal{E}_V^{d*}(g, B; s)$ must be equal to the Langlands-Eisenstein series of $O(d, d)$ with infinitesimal character $\rho - 2s\alpha_1$, according to the Siegel-Weil formula.
- The residue at $s = \frac{d}{2}$ is the minimal theta series, attached to the minimal representation of $SO(d, d)$ (functional dimension $2d - 3$).

Ginzburg Rallis Soudry; Kazhdan BP Waldron; Green Vanhove Miller

Modular integrals with unphysical tachyons I

- For many cases of interest, the integrand is NOT of moderate growth at the cusp, rather it grows exponentially, due to the heterotic unphysical tachyon.
- In mathematical terms, $\Phi(\tau) \in \mathbb{C}[\hat{E}_2, E_4, E_6, 1/\Delta]$ is a **weak almost holomorphic modular** form with weight $w = -k/2 \leq 0$.
- The RSZ method fails, however the unfolding trick could still work provided $\Phi(\tau)$ had a **uniformly convergent Poincaré representation**

$$\Phi(\tau) = \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} f(\tau)|_w \gamma$$

where the seed $f(\tau)$ is invariant under $\tau \rightarrow \tau + 1$ and

$$(f|_w \gamma)(\tau) = (c\tau + d)^{-w} f(\gamma \cdot \tau), \quad \gamma \cdot \tau = \frac{a\tau + b}{c\tau + d}.$$

Various Poincaré series representations I

- Naively, one requires $f(\tau) = 1/q^\kappa$ ($\kappa = 1$ for physics applications), however convergence requires $f(\tau) \ll \tau_2^{1-\frac{w}{2}}$ as $\tau_2 \rightarrow 0$. This is OK for $w > 2$ but fails for $w \leq 0$. We need to regularize.
- Any weak holomorphic modular form can be represented as a linear combination of **regularized holomorphic Poincaré series**

$$P(\kappa, w) = \frac{1}{2} \sum_{(c,d)=1}^{\dagger} (c\tau + d)^{-w} e^{-2\pi i \kappa \frac{a\tau+b}{c\tau+d}} R_w \left(\frac{2\pi i \kappa}{c(c\tau + d)} \right),$$

where $R_w(x) \sim x^{1-w}/\Gamma(2-w)$ as $x \rightarrow 0$ and approaches 1 as $x \rightarrow \infty$. However this is only conditionally convergent, and $P(\kappa, w)$ in general has **modular anomalies**.

Niebur; Knopp; Manschot Moore

Various Poincaré series representations II

- Another option is to insert a **non-holomorphic convergence factor** à la Hecke-Kronecker, i.e. choose $f(\tau) = \tau_2^{s-\frac{w}{2}} q^{-\kappa}$

$$E(s, \kappa, w) \equiv \frac{1}{2} \sum_{(c,d)=1} \frac{\tau_2^{s-\frac{w}{2}}}{|c\tau + d|^{2s-w}} (c\tau + d)^{-w} e^{-2\pi i \kappa \frac{a\tau+b}{c\tau+d}}$$

Selberg; Goldfeld Sarnak; Pribitkin

This converges absolutely for $\text{Re}(s) > 1$, but the analytic continuation to $s = \frac{w}{2}$ is tricky (no modular anomaly, but in general **holomorphic anomalies**).

- Moreover, $E(s, \kappa, w)$ is not an eigenmode of the Laplacian, rather $[\Delta_w + \frac{1}{2}s(1-s) + \frac{1}{8}w(w+2)] E(s, \kappa, w) = 2\pi\kappa (s - \frac{w}{2}) E(s+1, \kappa, w)$

Niebur-Poincaré series I

- There exist yet another regularization which does not require analytic continuation and is still an eigenmode of the Laplacian: the **Niebur-Poincaré series**

$$\mathcal{F}(s, \kappa, w) = \frac{1}{2} \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} \mathcal{M}_{s,w}(-\kappa\tau_2) e^{-2\pi i \kappa \tau_1} |w \gamma$$

Niebur; Hejhal; Bruinier Ono Bringmann...

where $\mathcal{M}_{s,w}(y)$ is proportional to a Whittaker function, so that

$$\left[\Delta_w + \frac{1}{2} s(1-s) + \frac{1}{8} w(w+2) \right] \mathcal{F}(s, \kappa, w) = 0$$

- The seed $f(\tau) = \mathcal{M}_{s,w}(-\kappa\tau_2) e^{-2\pi i \kappa \tau_1}$ satisfies

$$f(\tau) \sim_{\tau_2 \rightarrow 0} \tau_2^{\operatorname{Re}(s) - \frac{w}{2}} e^{-2\pi i \kappa \tau_1} \quad f(\tau) \sim_{\tau_2 \rightarrow \infty} \frac{\Gamma(2s)}{\Gamma(s + \frac{w}{2})} q^{-\kappa}$$

hence $\mathcal{F}(s, \kappa, w)$ converges absolutely for $\operatorname{Re}(s) > 1$.

Niebur-Poincaré series II

- For $s = 1 - \frac{w}{2}$ the eigenvalue coincides with that of a holomorphic modular form, and the seed simplifies to

$$f(\tau) = \Gamma(2 - w) \left(q^{-\kappa} - \bar{q}^{\kappa} \sum_{\ell=0}^{-w} \frac{(4\pi\kappa T_2)^\ell}{\ell!} \right)$$

- For $w < 0$, the value $s = 1 - \frac{w}{2}$ lies in the convergence domain. $\mathcal{F}(1 - \frac{w}{2}, \kappa, w)$ is in general NOT holomorphic, but rather a **weak harmonic Maass form**.
- For $s = \frac{w'}{2}$ and $w' > 0$, $\mathcal{F}(\frac{w'}{2}, \kappa, w')$ IS weakly holomorphic. For $w' = 2 - w$, it is the **Farey transform** (or the 'ghost') of the weak harmonic Maass form $\mathcal{F}(1 - \frac{w}{2}, \kappa, w)$.

Niebur-Poincaré series III

w	$\mathcal{F}(1 - \frac{w}{2}, 1, w)$	$\mathcal{F}(1 - \frac{w}{2}, 1, 2 - w)$
0	$j + 24$	$E_4^2 E_6 \Delta^{-1}$
-2	$3! E_4 E_6 \Delta^{-1}$	$E_4(j - 240)$
-4	$5! E_4^2 \Delta^{-1}$	$E_6(j + 204)$
-6	$7! E_6 \Delta^{-1}$	$E_4^2(j - 480)$
-8	$9! E_4 \Delta^{-1}$	$E_4 E_6(j + 264)$
-10	$11! \Phi_{-10}$	<i>(mess)</i>
-12	$13! \Delta^{-1}$	$E_4^2 E_6(j + 24)$
-14	$15! \Phi_{-14}$	<i>(mess)</i>

Niebur-Poincaré series IV

- Indeed, for $w = -10$, there does not exist any weak holomorphic modular form with a simple pole at the cusp. Rather, there exist a weak harmonic Maass form

$$\begin{aligned}\Phi_{-10} = & q^{-1} - \frac{65520}{691} q - 1842.89 q - 23274.08 q^2 + \dots \\ & + \sum_{m=1}^{\infty} m^{-11} \bar{b}_m \Gamma(11, 4\pi m\tau_2) q^{-m}\end{aligned}$$

Ono

with shadow $\sum b_m q^m$ proportional to the cusp form Δ .

- Theorem (Bruinier) : any weak holomorphic modular form of weight $w \leq 0$ with polar part $\Phi = \sum_{-\kappa \leq m < 0} a_m q^m + \mathcal{O}(1)$ can be represented as a **linear combination of Niebur-Poincaré series**

$$\Phi = \frac{1}{\Gamma(2-w)} \sum_{-\kappa \leq m < 0} a_m \mathcal{F}\left(1 - \frac{w}{2}, m, w\right) + a'_0 \delta_{w,0}$$

Niebur-Poincaré series V

- **Almost** weak holomorphic modular forms can be reached by **raising and lowering operators**

$$D_w = \frac{i}{\pi} \left(\partial_\tau - \frac{iw}{2\tau_2} \right), \quad \bar{D}_w = -i\pi \tau_2^2 \partial_{\bar{\tau}},$$


under which

$$D_w \cdot \mathcal{F}(s, \kappa, w) = 2\kappa \left(s + \frac{w}{2} \right) \mathcal{F}(s, \kappa, w + 2),$$
$$\bar{D}_w \cdot \mathcal{F}(s, \kappa, w) = \frac{1}{8\kappa} \left(s - \frac{w}{2} \right) \mathcal{F}(s, \kappa, w - 2).$$

The relevant values of s are $s = 1 - \frac{w}{2} + n$ with $n \geq 0$. E.g.

$$\frac{\hat{E}_2 E_4 E_6}{\Delta} = \mathcal{F}(2, 1, 0) - 5 \mathcal{F}(1, 1, 0) - 144$$

Niebur-Poincaré series VI



Unfolding the modular integral

- Using Bruinier's thm, any modular integral can be expressed as a linear combination of

$$\mathcal{I}_{d+k,d}(\mathbf{s}, \kappa;) = \text{R.N.} \int_{\mathcal{F}} d\mu \Gamma_{d+k,d}(G, B, Y) \mathcal{F}(\mathbf{s}, \kappa, -\frac{k}{2})$$

Unfolding the modular integral

- Using Bruinier's thm, any modular integral can be expressed as a linear combination of

$$\mathcal{I}_{d+k,d}(\mathbf{s}, \kappa;) = \text{R.N.} \int_{\mathcal{F}} d\mu \Gamma_{d+k,d}(G, B, Y) \mathcal{F}(\mathbf{s}, \kappa, -\frac{k}{2})$$

- Using the unfolding trick, one arrives at the **BPS state sum**

$$\begin{aligned} \mathcal{I}_{d+k,d}(\mathbf{s}, \kappa) &= (4\pi\kappa)^{1-\frac{d}{2}} \Gamma(\mathbf{s} + \frac{2d+k}{4} - 1) \\ &\times \sum_{\text{BPS}} {}_2F_1\left(\mathbf{s} - \frac{k}{4}, \mathbf{s} + \frac{2d+k}{4} - 1; 2\mathbf{s}; \frac{4\kappa}{p_L^2}\right) \left(\frac{p_L^2}{4\kappa}\right)^{1-\mathbf{s}-\frac{2d+k}{4}} \end{aligned}$$

Bruinier; Angelantonj Florakis BP

where $\sum_{\text{BPS}} \equiv \sum_{p_L, p_R} \delta(p_L^2 - p_R^2 - 4\kappa)$. This converges absolutely for $\text{Re}(\mathbf{s}) > \frac{2d+k}{4}$ and can be analytically continued to $\text{Re}(\mathbf{s}) > 1$ with a simple pole at $\mathbf{s} = \frac{2d+k}{4}$.

Unfolding the modular integral

- The result is manifestly **T-duality invariant**, and requires no choice of chamber in Narain modular space. Singularities on $G_{d+k,d}$ arise when $p_L^2 = 0$ for some lattice vector.
- For the relevant values $s = 1 - \frac{w}{2} + n$, the result can be written using elementary functions, e.g.

$$\mathcal{I}_{1,1}(1+n, \kappa) = \frac{1}{2} \sqrt{\pi} (16\kappa)^{1+n} \Gamma(n + \frac{1}{2}) \\ \times \sum_{\substack{p,q \in \mathbb{Z} \\ pq = \kappa}} \left(|pR + qR^{-1}| + |pR - qR^{-1}| \right)^{-1-2n}$$

$$\mathcal{I}_{2+k,2}(1 + \frac{k}{4}, \kappa) = -\Gamma(2 + \frac{k}{2}) \sum_{\text{BPS}} \left[\log \left(\frac{p_R^2}{p_L^2} \right) + \sum_{\ell=1}^{k/2} \frac{1}{\ell} \left(\frac{p_L^2}{4\kappa} \right)^{-\ell} \right]$$

One example

- Consider $\text{Het}/T^2 \times K3$ at \mathbb{Z}_2 orbifold point with gauge group broken to $E_8 \times E_7 \times SU(2)$. The gauge threshold for E_7 is

$$\Delta_{E_7} = -\frac{1}{12} \int_{\mathcal{F}} d\mu \Gamma_{2,2} \frac{\hat{E}_2 E_4 E_6 - E_4^3}{\Delta}$$

Expressing the elliptic genus as a linear combination

$$\frac{\hat{E}_2 E_4 E_6 - E_4^3}{\Delta} = \mathcal{F}(2, 1, 0) - 6 \mathcal{F}(1, 1, 0) - 864$$

one arrives at

$$\Delta_{E_7} = \sum_{\text{BPS}} \left[1 + \frac{p_R^2}{4} \log \left(\frac{p_R^2}{p_L^2} \right) \right] - 72 \log \left(4\pi e^{-\gamma} T_2 U_2 |\eta(T)\eta(U)|^4 \right)$$

Conclusion - Outlook

- Modular integrals can be efficiently computed using Rankin-Selberg type methods. While the time-honored 'orbit method' is still useful for studying large volume limits, our method makes T-duality and singularities manifest.

Conclusion - Outlook

- Modular integrals can be efficiently computed using Rankin-Selberg type methods. While the time-honored 'orbit method' is still useful for studying large volume limits, our method makes T-duality and singularities manifest.
- For application to orbifold models, it would be useful to extend this method to congruence subgroups of $SL(2, \mathbb{Z})$.

Conclusion - Outlook

- Modular integrals can be efficiently computed using Rankin-Selberg type methods. While the time-honored 'orbit method' is still useful for studying large volume limits, our method makes T-duality and singularities manifest.
- For application to orbifold models, it would be useful to extend this method to congruence subgroups of $SL(2, \mathbb{Z})$.
- The RSZ method also works at higher genus, at least for $g = 2, 3$ and $\Phi = 1$. It would be useful to derive Poincaré series representations for Siegel modular forms.

Conclusion - Outlook

- Modular integrals can be efficiently computed using Rankin-Selberg type methods. While the time-honored 'orbit method' is still useful for studying large volume limits, our method makes T-duality and singularities manifest.
- For application to orbifold models, it would be useful to extend this method to congruence subgroups of $SL(2, \mathbb{Z})$.
- The RSZ method also works at higher genus, at least for $g = 2, 3$ and $\Phi = 1$. It would be useful to derive Poincaré series representations for Siegel modular forms.
- The Niebur-Poincaré series may have useful applications for the Black Hole Farey Tale...

Conclusion - Outlook

- Modular integrals can be efficiently computed using Rankin-Selberg type methods. While the time-honored 'orbit method' is still useful for studying large volume limits, our method makes T-duality and singularities manifest.
- For application to orbifold models, it would be useful to extend this method to congruence subgroups of $SL(2, \mathbb{Z})$.
- The RSZ method also works at higher genus, at least for $g = 2, 3$ and $\Phi = 1$. It would be useful to derive Poincaré series representations for Siegel modular forms.
- The Niebur-Poincaré series may have useful applications for the Black Hole Farey Tale...
- Automorphic forms for exceptional groups are relevant to physics, and can be in principle constructed with similar methods, using other dual pairs such as $E_6 \times SL(3)$ in E_8 ...

$$E(s, \kappa, w) = \sum_{m \geq 0} b(s, \kappa, w, m) \mathcal{F}(s + m, \kappa, w),$$

$$b(s, \kappa, w, m) = \frac{2^{w-2s} (\pi \kappa)^{-s + \frac{w}{2}} \Gamma(2s + m - 1) \Gamma(s + m - \frac{w}{2})}{m! \Gamma(2s + 2m - 1) \Gamma(s - \frac{w}{2})}.$$

In the limit $s \rightarrow \frac{w}{2}$, for $w \leq 0$,

$$\begin{aligned} E\left(\frac{w}{2}, \kappa, w\right) &= \mathcal{F}\left(\frac{w}{2}, \kappa, w\right) + \sum_{m=1}^{-\frac{w}{2}-1} b'_m \operatorname{Res}_{s=\frac{w}{2}+m} \mathcal{F}(s, \kappa, w) \\ &+ \sum_{m=-\frac{w}{2}+1}^{1-w} b_m \mathcal{F}\left(\frac{w}{2} + m, \kappa, w\right) \end{aligned}$$