

Wall-crossing from quantum multi-centered BPS black holes

Boris Pioline

LPTHE, Jussieu




IPHT, Saclay, 23/5/2011

*based on work with J. Manschot and A. Sen,
arxiv:1011.1258, 1103.0261, 1103.1887*

Introduction I

- In $D = 4, N = 2$ supersymmetric field and string theories, the exact **spectrum of BPS states** can often be determined at weak coupling, and extrapolated to strong coupling.
- E.g., in pure $SU(2)$ Seiberg-Witten theory,



Seiberg Witten; Bilal Ferrari

Introduction II

- In following the BPS spectrum from weak to strong coupling, one must be wary of two issues:
 - short multiplets may pair up into a long multiplet,
 - single-particle states may decay into multi-particle states.
- The first issue can be avoided by considering a suitable index $\Omega(\gamma, t)$, designed such that contributions from long multiplets cancel. $\Omega(\gamma, t)$ is then a piecewise constant function of the charge vector γ and couplings/moduli t .
- To deal with the second issue, one must understand how $\Omega(\gamma, t)$ changes across a wall of marginal stability W , where a single-particle state with charge γ can decay into a multi-particle state with charges $\{\alpha_i\}$, such that $\gamma = \sum_i \alpha_i$.

Introduction III

- Initial progress came from physics, by noting that single-particle states (in a certain limit) can be represented by **multi-centered solitonic solutions**. Those exist only on one side of the wall and decay into the continuum of multi-particle states on the other side.
- The simplest decay $\gamma \rightarrow \gamma_1 + \gamma_2$, where γ_1, γ_2 are **primitive** charge vectors, involves only two-centered configurations, whose index is easily computed:

$$\Delta\Omega(\gamma \rightarrow \gamma_1 + \gamma_2) = (-1)^{\gamma_{12}+1} |\gamma_{12}| \Omega^+(\gamma_1) \Omega^+(\gamma_2)$$

Denef Moore

- In the **non-primitive** case $\gamma = M\gamma_1 + N\gamma_2$ where $M, N > 1$, many multi-centered configurations in general contribute, and computing their index is non-trivial.

- A general answer to this problem has come from the mathematical study of the wall-crossing properties of (generalized) **Donaldson-Thomas invariants** for Calabi-Yau three-folds, believed to be the mathematical translation of the BPS index $\Omega(\gamma)$ in type IIA CY vacua.
- Notably, **Kontsevich & Soibelman** (KS) and **Joyce & Song** (JS) gave two different-looking formulae for $\Delta\Omega(\gamma \rightarrow M\gamma_1 + N\gamma_2)$.
- The KS formula has already been derived/interpreted in several ways, e.g. by considering instanton corrections to the moduli space metric in 3D after compactification on a circle.

Gaiotto Moore Neitzke; Alexandrov BP Saueressig Vandoren

- Our goal will be to derive new wall-crossing formulae, based on the **quantization of multi-centered solitonic configurations**.

- 1 Introduction
- 2 Generalities, and a Boltzmannian view of wall-crossing
- 3 The Kontsevich-Soibelman-Joyce-Song formula
- 4 Non-primitive wall-crossing from localization
- 5 Away from the wall

- 1 Introduction
- 2 Generalities, and a Boltzmannian view of wall-crossing**
- 3 The Kontsevich-Soibelman-Joyce-Song formula
- 4 Non-primitive wall-crossing from localization
- 5 Away from the wall

- We consider $\mathcal{N} = 2$ supergravity in 4 dimensions (this includes field theories with rigid $\mathcal{N} = 2$ as a special case). Let $\Gamma = \Gamma_e \oplus \Gamma_m$ be the lattice of electric and magnetic charges, with antisymmetric (Dirac-Schwinger- Zwanziger) integer pairing

$$\langle \gamma, \gamma' \rangle = \langle (p^\Lambda, q_\Lambda), (p'^\Lambda, q'_\Lambda) \rangle \equiv q_\Lambda p'^\Lambda - q'_\Lambda p_\Lambda \in \mathbb{Z}$$

- BPS states preserve 4 out of 8 supercharges, and saturate the bound $M \geq |Z(\gamma, t^a)|$ where $Z(\gamma, t^a) = e^{\mathcal{K}/2}(q_\Lambda X^\Lambda - p^\Lambda F_\Lambda)$ is the central charge/stability data.
- We are interested in the index $\Omega(\gamma; t^a) = \text{Tr}_{\mathcal{H}'_\gamma(t^a)} (-1)^{2J_3}$ where $\mathcal{H}'_\gamma(t^a)$ is the Hilbert space of one-particle states with charge $\gamma \in \Gamma$ in the vacuum with vector moduli t^a .

- The BPS invariants $\Omega(\gamma; t^a)$ are locally constant functions of t^a , but may jump across codimension-one subspaces

$$W(\gamma_1, \gamma_2) = \{t^a / \arg[Z(\gamma_1)] = \arg[Z(\gamma_2)]\}$$

where γ_1 and γ_2 are two primitive (non-zero) vectors such that $\gamma = M\gamma_1 + N\gamma_2$, $M, N \geq 1$. Assume for definiteness that $\gamma_{12} < 0$.

- We choose γ_1, γ_2 such that $\Omega(\gamma; t^a)$ has support only on the **positive cone** (root basis property)

$$\tilde{\Gamma} : \{M\gamma_1 + N\gamma_2, \quad M, N \geq 0, \quad (M, N) \neq (0, 0)\} .$$

- Let c_{\pm} be the chamber in which $\arg(Z_{\gamma_1}) \geq \arg(Z_{\gamma_2})$. Our aim is to compute $\Delta\Omega(\gamma) \equiv \Omega^-(\gamma) - \Omega^+(\gamma)$ as a function of $\Omega^+(\gamma)$ (say).

Wall-crossing from semi-classical solutions I

- Assume that $M(\gamma_1), M(\gamma_2) \gg \Lambda, m_P$. Single-particle states which are potentially unstable across W are described by **classical configurations** with n centers of charge $\alpha_i = M_i\gamma_1 + N_i\gamma_2 \in \tilde{\Gamma}$, satisfying $(M, N) = \sum_i (M_i, N_i)$.
- Such bound states exist only on one side of the wall, and the distances r_{ij} diverge at the wall. Across the wall, the single-particle bound state has decayed into the continuum of multi-particle states. $\Delta\Omega(\gamma)$ is given by the index of such configurations.
- *In addition, in either chamber, there may be multi-centered configurations whose charge vectors do not lie in $\tilde{\Gamma}$. However, they remain bound across W and do not contribute to $\Delta\Omega(\gamma)$.*

Wall-crossing from semi-classical solutions II

- In $\mathcal{N} = 2$ supergravity (and presumably also in $\mathcal{N} = 2$ Abelian gauge theories), the locations of the centers are constrained by

$$\sum_{j=1 \dots n, j \neq i}^n \frac{\alpha_{ij}}{|\vec{r}_i - \vec{r}_j|} = c_i, \quad \begin{cases} c_i = 2 \operatorname{Im} [e^{-i\phi} Z(\alpha_i, t^a)] \\ \phi \equiv \arg[Z(\alpha_1 + \dots + \alpha_n, t^a)] \\ \alpha_{ij} \equiv \langle \alpha_i, \alpha_j \rangle \end{cases} \quad \text{Denef}$$

If all $\alpha_j \in \tilde{\Gamma}$, the constants c_i are given by $c_i = \Lambda \sum_{i \neq j} \alpha_{ij}$, with $\Lambda \rightarrow \infty$ near the wall.

- After factoring out an overall translational mode, the solution space is (generically) a $(2n - 2)$ -dimensional **symplectic manifold** $(\mathcal{M}_n(\alpha_{ij}, c_i), \omega)$, with $\omega = \frac{1}{2} \sum_{i < j} \alpha_{ij} \sin \theta_{ij} d\theta_{ij} \wedge d\phi_{ij}$.

de Boer El Showk Messamah Van den Bleeken

- Up to issues of statistics, $\Delta\Omega(\gamma)$ is equal to the index of the **SUSY quantum mechanics** on \mathcal{M}_n , multiplied by the index $\Omega(\gamma_i)$ of the internal d.o.f. carried by each center.

Wall-crossing from semi-classical solutions III

- For primitive decay $\gamma \rightarrow \gamma_1 + \gamma_2$, the quantization of the phase space $(\mathcal{M}_2, \omega) = (\mathbb{S}^2, \frac{1}{2}\gamma_{12} \sin \theta d\theta d\phi)$ reproduces the primitive WCF

$$\Delta\Omega(\gamma \rightarrow \gamma_1 + \gamma_2) = (-1)^{\gamma_{12}+1} |\gamma_{12}| \Omega^+(\gamma_1) \Omega^+(\gamma_2),$$

where $(-1)^{\gamma_{12}+1} |\gamma_{12}|$ is the index of the angular momentum multiplet of spin $j = \frac{1}{2}(\gamma_{12} - 1)$.

- This generalizes to **semi-primitive wall-crossing** $\gamma \rightarrow \gamma_1 + N\gamma_2$: unstable configurations consist of a **halo of n_s particles of charge $s\gamma_2$** , with total charge $\sum s n_s \gamma_2 = m\gamma_2$, orbiting around a **core of charge $\gamma_1 + (N - n)\gamma_2$** . The phase space is $\mathcal{M}_n = \prod_s (\mathcal{M}_2)^{n_s} / \mathbb{S}_{n_s}$.

Wall-crossing from semi-classical solutions IV

- Taking into account the **Bose/Fermi statistics** of the n_s identical particles, one arrives at a Mac-Mahon type partition function,

$$\frac{\sum_{N \geq 0} \Omega^-(1, N) q^N}{\sum_{N \geq 0} \Omega^+(1, N) q^N} = \prod_{k > 0} \left(1 - (-1)^{k\gamma_{12}} q^k \right)^{k|\gamma_{12}| \Omega^+(k\gamma_2)} .$$

Denef Moore

- E.g. for $\gamma \mapsto \gamma_1 + 2\gamma_2$,

$$\begin{aligned} \Delta\Omega(1, 2) = & (-1)^{\gamma_{12}} \gamma_{12} \Omega^+(0, 1) \Omega^+(1, 1) + 2\gamma_{12} \Omega^+(0, 2) \Omega^+(1, 0) \\ & + \frac{1}{2} \gamma_{12} \Omega^+(0, 1) (\gamma_{12} \Omega^+(0, 1) + 1) \Omega^+(1, 0) . \end{aligned}$$

In particular, the term $\frac{1}{2}d(d+1)$ with $d = \gamma_{12}\Omega^+(0, 1)$, reflects the projection on **(anti)symmetric wave functions**.

Wall-crossing from semi-classical solutions V

- It is instructive to rewrite the semi-primitive WCF using the **rational BPS invariants**, related to the usual integer invariants via

$$\bar{\Omega}(\gamma) \equiv \sum_{d|\gamma} \Omega(\gamma/d)/d^2, \quad \Omega(\gamma) = \sum_{d|\gamma} \mu(d) \bar{\Omega}(\gamma/d)/d^2$$

where $\mu(d)$ is the Möbius function.

- Using the identity $\prod_{d=1}^{\infty} (1 - q^d)^{\mu(d)/d} = e^{-q}$, or working backwards, one arrives at

$$\frac{\sum_{N \geq 0} \bar{\Omega}^-(1, N) q^N}{\sum_{N \geq 0} \bar{\Omega}^+(1, N) q^N} = \exp \left[\sum_{s=1}^{\infty} q^s (-1)^{\langle \gamma_1, s\gamma_2 \rangle} \langle \gamma_1, s\gamma_2 \rangle \bar{\Omega}^+(s\gamma_2) \right].$$

- The same result follows by treating particles in the halo as **distinguishable** (satisfying Boltzmann statistics), and attaching an effective index $\bar{\Omega}(s\gamma_2)$!

Wall-crossing from semi-classical solutions VI

- One advantage is that $\Delta\bar{\Omega}(\gamma)$ takes a simpler form, and makes charge conservation manifest. E.g for $\gamma \mapsto \gamma_1 + 2\gamma_2$,

$$\Delta\bar{\Omega}(1, 2) = (-1)^{\gamma_{12}}\gamma_{12}\bar{\Omega}^+(0, 1)\bar{\Omega}^+(1, 1) + 2\gamma_{12}\bar{\Omega}^+(0, 2)\bar{\Omega}^+(1, 0) \\ + \frac{1}{2}\gamma_{12}\bar{\Omega}^+(0, 1)^2\bar{\Omega}^+(1, 0) .$$

- The rational DT invariants $\bar{\Omega}(\gamma)$ appear in the JS formula, in constructions of **modular invariant black hole partition functions**, and in **instanton corrections to hypermultiplet moduli spaces**.

Joyce Song; Manschot; Alexandrov BP Saueressig Vandoren

The main conjecture I

- In general, we expect that the jump to be given by a finite sum

$$\Delta\bar{\Omega}(\gamma) = \sum_{n \geq 2} \sum_{\substack{\{\alpha_1, \dots, \alpha_n\} \in \tilde{\Gamma} \\ \gamma = \alpha_1 + \dots + \alpha_n}} \frac{g(\{\alpha_j\})}{|\text{Aut}(\{\alpha_j\})|} \prod_{i=1}^n \bar{\Omega}^+(\alpha_i),$$

over all **unordered** decompositions of the total charge vector γ into a sum of n vectors $\alpha_j \in \tilde{\Gamma}$. The symmetry factor $|\text{Aut}(\{\alpha_j\})|$ reflects **Boltzmannian statistics**.

- $g(\{\alpha_j\})$ are universal factors depending only on the charges α_j , which should be given by the **index of the supersymmetric quantum mechanics on \mathcal{M}_n** .
- The KS and JS formulae give a mathematical prediction for these coefficients $g(\{\alpha_j\})$, which we shall compare with the index.

- 1 Introduction
- 2 Generalities, and a Boltzmannian view of wall-crossing
- 3 The Kontsevich-Soibelman-Joyce-Song formula**
- 4 Non-primitive wall-crossing from localization
- 5 Away from the wall

The Kontsevich-Soibelman formula I

- Consider the Lie algebra \mathcal{A} spanned by abstract generators $\{e_\gamma, \gamma \in \Gamma\}$, satisfying the commutation rule

$$[e_{\gamma_1}, e_{\gamma_2}] = (-1)^{\langle \gamma_1, \gamma_2 \rangle} \langle \gamma_1, \gamma_2 \rangle e_{\gamma_1 + \gamma_2} .$$

- For a given charge vector γ and value of the VM moduli t^a , consider the operator $U_\gamma(t^a)$ in the Lie group $\exp(\mathcal{A})$

$$U_\gamma(t^a) \equiv \exp \left(\Omega(\gamma; t^a) \sum_{d=1}^{\infty} \frac{e_{d\gamma}}{d^2} \right)$$

- The operators e_γ / U_γ can be realized as **Hamiltonian vector fields** / **symplectomorphisms** of a twisted torus.

Gaiotto Moore Neitzke

The Kontsevich-Soibelman formula II

- The KS wall-crossing formula states that the product

$$A_{\gamma_1, \gamma_2} = \prod_{\substack{\gamma = M\gamma_1 + N\gamma_2, \\ M \geq 0, N \geq 0}} U_{\gamma},$$

ordered so that $\arg(Z_{\gamma})$ decreases from left to right, stays constant across the wall. As t^a crosses W , $\Omega(\gamma; t^a)$ jumps and the order of the factors is reversed, but the operator A_{γ_1, γ_2} stays constant. Equivalently,

$$\prod_{\substack{M \geq 0, N \geq 0, \\ M/N \downarrow}} U_{M\gamma_1 + N\gamma_2}^+ = \prod_{\substack{M \geq 0, N \geq 0, \\ M/N \uparrow}} U_{M\gamma_1 + N\gamma_2}^- ,$$

The Kontsevich-Soibelman formula III

- The algebra \mathcal{A} is infinite dimensional but filtered. The KS formula may be projected to any finite-dimensional algebra

$$\mathcal{A}_{M,N} = \mathcal{A} / \left\{ \sum_{m>M \text{ or } n>N} \mathbb{R} \cdot e_{m\gamma_1 + n\gamma_2} \right\} .$$

This projection is sufficient to infer $\Delta\Omega(m\gamma_1 + n\gamma_2)$ for any $m \leq M, n \leq N$, e.g. using the Baker-Campbell-Hausdorff formula.

- For example, the primitive WCF follows in $\mathcal{A}_{1,1}$ from

$$\begin{aligned} & \exp(\bar{\Omega}^+(\gamma_1)e_{\gamma_1}) \exp(\bar{\Omega}^+(\gamma_1 + \gamma_2)e_{\gamma_1 + \gamma_2}) \exp(\bar{\Omega}^+(\gamma_2)e_{\gamma_2}) \\ &= \exp(\bar{\Omega}^-(\gamma_2)e_{\gamma_2}) \exp(\bar{\Omega}^-(\gamma_1 + \gamma_2)e_{\gamma_1 + \gamma_2}) \exp(\bar{\Omega}^-(\gamma_1)e_{\gamma_1}) \end{aligned}$$

and the order 2 truncation of the BCH formula

$$e^X e^Y = e^{X+Y + \frac{1}{2}[X,Y]} .$$


The Kontsevich-Soibelman formula IV

- In some simple cases, one may work in the full algebra \mathcal{A} , and use the “pentagonal identity”

$$U_{\gamma_2} U_{\gamma_1} = U_{\gamma_1} U_{\gamma_1 + \gamma_2} U_{\gamma_2}, \quad \gamma_{12} = -1$$

- Using this identity repeatedly, one can e.g. establish the wall-crossing identity in pure Seiberg-Witten $SU(2)$ theory,

$$U_{2,-1} \cdot U_{0,1} = U_{0,1} \cdot U_{2,1} \cdot U_{4,1} \cdots U_{2,0}^{(-2)} \cdots U_{3,-1} \cdot U_{2,-1} U_{1,-1}$$



Denef Moore; Dimofte Gukov Soibelman

The Kontsevich-Soibelman formula V

- Noting that the operators $U_{k\gamma}$ for different $k \geq 1$ commute, one may combine them into a single factor

$$V_\gamma \equiv \prod_{k=1}^{\infty} U_{k\gamma} = \exp \left(\sum_{\ell=1}^{\infty} \bar{\Omega}(\ell\gamma) e_{\ell\gamma} \right), \quad \bar{\Omega}(\gamma) = \sum_{d|\gamma} \Omega(\gamma/d)/d^2.$$

and rewrite the KS formula as a product over **primitive** charge vectors only,

$$\prod_{\substack{M \geq 0, N \geq 0, \\ \gcd(M, N) = 1, M/N \downarrow}} V_{M\gamma_1 + N\gamma_2}^+ = \prod_{\substack{M \geq 0, N \geq 0, \\ \gcd(M, N) = 1, M/N \uparrow}} V_{M\gamma_1 + N\gamma_2}^-$$

The Kontsevich-Soibelman formula VI

- Using the BCH formula, one easily derives the semi-primitive wcf formula, and generalizations to $\gamma \rightarrow 2\gamma_1 + N\gamma_2, \dots$
- The fact that the algebra is graded by the charge lattice and the expression of V_γ guarantees that the jumps in the rational invariant will be of the form

$$\Delta\bar{\Omega}(\gamma) = \sum_{n \geq 2} \sum_{\substack{\{\alpha_1, \dots, \alpha_n\} \in \tilde{\Gamma} \\ \gamma = \alpha_1 + \dots + \alpha_n}} \frac{g(\{\alpha_j\})}{|\text{Aut}(\{\alpha_j\})|} \prod_{i=1}^n \bar{\Omega}^+(\alpha_i),$$

with some universal coefficients $g(\{\alpha_j\})$.

- The Joyce-Song wall-crossing formula expresses $g(\{\alpha_j\})$ as a complicated sum over trees, permutations, etc.

Generic decay I

- When α_j have generic phases, $g(\{\alpha_j\})$ can be computed by projecting the KS formula to the subalgebra spanned by $e_{\sum \alpha_j}$ where $\{\alpha_j\}$ runs over all subsets of $\{\alpha_j\}$.
- E.g., for $n = 3$, assuming that the phase of the charges are ordered according to

$$\alpha_1, \alpha_1 + \alpha_2, \alpha_1 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, \alpha_2, \alpha_2 + \alpha_3, \alpha_3,$$

we find

$$g(\{\alpha_1, \alpha_2, \alpha_3\}) = (-1)^{\alpha_{12} + \alpha_{23} + \alpha_{13}} \alpha_{12} (\alpha_{13} + \alpha_{23})$$

As we shall see later, this fits the macroscopic index of 3-centered configurations !

The motivic Kontsevich-Soibelman formula I

- KS have proposed a **quantum deformation** of their formula, which governs wall-crossing properties of **motivic DT invariants** $\Omega_{\text{ref}}(\gamma; y, t)$. Physically, these correspond to the “refined index”

$$\Omega_{\text{ref}}(\gamma, y) = \text{Tr}'_{\mathcal{H}(\gamma)}(-y)^{2J_3} \equiv \sum_{n \in \mathbb{Z}} (-y)^n \Omega_{\text{ref},n}(\gamma),$$

where J_3 is the angular momentum in 3 dimensions along the z axis. As $y \rightarrow 1$, $\Omega_{\text{ref}}(\gamma; y, t) \rightarrow \Omega(\gamma; t)$.

Dimofte Gukov; D G Soibelman

- Caution: this index (rather, a variant of it using a combination of angular momentum and $SU(2)_R$ quantum numbers) is protected in $\mathcal{N} = 2, D = 4$ field theories, but not in supergravity/string theory, where $SU(2)_R$ is generically broken.

Gaiotto Moore Neitzke

The motivic Kontsevich-Soibelman formula II

- To state the formula, consider the Lie algebra $\mathcal{A}(y)$ spanned by generators $\{\tilde{\mathbf{e}}_\gamma, \gamma \in \Gamma\}$, satisfying the commutation rule

$$[\tilde{\mathbf{e}}_{\gamma_1}, \tilde{\mathbf{e}}_{\gamma_2}] = \kappa(\langle \gamma_1, \gamma_2 \rangle) \tilde{\mathbf{e}}_{\gamma_1 + \gamma_2}, \quad \kappa(x) = \frac{(-y)^x - (-y)^{-x}}{y - 1/y}.$$

- To any primitive charge vector γ , attach the operator

$$\hat{V}_\gamma = \prod_{k \geq 1} \hat{U}_{k\gamma} = \exp \left[\sum_{\ell=1}^{\infty} \bar{\Omega}_{\text{ref}}(\ell\gamma, y) \tilde{\mathbf{e}}_{\ell\gamma} \right]$$

where $\bar{\Omega}_{\text{ref}}(\gamma, y)$ are the **“rational motivic invariants”**, defined by

$$\bar{\Omega}_{\text{ref}}^+(\gamma, y) \equiv \sum_{d|\gamma} \frac{(y - y^{-1})}{d(y^d - y^{-d})} \Omega_{\text{ref}}^+(\gamma/d, y^d).$$

The motivic Kontsevich-Soibelman formula III

- The motivic version of the KS wall-crossing formula states that

$$\prod_{\substack{M \geq 0, N \geq 0 > 0, \\ \gcd(M, N) = 1, M/N \downarrow}} \hat{V}_{M\gamma_1 + N\gamma_2}^+ = \prod_{\substack{M \geq 0, N \geq 0 > 0, \\ \gcd(M, N) = 1, M/N \uparrow}} \hat{V}_{M\gamma_1 + N\gamma_2}^- ,$$

- $\Delta \bar{\Omega}_{\text{ref}}(\gamma, y)$ can be computed using the same techniques as before, e.g. the primitive wcf read

$$\Delta \Omega_{\text{ref}}(\gamma_1 + \gamma_2, y) = \frac{(-y)^{\langle \gamma_1, \gamma_2 \rangle} - (-y)^{-\langle \gamma_1, \gamma_2 \rangle}}{y - 1/y} \Omega_{\text{ref}}(\gamma_1, y) \Omega_{\text{ref}}(\gamma_2, y)$$

- The general formula for $\Delta \bar{\Omega}_{\text{ref}}$ involves universal factors $g(\{\alpha_j\}, y)$, which reduce to $g(\{\alpha_j\})$ in the limit $y \rightarrow 1$. We expect that they are given by $\text{Tr}'(-y)^{2J_3}$ in the corresponding SUSY quantum mechanics.

- 1 Introduction
- 2 Generalities, and a Boltzmannian view of wall-crossing
- 3 The Kontsevich-Soibelman-Joyce-Song formula
- 4 Non-primitive wall-crossing from localization**
- 5 Away from the wall

Quantum mechanics of multi-centered solutions I

- The moduli space \mathcal{M}_n of BPS configurations with n centers in $\mathcal{N} = 2$ SUGRA is described by solutions to Denef's equations

$$\sum_{j=1 \dots n, j \neq i}^n \frac{\alpha_{ij}}{|\vec{r}_i - \vec{r}_j|} = c_i, \quad \begin{cases} c_i = 2 \operatorname{Im} [e^{-i\phi} Z(\alpha_i)] \\ \phi = \arg[Z(\alpha_1 + \dots + \alpha_n)] \end{cases} .$$


- \mathcal{M}_n is a **symplectic manifold** of dimension $2n - 2$, and carries an Hamiltonian action of $SU(2)$:

$$\omega = \frac{1}{2} \sum_{i < j} \alpha_{ij} \sin \theta_{ij} d\theta \wedge d\phi_{ij}, \quad \vec{J} = \frac{1}{2} \sum_{i < j} \alpha_{ij} \frac{\vec{r}_{ij}}{|\vec{r}_{ij}|}$$

de Boer El Showk Messamah Van den Bleeken

Quantum mechanics of multi-centered solutions II

- When the α_i 's lie in the positive cone $\tilde{\Gamma}$ (more generally, whenever $\text{sign}(\alpha_{ij})$ defines an ordering of the α_i , \mathcal{M}_n is **compact**, and the fixed points of J_3 are **isolated**.
- E.g for 3 centers with $\alpha_{12} > 0, \alpha_{23} > 0, \alpha_{13} > 0$, the domain of the plane $c_1 + c_2 + c_3 = 0$ allowed by Denef's equations is:



For fixed c_i , the range of r_{ij} is read off by intersecting the shaded area with a radial line which joins c_i to the origin. Here, 3-centered solutions only exist in the region $c_1 > 0, c_3 < 0$, and have r_{ij} bounded from below and from above. Fixed points of J_3 correspond to collinear solutions, and lie on the boundary of this domain.

Quantum mechanics of multi-centered solutions III

- The symplectic form $\omega/2\pi \in H^2(\mathcal{M}_n, \mathbb{Z})$ is the curvature of a **complex line bundle** \mathcal{L} over \mathcal{M}_n , with connection

$$\lambda = \frac{1}{2} \sum_{i < j} \alpha_{ij} (1 - \cos \theta_{ij}) d\phi_{ij}, \quad d\lambda = \omega.$$

- Assuming that \mathcal{M}_n is spin, let $S = S_+ \oplus S_-$ be the spin bundle. Let $D = D_+ \oplus D_-$ be the **Dirac operator** for the metric obtained by restricting the flat metric on \mathbb{R}^{3n-3} to \mathcal{M}_n , with $D_{\pm} : S_{\pm} \mapsto S_{\mp}$. The action of $SO(3)$ on \mathcal{M}_n lifts to an action of $SU(2)$ on S_{\pm} .
- We assume that BPS states correspond to **harmonic spinors**, i.e. sections of $S \otimes \mathcal{L}$ annihilated by the Dirac operator D .

Quantum mechanics of multi-centered solutions IV

- The ‘refined index’ is then given by

$$g_{\text{ref}}(\{\alpha_j\}; y) = \text{Tr}_{\text{Ker}D_+} (-y)^{2J_3} + \text{Tr}_{\text{Ker}D_-} (-y)^{2J_3} .$$

- We further assume that $\text{Ker}D_- = 0$, so that the refined index $g_{\text{ref}}(\{\alpha_j\}; y)$ reduces to the **equivariant index**

$$g_{\text{ref}}(\{\alpha_j\}; y) = \text{Tr}_{\text{Ker}D_+} (-y)^{2J_3} - \text{Tr}_{\text{Ker}D_-} (-y)^{2J_3} .$$

- The vanishing of $\text{Ker}D_-$ can be shown to hold in special cases where \mathcal{M}_n is Kähler. In gauge theories, the **protected spin character** presumably reduces to the equivariant index without further assumption.

Quantum mechanics of multi-centered solutions V

- The refined/equivariant index can be computed by the **Atiyah-Bott Lefschetz fixed point formula**:

$$g_{\text{ref}}(\{\alpha_j\}, y) = \sum_{\text{fixed pts}} \frac{y^{2J_3}}{\det((-y)^L - (-y)^{-L})}$$

where L is the matrix of the action of J_3 on the holomorphic tangent space around the fixed point.

- In the large charge limit, $\mathcal{L} \rightarrow k\mathcal{L}$ with $k \rightarrow \infty$, this reduces to the **Duistermaat-Heckmann formula** for the equivariant volume,

$$\frac{\int_{\mathcal{M}_n} \omega^{n-1} y^{2J_3}}{(2\pi)^{n-1} (n-1)!} = \sum_{\text{fixed pts}} \frac{y^{2J_3}}{\det(L \log(-y))}$$

- The fixed points of the action of J_3 are **collinear multi-centered configurations** along the z -axis, such that

$$\sum_{j=1 \dots n, j \neq i}^n \frac{\alpha_{ij}}{|z_i - z_j|} = c_i, \quad J_3 = \frac{1}{2} \sum_{i < j} \alpha_{ij} \text{sign}(z_j - z_i).$$

- Equivalently, fixed points are critical points of the ‘superpotential’

$$W(\lambda, \{z_i\}) = - \sum_{i < j} \text{sign}[z_j - z_i] \alpha_{ij} \ln |z_j - z_i| - \sum_i (c_i - \frac{\lambda}{n}) z_i$$

These are **isolated**, and classified by **permutations** describing the order of z_i along the axis.

- This leads to the **Coulomb branch formula**

$$g_{\text{ref}}(\{\alpha_j\}, y) = \frac{(-1)^{\sum_{i<j} \alpha_{ij} + n - 1}}{(y - y^{-1})^{n-1}} \sum_{p: \partial_l W(p)=0} s(p) y^{\sum_{i<j} \alpha_{ij} \text{sign}(z_j - z_i)}$$

where $s(p) = -\text{sign}(\det W_{IJ})$

- For $n \leq 5$, we find perfect agreement with JS/KS !

$$g_{\text{ref}}(\alpha_1, \alpha_2; y) = (-1)^{\alpha_{12}} \frac{\sinh(\nu \alpha_{12})}{\sinh \nu} \quad (y = e^\nu)$$

$$g_{\text{ref}}(\alpha_1, \alpha_2, \alpha_3; y) = (-1)^{\alpha_{13} + \alpha_{23} + \alpha_{12}} \frac{\sinh(\nu(\alpha_{13} + \alpha_{23})) \sinh(\nu \alpha_{12})}{\sinh^2 \nu}$$

Higgs branch picture I

- An alternative formula can be given using the **Higgs branch** description of the multi-centered configuration, namely the **quiver** with n nodes $\{1 \dots n\}$ of dimension 1 and α_{ij} arrows from i to j .
- Since α_i lie on a 2-dimensional sublattice $\tilde{\Gamma}$, the quiver has no oriented closed loop. **Reineke's formula** gives

$$g_{\text{ref}} = \frac{(-y)^{-\sum_{i<j} \alpha_{ij}}}{(y - 1/y)^{n-1}} \sum_{\text{partitions}} (-1)^{s-1} y^{2 \sum_{a \leq b} \sum_{j<i} \alpha_{ji} m_i^{(a)} m_j^{(b)}},$$

where \sum runs over all ordered partitions of $\gamma = \alpha_1 + \dots + \alpha_n$ into s vectors $\beta^{(a)}$ ($1 \leq a \leq s$, $1 \leq s \leq n$) such that

- 1 $\beta^{(a)} = \sum_i m_i^{(a)} \alpha_i$ with $m_i^{(a)} \in \{0, 1\}$, $\sum_a \beta^{(a)} = \gamma$
- 2 $\langle \sum_{a=1}^b \beta^{(a)}, \gamma \rangle > 0 \quad \forall \quad b$ with $1 \leq b \leq s-1$

Higgs branch picture II

- The Higgs branch formula agrees with KS/JS/Coulomb for $n = 2, 3, 4, 5$!
- The formula gives a prescription for what permutations are allowed in the Coulomb problem, and for their Morse index.
- It is perhaps not surprising that the Higgs branch formula agrees with KS/JS, since quiver categories are an example of Abelian categories. Unlike the JS formula, the Higgs branch formula works at $y \neq 1$.

- 1 Introduction
- 2 Generalities, and a Boltzmannian view of wall-crossing
- 3 The Kontsevich-Soibelman-Joyce-Song formula
- 4 Non-primitive wall-crossing from localization
- 5 Away from the wall**

- Having understood the jump $\Delta\Omega(\gamma; y)$ in terms of the index of multi-centered solutions, one would like to compute the BPS index $\Omega(\gamma; y, t^a)$ on either side of the wall, from the index $\Omega_S(\alpha_j)$ of **single-centered black holes**. Since spherically symmetric SUSY black holes cannot decay and carry zero angular momentum, $\Omega_S(\alpha_j)$ must be independent of t^a and y .

Manchot BP Sen II


- Naively, one may expect

$$\bar{\Omega}(\gamma; y, t^a) = \sum_{n \geq 2} \sum_{\substack{\{\alpha_1, \dots, \alpha_n\} \in \Gamma \\ \gamma = \alpha_1 + \dots + \alpha_n}} \frac{g(\{\alpha_j\}; y, c_j)}{|\text{Aut}(\{\alpha_j\})|} \prod_{j=1}^n \bar{\Omega}_S(\alpha_j),$$

where $g(\{\alpha_j\}; y, c_j)$ is the refined index of the SUSY quantum mechanics on $\mathcal{M}_n(\alpha_j, c_j)$. This is similar to the formula for $\Delta\bar{\Omega}(\gamma)$, but with some important differences.

- Unlike the formula for $\Delta\bar{\Omega}(\gamma)$, the charges α_i of the constituents are **no longer restricted to a two-dimensional subspace** of the charge lattice, and there are a priori an infinite number of possible splittings $\gamma = \sum \alpha_i$. It is plausible that requiring that the multi-centered solution be **regular** may leave only a finite number of splittings. In addition, for a given splitting, the regularity constraint may rule out certain components of $\mathcal{M}_n(\alpha_i, c_i)$.
- The space $\mathcal{M}_n(\alpha_i, c_i)$ is in general **no longer compact**. E.g, in the 3-body case with $\alpha_{12} > 0, \alpha_{23} > 0, \alpha_{13} < 0$, the allowed values of c_i (and therefore r_{ij}) are plotted below:

Away from the wall III



- In particular, there can be **scaling regions** in \mathcal{M}_n , when some or all of the n centers approach each other at arbitrary small distances. Classically, these scaling solutions carry zero angular momentum and are invariant under $SO(3)$.
- Some of the distances r_{ij} can also diverge on walls of marginal stability, but the formula for $\Omega(\gamma, t^a)$ is by construction consistent with wall-crossing.

- In the presence of scaling solutions, it appears that \mathcal{M}_n admits a compactification $\overline{\mathcal{M}}_n$ with finite volume. However, this introduces new (non-collinear) fixed points of the action of J_3 which are **no longer isolated**, leading to additional contributions to the equivariant index.
- Rather than trying to compute these new contributions directly, we propose to determine them by requiring 1) that the resulting $\Omega(\gamma; \mathbf{y}, t^a)$ is a **finite Laurent polynomial in \mathbf{y}** and 2) that they carry the **minimal angular momentum J_3** compatible with condition 1). This **minimal modification hypothesis** fixes $\Omega(\gamma; \mathbf{y}, t^a)$ uniquely.
- We have checked that the minimal modification hypothesis works for an infinite class of ‘dipole halo’ configurations, where \mathcal{M}_n is a toric manifold and can be quantized directly.

Conclusion I

- Multi-centered solitonic configurations provide a simple picture to derive and understand wall-crossing formulae for the BPS (refined) index.
- We have not proven the equivalence between the Coulomb branch, Higgs branch, JS and KS wall-crossing formula, but there is overwhelming evidence that they all agree.
- Our derivation was made in the context of $\mathcal{N} = 2$ supergravity, it would be interesting to develop our understanding of multi-centered dyonic solutions in $\mathcal{N} = 2$ gauge theories.

Lee Yi

- In principle, our formulae can be used to extract the degeneracies $\Omega_S(\gamma)$ of single-centered black holes from the moduli-dependent BPS index $\Omega(\gamma)$. The former is the one that should be compared with Sen's quantum entropy function.

THANK YOU !