

Lecture II: Integrability in Strings on Coset Spaces

- Strings in Flat Space
- Strings on Coset Spaces
- Formalism: Currents, Action, Equations of Motion
- Lax Connection, Integrability
- Wilson Lines, Monodromy, Spectral Curve
- Properties of the Curve
- Moduli of the Curve, Cycles
- Ansätze
- Spectral Curves for various Models

Strings in Flat Space

Fields: $\vec{X}: \mathbb{R}^{1,1} \rightarrow \mathbb{R}^{D-1,1}$
 $\uparrow \quad \quad \quad \uparrow$
 2D World Sheet $\quad$ D-dim Target Space

Periodicity: $\vec{X}(\tau, \sigma + 2\pi) = \vec{X}(\tau, \sigma)$

Action (Polyakov)

$$S \sim \int d\tau \int_0^{2\pi} d\sigma \quad dX_\mu \wedge * dX^\mu$$

EOM: $d * d\vec{X} = 0$

General Solution of EOM (Fourier mode decomposition)

$$\vec{X}(\tau, \sigma) = \vec{X}_0 + \vec{P}\tau + \sum_{n \neq 0} \text{Re } c_n \exp(in\sigma + i\tau \ln|n|)$$

Easily quantised: set of independent HO modes

String theory has also WS-metric γ_{ab} (in *)

Virasoro constraint: $(\partial_\pm \vec{X})^2 = 0$ from variation of γ_{ab}
 $\uparrow$
 $\partial_\pm = \partial_\tau \pm \partial_\sigma$

Constraints imposed on quantised system.

Strings on Coset Spaces

eg. $AdS_5 \times S^5$, Non-Linear Sigma Models, Non-Trivial models

$$S_{N-1} = SO(N)/SO(N-1)$$

$$AdS_{N-1} = SO(N-2,2)/SO(N-2,1) \text{ (universal cover)}$$

$$\left. \begin{array}{l} S_{N-1} \\ AdS_{N-1} \end{array} \right\} G/H$$

Field: $g(\tau, \sigma) \in G$ eg. $N \times N$ orthogonal matrix for S_{N-1}

Coset: $g(\tau, \sigma) \hat{=} g(\tau, \sigma) h(\tau, \sigma)$, $h \in H$ gauge transformation

Periodicity: $g(\tau, \sigma + 2\pi) = g(\tau, \sigma) h(\tau, \sigma) \hat{=} g(\tau, \sigma)$, $h \in H$

Currents (moving frame)

Full Current: $J = -g^{-1} dg \in \mathfrak{g}$ (algebra of G)

Decompose $J = B + P$, $B \in \mathfrak{h}$ (algebra of H) gauge field, $P \perp \mathfrak{h}$ momentum (unphysical) (physical)

Action $S = \frac{\sqrt{\lambda}}{2\pi} \int d\tau \int_0^{2\pi} d\sigma \frac{1}{2} \text{tr}(P_\mu \star P^\mu)$ Currents (static frame)
lowercase $j = g^{-1} J g^{-1}$ easier
 $J = g^{-1} j g$

EOM: $d \star P = B_\mu \star P^\mu + \star P_\mu B^\mu$

Jacobi identities: $dJ = J_\mu J^\mu \rightarrow \begin{cases} dB = B_\mu B^\mu + P_\mu P^\mu \\ dP = B_\mu P^\mu + P_\mu B^\mu \end{cases}$

EOM: $d \star p = 0$ (p is Noether cons.)

Jacobi: $dj = -j_\mu j^\mu \rightarrow \begin{cases} db = -b_\mu p^\mu - p_\mu b^\mu \\ dp = -2p_\mu p^\mu \end{cases}$
 p is gauge invariant

String Theory $\rightarrow$ WS-metric

Virasoro constraint: $\text{tr}(P_\pm)^2 = 0$ Virasoro: $\text{tr}(p_\pm)^2 = 0$

will be imposed later

Action: $S = \frac{\sqrt{\lambda}}{2\pi} \int d\tau \int_0^{2\pi} d\sigma \frac{1}{2} \text{tr}(p_\mu \star p^\mu)$

Integrable Structures on The Worldsheet

Useful for extracting physical data from WS

Let's construct a flat connection $D = d + a$, ie $D^2 = da + a \wedge a = 0$

Ansatz: $a = \alpha p + \beta *p$ $\xrightarrow{\text{Jacobi}} 2p \wedge p = 0$ by sym

$$\begin{aligned} \text{Flatness: } D^2 &= \alpha dp + \beta d*p - \alpha^2 p \wedge p - \alpha\beta p \wedge *p - \alpha\beta *p \wedge p - \beta^2 *p \wedge *p \\ &= (-2\alpha - \alpha^2 + \beta^2) p \wedge p \stackrel{!}{=} 0 \end{aligned}$$

Solution: $\alpha = \frac{2}{x^2-1}, \quad \beta = \frac{2x}{x^2-1}$


Lax connection: $a(x) = \frac{2}{x^2-1} p + \frac{2x}{x^2-1} *p$

Family of flat connections $(D(x))^2 = 0 \quad da(x) = a(x) \wedge a(x)$

Wilson Lines on the WS & Monodromy


Parallel transport of connection $d - a(x)$ ^{fixed "spectral" par.} along curve γ on WS

Wilson line: $\omega(\gamma, x) = \text{Pexp} \int_{\gamma} (-a(x))$



Connection $d - a(x)$ is flat $\Rightarrow \omega$ indep. of deformations of γ

$$\omega(\gamma, x) = \omega(\gamma', x) \text{ if } \gamma' \text{ deformed } \gamma$$



define Monodromy $\omega(x) = \omega(\gamma, x)$ where γ once around string

$$\text{eg. } \omega(x) = \text{Pexp} \oint_0^{2\pi} (-a_{\sigma}(x)) d\sigma$$



$\omega(x)$ indep. of γ , but not of endpoints $\gamma(0), \gamma(1)$



$$\omega'(x) = \omega(\tilde{\gamma}^{-1} \gamma \tilde{\gamma}, x) = \omega(\tilde{\gamma}, x)^{-1} \omega(x) \omega(\tilde{\gamma}, x)$$

similarity transformation $\omega'(x) \cong \omega(x)$ by $\omega(\tilde{\gamma}, x)$

Eigenvalues of monodromy indep. of WS!

$$\omega(x) \cong \text{diag}(\omega_1(x), \dots, \omega_N(x))$$

Eigenvalues $\omega_k(x)$ depend on spectral parameter x only.

Still a lot of data on string, but only conserved qts.

Have transformed $g(\tau, \sigma)$ to $\omega_k(x)$

string embedding

~~the~~ conserved quantities

Investigate properties of $\omega_k(x)$

Diagonalisation and Branch Points

$\det(\lambda - \omega(x)) = (\lambda - \omega_1(x)) \dots (\lambda - \omega_N(x))$
 $\{\omega_k\}$ form N Riemann sheets of a function $\omega(x)$
 when two EV degenerate. consider 2×2 submatrix


$$\omega(x) = \begin{pmatrix} a(x) & b(x) \\ c(x) & d(x) \end{pmatrix} \quad \omega_{1,2}(x) = \frac{1}{2}(a+d) \pm \sqrt{\frac{1}{4}(a-d)^2 + bc}$$

Typically at degeneracy point $x_k \sim (x - x_k)$

$$\omega_{1,2}(x) = \omega_k(x) \pm \alpha_k \sqrt{x - x_k}$$

Two sheets join at degeneracy point x_k

Branch cut originates from x_k (and ends at x_l).



2 Riemann sheets

$\omega(x)$ has N sheets connected by branch cuts.

Singularities at $x = \pm 1$

$$a(x) = \frac{2}{x^2-1} p + \frac{2x}{x^2-1} * p$$

$$a_\sigma(x) \stackrel{x \rightarrow 1}{\sim} \frac{1}{x-1} p_\sigma + \frac{1}{x-1} p_\tau + O((x-1)^0) = \frac{1}{x-1} p_+ + \dots$$

Diagonalise $p_+ = S \tilde{p}_+ S^{-1}$

Then

$$S^{-1} (\partial_\sigma + a_\sigma(x)) S = \frac{1}{x-1} \tilde{p}_+ \frac{\partial}{\partial \sigma} + O((x-1)^0)$$

$$\omega(x) \stackrel{x \rightarrow 1}{\sim} \exp \left(-\frac{1}{x-1} \int_0^{2\pi} \tilde{p}_+ d\sigma + O((x-1)^0) \right)$$

Exponential singularity!

Consider "quasi-momenta" $\omega_k(x) = \exp i q_k(x)$

Single poles in $q_k(x)$ at $x = \pm 1$

Residues linked by Virasoro $h(p_\pm)^2 = 0$

Moduli of Spectral Curves

$w(x)$ is single-valued on the curve

$q(x) = -i \log w(x)$ is single-valued modulo 2π .


$q'(x) = -i w'(x)/w(x)$ is single-valued

(closed cycles) $\oint dq = \oint q'(x) dx \in 2\pi \mathbb{Z}$

Can arrange branch cuts s.t.

A -cycles vanish

$$\oint_{A_k} dq = 0$$



Moduli:

Mode numbers: $n_k = \frac{1}{2\pi} \oint_{B_k} dq \in \mathbb{Z}$ discrete!

Filling $k_k = \frac{\sqrt{\lambda}}{8\pi^2 i} \oint_{A_k} (x + 1/x) dq \sim \text{length of cut } C_k$

Compare to flat space

cut C_k corresponds to excitation of mode n_k
with amplitude k_k

$\Rightarrow$ Structure qualitatively similar to flat space!

Global Charges at $x = \infty$

Expand at $x = \infty$

$$a(x) = \frac{2}{x} * p + \mathcal{O}(1/x^2)$$

Monodromy

$$\omega(x) = 1 + \frac{2}{x} \oint * p + \dots$$

Noether charge

$$Q = \frac{\sqrt{\lambda}}{2\pi} \oint * p$$

$$\omega(x) = \exp\left(-\frac{2\pi}{\sqrt{\lambda}x} Q + \dots\right)$$

Some more properties depending on G/H

Ansätze for Spectral Curves.

Consider $y(x) = (x^2 - 1)^2 q'(x)$ (to remove poles at $x = \pm 1$)

$y(x)$ has

- branch points $\sim \frac{1}{\sqrt{x-x_k}}$
- no branch points $\sim \sqrt{x-x_k}$
- no single poles or double poles.
- analytic otherwise

$\Rightarrow y(x)$ is an algebraic curve if finite genus.

Finite genus - finite number of branch cuts, "finite gap"

Ansatz for algebraic curve: $F_k(x)$ polynomials

$$F_N(x) y^N + F_{N-1}(x) y^{N-1} + F_{N-2}(x) y^{N-2} + \dots + F_0(x) = 0$$

Branch point $\frac{1}{\sqrt{x-x_k}}$ requires $F_N(x_k) = F_{N-1}(x_k) = 0$, $F_{N-2}(x_k) \neq 0$
 $F'_N(x_k) \neq 0$ ~~$F'_{N-1}(x_k) \neq 0$~~

$$F_N(x_k) \sim \prod (x - x_k), \quad F_{N-1}(x) = 0 \text{ typically.}$$

Restrict other coefficients of polynomials $F_k(x)$ by

- Absence of branch points $\sqrt{x-x_k}$ or poles
- Behaviour at $x = 0, \infty, \pm 1$
- A-cycles, mode numbers, fillings of cuts
- Further properties of sigma model on G/H .

Can fix all coefficients.

General Spectral Curves

Monodromy in N -dim representation of symmetry group

$\leadsto$ Spectral curve of degree N (N Riemann sheets)

Conjugation class of representation

$\leadsto$ Symmetries in curve, eg. $x \rightarrow 1/x$, $x \rightarrow -x$

For $AdS_5 \times S^5$ superstrings 4+4-dim repr. of $PSU(2,2|4)$

