

Counting Calabi-Yau black holes with mock modular forms

Boris Pioline



"Higher structures, moduli spaces and integrability"
Opening conference of CRC 1624
Hamburg, 1/4/2025

- A central goal for any theory of quantum gravity is to provide a **microscopic explanation** of the **thermodynamical entropy of black holes** in General Relativity [*Bekenstein'72, Hawking'74*]

$$S_{BH} = \frac{A}{4G_N}$$



$$S_{BH} \stackrel{?}{=} \log \Omega$$

- As shown by [*Strominger Vafa'96,...*], String Theory provides a quantitative description in the case of **BPS black holes in vacua with extended SUSY**: at weak string coupling, black hole micro-states arise as **bound states of D-branes** wrapped on cycles of the internal manifold.
- Besides confirming the consistency of string theory as a theory of quantum gravity, this has opened up many fruitful connections with mathematics.

- In the context of type IIA strings compactified on a Calabi-Yau three-fold X , BPS states are described mathematically by **stable objects in the derived category of coherent sheaves** $\mathcal{C} = D^b\text{Coh}X$. The Chern character $\gamma = (\text{ch}_0, \text{ch}_1, \text{ch}_2, \text{ch}_3)$ is identified as the electromagnetic charge, or D6-D4-D2-D0-brane charge.
- The problem becomes a question in **Donaldson-Thomas theory**: for fixed $\gamma \in K(X)$, compute the **generalized DT invariant** $\Omega_z(\gamma)$ counting **(semi)stable objects** of class γ , and determine its growth as $|\gamma| \rightarrow \infty$.
- Importantly, $\Omega_z(\gamma)$ depends on the moduli of X , or more generally on a choice of **Bridgeland stability condition** $z \in \text{Stab}\mathcal{C}$. The **chamber structure** is fairly simple for $X = T^6$ or $X = K3 \times T^2$, but very intricate for a general CY 3-fold.

Modularity of Donaldson-Thomas invariants

- Physical arguments predict that **suitable generating series of DT invariants** (those counting D4-D2-D0 bound states in a suitable chamber) should have specific **mock modular properties**. This gives very good control on their asymptotic growth, and allows to test agreement with the BH prediction $\Omega_z(\gamma) \simeq e^{S_{BH}(\gamma)}$.
- More precisely, these generating series are expected to be **mock modular**, similar to Ramanujan's mock theta series. The modular anomaly can be repaired by adding a universal non-holomorphic correction, determined recursively from generating series with lower D4-brane charge [*Alexandrov BP Manschot'16-20*].

In this talk, I will explain how to combine knowledge of standard **Gromov-Witten invariants** (counting curves in X) and **wall-crossing arguments** to **rigorously compute many DT invariants**, and check mock modularity to **high precision**

S. Alexandrov, S. Feyzbakhsh, A. Klemm, BP, T. Schimannek, arXiv:2301.08066

S. Alexandrov, S. Feyzbakhsh, A. Klemm, BP, arXiv:2312.12629

- 1 Reminder of enumerative invariants on CY3: GW, GV, DT, PT...
- 2 Mock modularity of D4-D2-D0 generating series
- 3 From rank 1 to rank 0 DT invariants, and back
- 4 Testing modularity on X_5 and other hypergeometric models
- 5 Conclusion and open problems

- Let X be a smooth, projective CY threefold. The **Gromov-Witten invariants** $\text{GW}_\beta^{(g)}$ count genus g curves Σ with class $\beta \in H_2(X, \mathbb{Z})$. They depend only on the symplectic structure (or Kähler moduli) of X and in general take rational values.
- Physically, they determine certain **higher-derivative couplings** in the low energy effective action, which depend only on the (complexified) Kähler moduli t and receive **worldsheet instanton corrections**: $F_g(t) = \sum_\beta \text{GW}_\beta^{(g)} e^{2\pi i t \cdot \beta}$ [Antoniadis Gava Narain Taylor'93]
- The first two F_0 and F_1 can be computed using **mirror symmetry**. **Holomorphic anomaly equations** along with suitable boundary conditions allow to determine $F_{g \geq 2}$ up to a certain genus g_{int} ($= 53$ for the quintic threefold X_5) [Bershadsky Cecotti Ooguri Vafa'93; Huang Klemm Quackenbush'06]

Gopakumar-Vafa invariants

- Gromov-Witten invariants turn out to be determined by a set of integer invariants $GV_{\beta}^{(g)}$ via [Gopakumar Vafa'98, Ionel Parker'13]

$$\sum_{g=0}^{\infty} \sum_{\beta} GW_{\beta}^{(g)} \lambda^{2g-2} e^{2\pi i t \cdot \beta} = \sum_{g=0}^{\infty} \sum_{k=1}^{\infty} \sum_{\beta} \frac{GV_{\beta}^{(g)}}{k} \left(2 \sin \frac{k\lambda}{2}\right)^{2g-2} e^{2\pi i k t \cdot \beta}$$

For $g = 0$, this reduces to [Candelas de la Ossa Greene Parkes'93]

$$GW_{\beta}^{(0)} = \sum_{k|\beta} \frac{1}{k^3} GV_{\beta/k}^{(0)}$$


- Physically, $GV_{\beta}^{(0)}$ counts **D2-D0 brane bound states** with D2 charge β , and arbitrary D0 charge n , while higher genus GV invariants keep track of their **angular momentum**.
- Importantly, $GV_{\beta}^{(g)}$ vanishes for large enough $g \geq g_{\max}(\beta)$ (Castelnuovo bound).

GV invariants and 5D rotating black holes

- Viewing type II string theory as M-theory on a circle, D2-branes lift to M2-branes wrapped on curve inside X , yielding **BPS black holes in $\mathbb{R}^{1,4}$** . These carry in general two angular momenta (j_L, j_R) .
- Tracing over j_R , the number of BPS states with $m = j_L^Z$ is

$$\Omega_{5D}(\beta, m) = \sum_{g=0}^{g_{\max}(\beta)} \binom{2g+2}{g+1+m} \text{GV}_{\beta}^{(g)}$$

Katz Klemm Vafa'99



- There is some numerical evidence that $\Omega(\beta, m) \sim e^{2\pi\sqrt{\beta^3 - m^2}}$ for large β keeping m^2/β^3 fixed, in agreement with the BH entropy of 5D black holes [*Klemm Marino Tavanfar'07*], with a transition to black rings at large angular momentum [*Halder Lin'23*].

Generalized Donaldson-Thomas invariants

- More generally, bound states of D6-D4-D2-D0 branes are described by stable objects in the **bounded derived category of coherent sheaves** $\mathcal{C} = D^b\text{Coh}(X)$ [Kontsevich'95, Douglas'01]. Objects are bounded complexes $E = (\cdots \rightarrow \mathcal{E}_{-1} \rightarrow \mathcal{E}_0 \rightarrow \mathcal{E}_1 \rightarrow \cdots)$ of coherent sheaves \mathcal{E}_k , graded by the total Chern character $\gamma(E) = \sum_k (-1)^k \text{ch } \mathcal{E}_k \in \Gamma$
- Stability depends on a choice of stability condition $\sigma = (Z, \mathcal{A})$, where the central charge $Z \in \text{Hom}(\Gamma, \mathbb{C})$ and the heart $\mathcal{A} \subset \mathcal{C}$ satisfy various axioms [Bridgeland 2007], in particular
 - 1 $\forall E \in \mathcal{A}, \text{Im}Z(E) \geq 0$
 - 2 $\forall E \in \mathcal{A}, \text{Im}Z(E) = 0 \Rightarrow \text{Re}Z(E) < 0$
- The **generalized Donaldson-Thomas invariant** $\Omega_\sigma(\gamma)$ is roughly the weighted Euler number of the moduli space $M_\sigma(\gamma)$ of **semi-stable objects** $E \in \mathcal{A}$ with $\text{ch } E = \gamma$, where semi-stability means that $\arg Z(E') \leq \arg Z(E)$ for any subobject $E' \subset E$.


- The space of stability conditions $\text{Stab } \mathcal{C}$ is a complex manifold of dimension $\dim K_{\text{num}}(X) = 2b_2(X) + 2$, unless it is empty [Bridgeland'07].
- Stability conditions in the vicinity of the large volume point can be constructed subject to a conjectural **Bogomolov-Gieseker-type inequality** introduced in [Bayer Macri Toda'11] – more on this later.
- The BMT inequality is very hard to prove for a general compact CY3, but has been proven for the quintic threefold X_5 [Li'18] and a couple of other examples [Koseki'20, Liu'21].

Generalized Donaldson-Thomas invariants

- $\Omega_\sigma(\gamma)$ may **jump** on co-dimension 1 walls in $\text{Stab } \mathcal{C}$ where some the central charge $Z(\gamma')$ of a subobject $E' \subset E$ becomes aligned with $Z(\gamma)$. The jump is governed by a universal **wall-crossing formula** [Joyce Song'08, Kontsevich Soibelman'08]. In simplest primitive case,

$$\Delta \Omega_\sigma(\gamma_1 + \gamma_2) = \langle \gamma_1, \gamma_2 \rangle \Omega_\sigma(\gamma_1) \Omega_\sigma(\gamma_2)$$

corresponding physically to the (dis)appearance of multi-centered black hole bound states [Denef Moore'07; Andriyash Denef Jafferis Moore'10; Manschot BP Sen'10]



- For $\gamma = (0, 0, \beta, n)$, $\Omega_\sigma(\gamma)$ coincides with $GV_\beta^{(0)}$ at large volume.

- For $\gamma = (-1, 0, \beta, -n)$ at large volume and B -field, stable objects have a much simpler mathematical description in terms of **stable pairs** $E : \mathcal{O}_X \xrightarrow{s} F$ [Pandharipande Thomas'07]:

- 1 F is a pure 1-dimensional sheaf with $\text{ch}_2 F = \beta$ and $\chi(F) = n$
- 2 the section s has zero-dimensional kernel

The **PT invariant** $\text{PT}(\beta, n)$ is defined as the (weighted) Euler characteristic of the corresponding moduli space.

- Since a single D6-brane lifts to a Taub-NUT space in M-theory, which is locally flat, one expects that PT invariants are computable from GV invariants [Dijkgraaf Vafa Verlinde'06].

GV invariants and D6-brane bound states

- More precisely, PT invariants are related to GV invariants by [Maulik Nekrasov Okounkov Pandharipande'06]

$$\sum_{\beta, n} \text{PT}(\beta, n) e^{2\pi i t \cdot \beta} q^n = \text{Exp} \left(\sum_{\beta, g} \text{GV}_{\beta}^{(g)} (\sqrt{q} - 1/\sqrt{q})^{2g-2} e^{2\pi i t \cdot \beta} \right)$$

where $\text{Exp}(f(q)) = \exp(\sum_{n \geq 1} f(q^n))$ is the plethystic exponential.

- Under this relation, the Castelnuovo bound $\text{GV}_{\beta}^{(g \geq g_{\max}(\beta))} = 0$ is mapped to $\text{PT}(\beta, n \leq 1 - g_{\max}(\beta)) = 0$
- The main interest in this talk will be on **rank 0 DT invariants** $\Omega(0, p, \beta, n)$ counting D4-D2-D0 brane bound states supported on a divisor \mathcal{D} with class $[\mathcal{D}] = p \in H_4(X, \mathbb{Z})$.

D4-D2-D0 indices as rank 0 DT invariants

- Viewing IIA= M/S^1 , D4-D2-D0 branes on \mathcal{D} arise from **M5-branes** wrapped on $\mathcal{D} \times S^1$. In the limit where S^1 is much larger than X , they are described by a **two-dimensional superconformal field theory** with (0, 4) SUSY. [*Maldacena Strominger Witten'97*]
- D4-D2-D0 indices occur as Fourier coefficients in the **elliptic genus** $\text{Tr}(-1)^F q^{L_0 - \frac{c_L}{24}} e^{2\pi i q_a z^a}$. If the SCFT has a discrete spectrum, after theta series decomposition with respect to the elliptic variables z^a , one obtains a vector-valued modular form

$$h_{\rho, \mu}(\tau) := \sum_n \bar{\Omega}(0, \rho, \mu, n) q^{n - \frac{\chi(\mathcal{D})}{24} + \frac{1}{2}\mu^2 - \frac{1}{2}\rho\mu}$$

where μ takes values in the finite discriminant group Λ^* / Λ associated to $\Lambda = (H_4(X, \mathbb{Z}), \kappa_{ab} := \kappa_{abc} \rho^c)$.

Modularity of rank 0 DT invariants

- When \mathcal{D} is **very ample** and **irreducible**, there are no walls extending to large volume, so the choice of chamber is irrelevant. The central charges are given by [Maldacena Strominger Witten'97]

$$\begin{cases} c_L = p^3 + c_2(TX) \cdot p = \chi(\mathcal{D}), \\ c_R = p^3 + \frac{1}{2}c_2(TX) \cdot p = 6\chi(\mathcal{O}_{\mathcal{D}}) \end{cases}$$

Cardy's formula predicts a growth $\Omega(0, p, \beta, n \rightarrow \infty) \sim e^{2\pi\sqrt{p^3 n}}$ in perfect agreement with Bekenstein-Hawking formula !

- Moreover, since the space of vector-valued weakly holomorphic modular form has finite dimension, the full series is completely determined by its **polar coefficients**, with $n + \frac{1}{2}\mu^2 - \frac{1}{2}p\mu < \frac{\chi(\mathcal{D})}{24}$. (Actually, the dimension can be smaller than the number of polar terms).

Mock modularity of rank 0 DT invariants

- When \mathcal{D} is reducible, the generating series $h_{p^a, \mu_a}(\tau)$ in a suitable ("large volume attractor") chamber is expected to be a **mock modular form of higher depth** [Alexandrov BP Manschot'16-20])
- Namely, there exists explicit, universal **non-holomorphic theta series** $\Theta_n(\{p_i\}, \tau, \bar{\tau})$ such that (ignoring the μ 's for simplicity)

$$\widehat{h}_p(\tau, \bar{\tau}) = h_p(\tau) + \sum_{p=\sum_{i=1}^{n \geq 2} p_i} \Theta_n(\{p_i\}, \tau, \bar{\tau}) \prod_{i=1}^n h_{p_i}(\tau)$$

transforms as a modular form. The completed series satisfy the **holomorphic anomaly equation**,

$$\partial_{\bar{\tau}} \widehat{h}_p(\tau, \bar{\tau}) = \sum_{p=\sum_{i=1}^{n \geq 2} p_i} \widehat{\Theta}_n(\{p_i\}, \tau, \bar{\tau}) \prod_{i=1}^n \widehat{h}_{p_i}(\tau, \bar{\tau})$$


Mock modularity of rank 0 DT invariants

- For binary splittings, this reduces to mock modular forms encountered in the study of BPS dyons in Type II on $K3 \times T^2$, or in heterotic string on T^6 [Dabholkar Murthy Zagier'12].
- The modular completion is constructed using similar ideas as in Zwegers's work on Ramanujan's mock theta series, namely replacing "step functions" with "generalized error functions" [Alexandrov Banerjee BP Manschot'16].
- Our derivation relied on the study of **instanton corrections to the QK metric** on the moduli space after compactifying on a circle, and implementing $SL(2, \mathbb{Z})$ symmetry manifest from $IIA/S^1 = M/T^2$. A nice spin off of earlier research on hypermultiplet moduli spaces !

Alexandrov Banerjee Persson BP Manschot Saueressig Vandoren, 2008-19

Non-holomorphic completion from Witten index

- Physically, the non-holomorphic corrections arise from the spectral asymmetry in the continuum of scattering states in the supersymmetric quantum mechanics of n BPS black holes.



BP 2015; Murthy BP 2018; BP Raj, in progress

Testing mock modularity for one-parameter models

- In the remainder of this talk, we shall test these modularity predictions for CY threefolds with Picard rank 1, by computing the first few coefficients in the q -expansion and determine the putative vector-valued (mock) modular form.
- This was first attempted by [*Gaiotto Strominger Yin '06-07*] for the quintic threefold X_5 and a few other hypergeometric models. They were able to guess the first few terms for unit D4-brane charge, and found a unique modular completion.
- We shall compute many terms rigorously, using recent results by [*Soheyla Fezbakhsh and Richard Thomas'20-22*] relating rank r DT invariants (including $r = 0$, counting D4-D2-D0 bound states) to PT invariants, hence to GV invariants.

Alexandrov, Fezbakhsh, Klemm, BP, Schimannek'23

From rank 1 to rank 0 DT invariants

- The key idea is to study wall-crossing in the space of Bridgeland stability conditions, away from the physical slice. For any $b + it \in \mathbb{H}$, consider the central charge


$$Z_{b,t}(E) = \frac{i}{6} t^3 \text{ch}_0^b(E) - \frac{1}{2} t^2 \text{ch}_1^b(E) - it \text{ch}_2^b(E) + \mathbf{0} \text{ch}_3^b(E)$$

with $\text{ch}_k^b(E) := \int_X H^{3-k} e^{-bH} \text{ch}(E)$. With a suitable choice of heart (defined by tilting with respect to the slope $\frac{\text{ch}_1^b(E)}{\text{rk}(E)}$), this defines a **weak** stability condition called **tilt-stability**.

- Note that $Z_{b,t}(E)$ is obtained from $Z^{\text{LV}}(E) = - \int_X e^{(b+it)H} \text{ch}(E)$ by setting by hand the coefficient of ch_3^b to $\mathbf{0}$. In fact, tilt-stability is the first step in constructing genuine stability conditions near the large volume point [*Bayer Macri Toda'11*]
- The KS/JS wall-crossing formulae still hold for such weak stability conditions.

Rank 0 DT invariants from GV invariants

- Tilt stability agrees with slope stability at large volume, but the chamber structure is much simpler: walls are **nested half-circles** in the Poincaré upper half-plane spanned by $z = b + i\frac{t}{\sqrt{3}}$.



- Importantly, for any tilt-semistable object E there is a **conjectural inequality** on Chern classes $C_i := \int_X \text{ch}_i(E) \cdot H^{3-i}$ [Bayer Macri Toda'11; Bayer Macri Stellari'16]

$$(C_1^2 - 2C_0C_2)\left(\frac{1}{2}b^2 + \frac{1}{6}t^2\right) + (3C_0C_3 - C_1C_2)b + (2C_2^2 - 3C_1C_3) \geq 0$$

Rank 0 DT invariants from GV invariants

- In particular, if the discriminant $\Delta(C)$ at $t = 0$ is positive, there exists an empty chamber ! $\Delta(\gamma)$ is quartic in the charges,

$$\Delta(C) = 8C_0C_2^3 + 6C_1^3C_3 + 9C_0^2C_3^2 - 3C_1^2C_2^2 - 18C_0C_1C_2C_3 \geq 0$$

- Remarkably, $\Delta(C)$ is proportional to (minus) the quartic invariant $I_4(Q)$ which determines the entropy $S_{BH} \sim \pi \sqrt{I_4(Q)}$ of single-centered black holes ! In particular, *an empty chamber exists whenever single-centered black hole are ruled out !*
- Consider an anti-D6-brane with charge $\gamma = (-1, 0, \beta, -n)$ such that $\Delta(C) > 0$. By studying wall-crossing between the empty chamber where $\Omega_{b,t}(\gamma) = 0$ and the large volume chamber where $\Omega_{b,t}(\gamma) = \text{PT}(\beta, m)$, one can extract the indices of the D4-D2-D0 branes emitted at each wall !

A new explicit formula (S. Feyzbakhsh'23)

Theorem Let (X, H) be a smooth polarised CY threefold with $\text{Pic}(X) = \mathbb{Z}.H$ satisfying the BMT conjecture. There is $f(x)$ such that

- If $\frac{m}{\beta.H} < f\left(\frac{\beta.H}{H}\right)$ then the stable pair invariant $\text{PT}(\beta, m) =$

$$\sum_{(m', \beta')} (-1)^{\chi_{m', \beta'}} \chi_{m', \beta'} \text{PT}(\beta', m') \Omega\left(0, 1, \frac{H^2}{2} - \beta' + \beta, \frac{H^3}{6} + m' - m - \beta'.H\right)$$

where $\chi_{m', \beta'} = \beta.H + \beta'.H + m - m' - \frac{H^3}{6} - \frac{1}{12}c_2(X).H$.

- The sum runs over $(\beta', m') \in H_2(X, \mathbb{Z}) \oplus H_0(X, \mathbb{Z})$ such that

$$\begin{aligned} 0 \leq \beta'.H &\leq \frac{H^3}{2} + \frac{3mH^3}{2\beta.H} + \beta.H \\ -\frac{(\beta'.H)^2}{2H^3} - \frac{\beta'.H}{2} \leq m' &\leq \frac{(\beta.H - \beta'.H)^2}{2H^3} + \frac{\beta.H + \beta'.H}{2} + m \end{aligned}$$

In particular, $\beta'.H < \beta.H$.

Corollary (Castelnuovo bound): $\text{PT}(\beta, m) = 0$ unless $m \geq -\frac{(\beta.H)^2}{2H^3} - \frac{\beta.H}{2}$

Modularity for one-modulus compact CY

- Using the theorem above and known GV invariants, we could compute a large number of coefficients in the generating series of Abelian (=unit D4-brane charge) rank 0 DT invariants in **one-parameter hypergeometric threefolds**, including the quintic X_5 .
- In all cases (except $X_{3,2,2}$, $X_{2,2,2,2}$ where current knowledge of GV invariants is insufficient), we found a linear combination of the following vv modular forms matching all computed coeffs:

$$\frac{E_4^a E_6^b}{\eta^{4\kappa+c_2}} D^\ell(\vartheta_\mu^{(\kappa)}) \quad \text{with} \quad \vartheta_\mu^{(\kappa)} = \sum_{k \in \mathbb{Z} + \frac{\mu}{\kappa} + \frac{1}{2}} q^{\frac{1}{2}\kappa k^2}, \quad \kappa := H^3$$

where $D = 2\pi i \partial_\tau - \frac{w}{12} E_2$, and $4a + 6b + 2\ell - 2\kappa - \frac{1}{2}c_2 = -2$.

Modularity for one-modulus compact CY

X	χ_X	κ	$c_2(TX)$	$\chi(\mathcal{O}_D)$	n_1	C_1
$X_5(1^5)$	-200	5	50	5	7	0
$X_6(1^4, 2)$	-204	3	42	4	4	0
$X_8(1^4, 4)$	-296	2	44	4	4	0
$X_{10}(1^3, 2, 5)$	-288	1	34	3	2	0
$X_{4,3}(1^5, 2)$	-156	6	48	5	9	0
$X_{4,4}(1^4, 2^2)$	-144	4	40	4	6	1
$X_{6,2}(1^5, 3)$	-256	4	52	5	7	0
$X_{6,4}(1^3, 2^2, 3)$	-156	2	32	3	3	0
$X_{6,6}(1^2, 2^2, 3^2)$	-120	1	22	2	1	0
$X_{3,3}(1^6)$	-144	9	54	6	14	1
$X_{4,2}(1^6)$	-176	8	56	6	15	1
$X_{3,2,2}(1^7)$	-144	12	60	7	21	1
$X_{2,2,2,2}(1^8)$	-128	16	64	8	33	3

Modular predictions for the quintic threefold

- Using known $GV_{\beta}^{(g \leq 53)}$ we can compute more than 20 terms:

$$h_0 = q^{-\frac{55}{24}} \left(\underline{5 - 800q + 58500q^2 + 5817125q^3 + 75474060100q^4} \right. \\ \left. + 28096675153255q^5 + 3756542229485475q^6 \right. \\ \left. + 277591744202815875q^7 + 13610985014709888750q^8 + \dots \right),$$

$$h_{\pm 1} = q^{-\frac{55}{24} + \frac{3}{5}} \left(\underline{0 + 8625q - 1138500q^2 + 3777474000q^3} \right. \\ \left. + 3102750380125q^4 + 577727215123000q^5 + \dots \right)$$

$$h_{\pm 2} = q^{-\frac{55}{24} + \frac{2}{5}} \left(\underline{0 + 0q - 1218500q^2 + 441969250q^3 + 953712511250q^4} \right. \\ \left. + 217571250023750q^5 + 22258695264509625q^6 + \dots \right)$$

Modular predictions for the quintic threefold

- The space of v.v modular forms has dimension 7. Remarkably, all terms above are reproduced by *[Gaiotto Strominger Yin'06]*

$$h_{\mu} = \frac{1}{\eta^{55+15}} \left[-\frac{222887E_4^8+1093010E_4^5E_6^2+177095E_4^2E_6^4}{35831808} + \frac{25(458287E_4^6E_6+967810E_4^3E_6^3+66895E_6^5)}{53747712} D + \frac{25(155587E_4^7+1054810E_4^4E_6^2+282595E_4E_6^4)}{8957952} D^2 \right] \vartheta_{\mu}^{(5)}$$

- Physically, polar coefficients are expected arise as **bound states of D6-brane and anti D6-branes** *[Denef Moore'07]*. Indeed, they are **often** consistent with the naive ansatz *[Alexandrov Gaddam Manschot BP'22]*

$$\Omega(0, 1, \beta, n) = \pm(\chi(\mathcal{O}_{\mathcal{D}}) - \beta.H - n) DT(\beta, n) PT(0, 0)$$

but deviations do occur ! *[Collinucci Wyder'08, van Herck Wyder'09]*

Mock modularity for non-Abelian D4-D2-D0 indices

- For D4-D2-D0 indices with $N = 2$ units of D4-brane charge, $\{h_{2,\mu}, \mu \in \mathbb{Z}/(2\kappa\mathbb{Z})\}$ should transform as a **vector-valued mock modular form** with modular completion

$$\widehat{h}_{2,\mu}(\tau, \bar{\tau}) = h_{2,\mu}(\tau) + \sum_{\mu_1, \mu_2=0}^{\kappa-1} \delta_{\mu_1+\mu_2-\mu}^{(\kappa)} \Theta_{\mu_2-\mu_1+\kappa}^{(\kappa)} h_{1,\mu_1} h_{1,\mu_2}$$

where (denoting $\beta(x) = 2|x|^{-1/2}e^{-\pi x} - 2\pi\text{Erfc}(\sqrt{\pi|x|})$)

$$\Theta_{\mu}^{(\kappa)}(\tau, \bar{\tau}) := \frac{(-1)^{\mu}}{8\pi} \sum_{k \in 2\kappa\mathbb{Z} + \mu} |k| \beta\left(\frac{\tau_2 k^2}{\kappa}\right) e^{-\frac{\pi i \tau}{2\kappa} k^2},$$

$$\partial_{\bar{\tau}} \Theta_{\mu}^{(\kappa)} = \frac{(-1)^{\mu} \sqrt{\kappa}}{16\pi i \tau_2^{3/2}} \sum_{k \in 2\kappa\mathbb{Z} + \mu} e^{-\frac{\pi i \bar{\tau}}{2\kappa} k^2} k^2$$

Mock modularity for non-Abelian D4-D2-D0 indices

- Suppose there exists a holomorphic function $g_\mu^{(\kappa)}$ such that $\Theta_\mu^{(\kappa)} + g_\mu^{(\kappa)}$ transforms as a $\nu\nu$ modular form. Then

$$\tilde{h}_{2,\mu}(\tau, \bar{\tau}) = h_{2,\mu}(\tau) - \sum_{\mu_1, \mu_2=0}^{\kappa-1} \delta_{\mu_1+\mu_2-\mu}^{(\kappa)} g_{\mu_2-\mu_1+\kappa}^{(\kappa)} h_{1,\mu_1} h_{1,\mu_2}$$

will be an ordinary weak holomorphic $\nu\nu$ modular form, hence uniquely determined by its polar part.

- For $\kappa = 1$, the series $\Theta_\mu^{(1)}$ is the one appearing in the modular completion of the generating series of **Hurwitz class numbers** [Hirzebruch Zagier 1973] (or **rank 2 Vafa-Witten invariants on \mathbb{P}^2**)

$$H_0(\tau) = -\frac{1}{12} + \frac{1}{2}q + q^2 + \frac{4}{3}q^3 + \frac{3}{2}q^4 + \dots$$
$$H_1(\tau) = q^{\frac{3}{4}} \left(\frac{1}{3} + q + q^2 + 2q^3 + q^4 + \dots \right)$$

Thus we can choose $g_\mu^{(1)} = H_\mu(\tau)$.

Mock modularity for non-Abelian D4-D2-D0 indices

X	χ_X	κ	c_2	$\chi(\mathcal{O}_{2D})$	n_2	C_2
$X_5(1^5)$	-200	5	50	15	36	1
$X_6(1^4, 2)$	-204	3	42	11	19	1
$X_8(1^4, 4)$	-296	2	44	10	14	1
$X_{10}(1^3, 2, 5)$	-288	1	34	7	7	0
$X_{4,3}(1^5, 2)$	-156	6	48	16	42	0
$X_{4,4}(1^4, 2^2)$	-144	4	40	12	25	1
$X_{6,2}(1^5, 3)$	-256	4	52	14	30	1
$X_{6,4}(1^3, 2^2, 3)$	-156	2	32	8	11	1
$X_{6,6}(1^2, 2^2, 3^2)$	-120	1	5	2	5	0
$X_{3,3}(1^6)$	-144	9	54	21	78	3
$X_{4,2}(1^6)$	-176	8	56	20	69	3
$X_{3,2,2}(1^7)$	-144	12	60	26	117	0
$X_{2,2,2,2}(1^8)$	-128	16	64	32	185	4

Mock modularity for non-Abelian D4-D2-D0 indices

- For X_{10} , we computed the 7 polar terms + 4 non-polar terms and found a unique mock modular form reproducing this data:

$$h_{2,\mu} = \frac{5397523E_4^{12} + 70149738E_4^9E_6^2 - 12112656E_4^6E_6^4 - 61127530E_4^3E_6^6 - 2307075E_6^8}{46438023168\eta^{100}} \mathcal{V}_\mu^{(1,2)} \\ + \frac{-10826123E_4^{10}E_6 - 14574207E_4^7E_6^3 + 20196255E_4^4E_6^5 + 5204075E_4E_6^7}{1934917632\eta^{100}} D\mathcal{V}_\mu^{(1,2)} \\ + (-1)^{\mu+1} H_{\mu+1}(\tau) h_1(\tau)^2$$


with $h_1 = \frac{203E_4^4 + 445E_4E_6^2}{216\eta^{35}} = q^{-\frac{35}{24}} (3 - 575q + \dots)$, leading to integer DT invariants

$$h_{2,0}^{(\text{int})} = q^{-\frac{19}{6}} \left(\underline{7 - 1728q + 203778q^2 - 13717632q^3} - 23922034036q^4 + \dots \right)$$


$$h_{2,1}^{(\text{int})} = q^{-\frac{35}{12}} \left(\underline{-6 + 1430q - 1086092q^2 + 208065204q^3} + \dots \right)$$

- Similar results for X_8 [*S. Alexandrov, S. Feyzbakhsh, A. Klemm'23*]

Computing the leading term in $h_{2,0}$ for X_{10}



Quantum geometry from stability and modularity



Alexandrov Feyzbakhsh Klemm BP Schimannek'23

Quantum geometry from stability and modularity

X	χ_X	κ	type	g_{integ}	$g_{\text{mod}}^{(1)}$	$g_{\text{mod}}^{(2)}$	g_{avail}
$X_5(1^5)$	-200	5	F	53	69	80	64
$X_6(1^4, 2)$	-204	3	F	48	66	84	48
$X_8(1^4, 4)$	-296	2	F	60	84	112	66
$X_{10}(1^3, 2, 5)$	-288	1	F	50	70	95	72
$X_{4,3}(1^5, 2)$	-156	6	F	20	24		24
$X_{6,4}(1^3, 2^2, 3)$	-156	2	F	14	17		17
$X_{6,6}(1^2, 2^2, 3^2)$	-120	1	K	18	22		26
$X_{4,4}(1^4, 2^2)$	-144	4	K	26	34		34
$X_{3,3}(1^6)$	-144	9	K	29	33		33
$X_{4,2}(1^6)$	-176	8	C	50	66		64
$X_{6,2}(1^5, 3)$	-256	4	C	63	78		49

<http://www.th.physik.uni-bonn.de/Groups/Klemm/data.php>

Summary and open questions

- We provided overwhelming evidence that D4-D2-D0 indices exhibit mock modular properties. Where does it come from mathematically ? Is there some VOA acting on the cohomology of moduli space of stable objects, à la *[Nakajima'94]* ?
- Can one test modularity in multi-parameter models, for example in genus-one fibrations or K3-fibrations ? Can one follow D4-D2-D0 invariants through extremal transitions ?
- Similar wall-crossing arguments also allow to compute higher rank DT invariants. Is there some higher rank version of *[MNOP'03]* ?
- A long-standing problem: incorporate NS5-instanton corrections to the QK metric on hypermultiplet moduli space, consistently with S-duality, beyond the linear analysis of *[Alexandrov Persson BP'10]*.
- Thanks for your attention !

Back up slide: Modularity from geometry

- While modularity of D4-D2-D0 invariants is clear physically from the M5-brane picture, its mathematical origin is in general mysterious.
- When X admits a **K3-fibration**, using the relation to **Noether-Lefschetz invariants** one can show that modularity holds for **vertical** D4-brane charge. The modular anomaly disappears entirely due to $\kappa_{ab}p^b = 0$. [*Bouchard Creutzig Diaconescu Doran Quigley Sheshmani'16; Doran BP Schimannek'24*]
- Similarly, when X admits a **genus-one fibration**, one can relate D4-D2-D0 invariants for a D4-brane wrapping the fiber to GW invariants via monodromy. Generating series of GW invariants are quasi-modular forms, consistent with $\kappa_{ab}p^a p^b = 0$. [*Klemm Manschot Wotschke'12; BP Schimannek, to appear.*]