

Asymptotics of higher genus string integrands

Boris Pioline



“String theory from a worldsheet perspective”,
GGI, FLorence, 16/4/2019

*based on 1712.06135, 1806.02691, 1810.11343
in collaboration with Eric d'Hoker and Michael B. Green*

Modular graph functions beyond genus one I

- In perturbative closed string theory, scattering amplitudes of N external states at h loop involve an integral over the moduli space $\mathcal{M}_{h,N}$ of genus h curves Σ with N punctures z_1, \dots, z_N . At fixed complex structure, this reduces to an integral over N copies of Σ .
- At genus one, the low energy expansion of the resulting integrand produces an infinite family of real-analytic modular forms labelled by certain graphs, known as **modular graph functions**. They exhibit remarkable asymptotics near the cusp (*finite Laurent polynomial plus exponential corrections*) and transcendental properties.
- My goal will be to describe some hints of a similar structure at higher genus, primarily for $N = 4, g = 2$.

String integrand I

- Recall that the genus h contribution to N -point scattering amplitude is of the form

$$\mathcal{A}_h(\{k_i, \epsilon_i\}) = \int_{\mathcal{M}_{h,N}} \mathcal{I}_h(\{k_i, \epsilon_i\})$$

where $\mathcal{I}_h(\{k_i, \epsilon_i\}) \sim \langle \prod_{i=1}^N V_{k_i, \epsilon_i}(z_i) \prod_{j=1}^{3h-3} |(b, \mu)|^2 \rangle$ is a correlator in the worldsheet conformal field theory.

- The moduli space $\mathcal{M}_{h,N}$ is fibered over $\mathcal{M}_h \equiv \mathcal{M}_{h,0}$, with fiber $\Sigma_1 \times \cdots \times \Sigma_N$. The vertex operators V_{k_i, ϵ_i} are $(1, 1)$ -forms over Σ_i , while the ghost part produces a volume form on the base \mathcal{M}_h .
- I will focus on $1 \leq h \leq 3$, where \mathcal{M}_h is isomorphic to a fundamental domain \mathcal{F}_h for the action of $Sp(h, \mathbb{Z})$ on the Siegel upper half-plane \mathcal{H}_h (away from some divisors).

- In the RNS formalism, \mathcal{I}_h originates from an integrand \mathcal{J}_h over the moduli space of super Riemann curves $\mathfrak{M}_{h,N}$, after summing over spin structures and integrating over fermionic moduli. There is no canonical projection $\mathfrak{M}_{h,N} \rightarrow \mathcal{M}_{h,n}$, leading to total derivative ambiguities [Donagi Witten'13]. State of the art: $h = 2, N = 4$ [D'Hoker Phong '05]
- In Berkovits' pure spinor formalism, b is a composite field involving powers of $(\lambda\bar{\lambda})^{-1}$, leading to potential divergences from the region $\lambda\bar{\lambda} \rightarrow 0$. State of the art: idem, with partial results for $h = 2, N = 5$ [Gomez Mafrá Schlotterer '15] and $h = 3, N = 4$ [Gomez Mafrá '13]

Modular graph functions beyond genus one I

- Although we do not know the integrand \mathcal{I}_h in general, for type II strings on T^d we expect a result of the form

$$\mathcal{I}_h = \delta\left(\sum k_i\right) Z_{KN}(\{k_i\}) Z_{T^d} P_h(\{k_i, \epsilon_i, z_i\}) \mu_h$$

where

- $Z_{KN}(\{k_i, z_i\}) = \exp(\sum_{i < j} s_{ij} G(z_i, z_j))$ is the **Koba-Nielsen factor**, with $s_{ij} = -\frac{\alpha'}{2} k_i \cdot k_j$ the Mandelstam variables and $G(z_i, z_j)$ the **scalar Green function**.
- Z_{T^d} is the genus- h **Siegel theta series** for a lattice of signature (d, d)
- $P_h(\{k_i, \epsilon_i, z_i\})$ is a **(1,1)-form** in each position z_i , and a Lorentz invariant homogeneous polynomial in k_i, ϵ_i
- μ_h is the Siegel volume form over $\mathcal{M}_h \simeq \mathcal{F}_h$

Canonical forms on compact curves I

- Let A_I, B_I ($I = 1 \dots h$) be a canonical basis of $H_1(\Sigma)$, such that $A_I \cap A_J = B_I \cap B_J = 0, A_I \cap B_J = \delta_{IJ}$. Choose ω_I a basis of holomorphic differentials on Σ such that $\int_{A_I} \omega_J = \delta_{IJ}$. The period matrix $\tau_{IJ} = \int_{B_I} \omega_J$ satisfies $\tau > 0$, hence $\tau \in \mathcal{H}_h$.
- Under $Sp(h, \mathbb{Z})$ change of basis, $\tau \mapsto (a\tau + b)(c\tau + d)^{-1}$. The Siegel volume form $\mu_h = \frac{\prod_{I < J} |d\tau_{IJ}|^2}{(\det \text{Im}\tau)^{h+1}}$ is invariant.
- An example of $(1,1)$ form on Σ_i is the canonical Kähler form $\kappa(z_i) = \frac{i}{2h} \text{Im}\tau^{IJ} \omega_I(z_i) \bar{\omega}_J(z_i)$.
- Examples of $(1,1)$ forms on $\Sigma_i \times \Sigma_j$ are $\kappa(z_i)\kappa(z_j)$ and $|\nu(z_i, z_j)|^2$ with $\nu(z_i, z_j) = \text{Im}\tau^{IJ} \omega_I(z_i) \bar{\omega}_J(z_j)$. Another example on h copies of Σ is $|\Delta(z_1, \dots, z_h)|^2$ where $\Delta(z_1, \dots, z_h) = \epsilon^{I_1 \dots I_h} \omega_{I_1}(z_1) \dots \omega_{I_h}(z_h)$ is a $(1,0)$ -form in each variable.

Canonical forms on compact curves II

- The Arakelov Green function $G(z, w)$ is a symmetric function on $\Sigma \times \Sigma$ defined by

$$\begin{aligned}\partial_z \bar{\partial}_z G(z, w) &= -\pi \delta^{(2)}(z, w) + \pi \kappa(z), \\ \int_{\Sigma} G(z, w) \kappa(w) &= 0,\end{aligned}$$

The r.h.s. integrates to zero thanks to $\int_{\Sigma} \kappa = 1$, allowing G to be globally well-defined.

- In genus one, setting $v = \int_z^w \omega = \alpha + \beta\tau$,

$$G(z, w) = -\log \left| \frac{\vartheta_1(v, \tau)}{\eta} \right|^2 + \frac{2\pi}{\tau_2} (\operatorname{Im} v)^2 = \sum'_{(m, n) \in \mathbb{Z}^2} \frac{\tau_2 e^{2\pi i(m\beta - n\alpha)}}{\pi |m + n\tau|^2}$$

Genus-one modular graph functions I

- After Taylor expanding in powers of s_{ij} , and integrating over the positions z_i of the vertex operators, the integrand is a linear combination of **genus one modular graph functions**, of the type

$$\mathcal{B}_\Gamma(\tau) = \int_{\Sigma^N/\Sigma} \prod_{(i,j) \in \Gamma} G(z_i, z_j) \prod_i \kappa(z_i) \quad \kappa(z) = \frac{dzd\bar{z}}{2i \operatorname{Im}\tau}$$

where (i, j) runs over the edges of the graph Γ (possibly dressed with derivatives wrt z_i) [*d'Hoker Green Gurdogan Vanhove '15*]

- By construction, \mathcal{B}_Γ are real analytic modular functions. e.g.

$$D_k(\tau) = \int_{\Sigma} G(v)^k \kappa(v) \quad \img alt="Diagram of a genus-one graph with two vertices and multiple edges." data-bbox="625 687 761 745"/>$$

gives E_2 for $k = 2$, $E_3 + \zeta(3)$ for $k = 3$.

Genus-one modular graph functions II

- Modular graph functions satisfy an intricate set of algebraic and differential equations. Their expansion near the cusp $\tau \rightarrow i\infty$ has the remarkable form

$$\mathcal{B}_\Gamma(\tau) = \sum_{n=w}^{1-w} a_n (\pi\tau_2)^n + \mathcal{O}(e^{-2\pi\tau_2})$$

where w is the number of edges (i.e. degree in G). The coefficients a_n are **single valued elliptic multizeta values** of transcendental degree $w - n$.

*d'Hoker Green Gurdogan Vanhove Zerbini Kaidi Basu Kleinschmidt Verschinin Gerken
Schlotterer Duke ...*

Genus-one modular graph functions III

- After integrating over $\tau \in \mathcal{F}_1$, the coefficient of the effective interaction $\sigma_2^p \sigma_3^q \mathcal{R}^4$ at one-loop (where $\sigma_n = s^n + t^n + u^n$) is given by a **regularized theta lifting**,

$$\mathcal{E}_{(p,q)}^{(1,d)} = \sum_{\Gamma} \alpha_{\Gamma} \int_{\mathcal{F}_1(L)} \mathcal{B}_{\Gamma}(\tau) Z_{T^d}(G_{ij}, B_{ij}; \tau) \mu_1$$

where G_{ij}, B_{ij} parametrize the Narain moduli space $\frac{O(d,d)}{O(d) \times O(d)}$ and $\mathcal{F}_1(L) = \mathcal{F}_1 \cap \{\tau_2 < L\}$ is the truncated fundamental domain.

- The powerlike terms in $\mathcal{B}_{\Gamma}(\tau)$ are responsible for infrared divergences as $L \rightarrow \infty$. The full amplitude including non-analytic terms is finite when $D > 4$. More on this later.

Higher genus modular graph functions I

- For $h = 2$, $N = 4$, the precise integrand is

d'Hoker Phong '05

$$\mathcal{I}_2 = \delta\left(\sum k_i\right) \mathcal{R}^4 |\mathcal{Y}|^2 Z_{KN} Z_{T^d} \det \text{Im} \tau \mu_2$$

where

$$\mathcal{Y} = (t - u) \Delta(z_1, z_2) \Delta(z_3, z_4) + \text{perm}$$

$$Z_{KN} = e^s [G(z_1, z_2) + G(z_3, z_4)] + t [G(z_1, z_4) + G(z_2, z_3)] + u [G(z_1, z_3) + G(z_2, z_4)]$$

- At leading order, $Z_{KN} \simeq 1$ and $\int_{\Sigma^4} |\Delta(1, 2) \Delta(3, 4)|^2 \propto 1 / \det \text{Im} \tau$.
The coefficient of the $\nabla^4 \mathcal{R}^4$ interaction at genus 2 is then

$$\mathcal{E}_{(1,0)}^{(d,2)} = \frac{\pi}{2} \int_{\mathcal{F}_2(L)} Z_{T^d} \mu_2 \propto E_{\frac{d-3}{2} \Lambda_2}^{O(d,d)}$$

Higher genus modular graph functions II

- At next-to-leading order, one power of $G(z_1, z_2)$ comes down from Z_{KN} . The integral over z_3, z_4 is easy. The coefficient of the $\nabla^6 \mathcal{R}^4$ interaction at genus-two is [d'Hoker Green'13]

$$\mathcal{E}_{(0,1)}^{(d,2)} = \pi \int_{\mathcal{F}_2(L)} \varphi_{KZ} Z_{T^d} \mu_2$$

where φ_{KZ} is the **Kawazumi-Zhang invariant**, defined by

$$\varphi_{KZ} = -\frac{1}{4} \int_{\Sigma \times \Sigma} |\nu(z_1, z_2)|^2 G(z_1, z_2)$$

where $\nu(z_1, z_2) = \text{Im} \tau^{\mu} \omega_I(z_1) \bar{\omega}_J(z_2)$.

- This is the simplest genus-two modular graph function, associated to the graph $1 \text{---} 2$ (but the measure need to be specified)

Higher genus modular graph functions III

- Using standard theory of complex structure deformations, one may show that φ_{KZ} is an eigenmode of the Laplacian on \mathcal{M}_2 ,

$$[\Delta_{Sp(4)} - 5] \varphi_{KZ} = -2\pi \det \text{Im} \tau \delta^{(2)}(\nu)$$

which makes it easy to compute $\int_{\mathcal{F}_2} \varphi_{KZ} \mu_2$ [d'Hoker Green BP Russo'14]

- Integrating against Z_{T^d} , it follows that [BP Russo'15]

$$[\Delta_{O(d,d)} - (d+2)(5-d)] \mathcal{E}_{(0,1)}^{(d,2)} = - \left[\mathcal{E}_{(0,0)}^{(d,1)} \right]^2 \propto \left[E_{\frac{d-2}{2}\Lambda_1}^{O(d,d)} \right]^2$$

hence $\mathcal{E}_{(0,1)}^{(d,2)}$ is not an Eisenstein series.

- Combined with information about asymptotics (see below), this shows that φ_{KZ} can be obtained as a **Borcherds' type theta lift** of the weak Jacobi form θ_1^2/η^6 from $SL(2)$ to $SO(3,2) = Sp(4)$. This gives access to the full Fourier expansion [BP'15].


Higher genus modular graph functions IV

- At NNLO, one needs to bring down two Green functions from Z_{KN} . The coefficient of the $\nabla^8 \mathcal{R}^4$ interaction at genus two is then

$$\mathcal{E}_{(2,0)}^{(d,2)} = \frac{\pi}{64} \int_{\mathcal{F}_2(L)} \mathcal{B}_{(2,0)} Z_{Td} \mu_2$$

where $\mathcal{B}_{(2,0)}$ is a linear combination of three genus-two modular graph functions,

$$\mathcal{B}_{(2,0)} = \int_{\Sigma^4} \frac{|\Delta(1,3)\Delta(2,4)|^2}{(\det \text{Im}\tau)^2} \left\{ \begin{array}{l} G(1,2)^2 - 2G(1,2)G(1,4) \\ + G(1,2)G(3,4) \end{array} \right\}$$



Higher genus modular graph functions V

- At NNNLO order, the $\nabla^{10}\mathcal{R}^4$ interaction involves modular graph functions with 4 vertices and 3 edges, etc.
- For higher point scattering amplitudes, one expects a similar hierarchy of integrands with N vertices and $k \geq 0$ edges, but the integration measure on Σ^N is not known for $N > 5$. Presumably it should be built from holomorphic one-forms ω_I , derivatives $\partial G(z, w)$ and their complex conjugate.
- For $h = 3$, $N = 4$, the leading $\nabla^6\mathcal{R}^4$ integrand computed in pure spinor formalism is given by *[Gomez Mafra 2013]*

$$\mathcal{E}_{(0,1)}^{(3,d)} = \frac{5}{16} \int_{\mathcal{F}_3(L)} Z_{T^d} \mu_3 \propto E_{\frac{d-4}{2}\Lambda_3}^{O(d,d)}$$

but the $\nabla^8\mathcal{R}^4$ integrand remains to be computed *[Gomez Mafra 20??]*.

Arakelov-Green function at higher genus I

- Recall that the **Arakelov Green function** $G(z, w)$ is defined by

$$\partial_z \bar{\partial}_z G(z, w) = -\pi \delta^{(2)}(z, w) + \pi \kappa(z), \quad \int_{\Sigma} G(z, w) \kappa(w) = 0$$

- After cutting Σ along $2h$ cycles to make it simply connected, $G(z, w)$ can be expressed in terms of the **string Green function**

$$\mathcal{G}(z, w) = -\log |E(z, w)|^2 + 2\pi i \operatorname{Im} v_I \operatorname{Im} \tau^{IJ} \operatorname{Im} v_J$$

where $E(z, w)$ is the **prime form** and $v_I = \int_z^w \omega_I$. $\mathcal{G}(z, w)$ satisfies

$$\partial_z \bar{\partial}_z \mathcal{G}(z, w) = -\pi \delta^{(2)}(z, w) + \pi h \kappa(z)$$


- $G(z, w)$ is obtained from $\mathcal{G}(z, w)$ via

$$G(z, w) = \mathcal{G}(z, w) - \gamma(z) - \gamma(w) + \gamma_0$$

with $\gamma(z) = \int_{\Sigma} \mathcal{G}(z, w) \kappa(w)$ and $\gamma_0 = \int_{\Sigma} \gamma(z) \kappa(z)$.

Arakelov-Green function at higher genus II

- Using the theta series representation of the prime form $E(z, w)$, one can obtain accurate asymptotics of $G(z, w)$ near boundaries of moduli space.
- In the non-separating degeneration, a genus $h + 1$ curve Σ_{h+1} degenerates into a genus h curve Σ_h with two marked points ρ_a, ρ_b , linked by a thin handle of proper length $t = \det \operatorname{Im} \tau_{h+1} / \det \operatorname{Im} \tau$.




Arakelov-Green function at higher genus III

- Theorem [Wentworth'91, d'H G BP '17]: In the limit $t \rightarrow \infty$,

$$\begin{aligned} G_{h+1}(z, w) &= \frac{\pi t}{3(h+1)^2} + \left[G_h(z, w) + \frac{G_h(p_a, p_b)}{(h+1)^2} \right. \\ &\quad \left. - \frac{1}{2(h+1)} (G_h(z, p_a) + G_h(z, p_b) + G_h(w, p_a) + G_h(w, p_b)) \right] \\ &\quad + \frac{1}{8\pi t} \left[\frac{f(z)^2 + f(w)^2}{(h+1)^2} - 2f(z)f(w) - \frac{2h}{(h+1)^2} \int_{\Sigma_h} f(x)^2 \kappa(x) \right] + \mathcal{O}(e^{-2\pi t}) \end{aligned}$$

- Here $f(z) := G_h(z, p_b) - G_h(z, p_a)$ is a single-valued real function on Σ_h , interpolating from $f(p_a) = -2\pi t$ to $f(p_b) = +2\pi t$.



Asymptotics of higher genus integrands I

- This gives access to the asymptotics of genus-two string invariants, e.g. for $\tau = \begin{pmatrix} \rho & v \\ v & \sigma_1 + i(t + \frac{v_2^2}{\rho_2}) \end{pmatrix}$, $t \rightarrow \infty$,

$$\varphi_{KZ}(\tau) = \frac{\pi t}{6} + \frac{1}{2}g_1(v) + \frac{5}{4\pi t}(g_2(v) - E_2) + \mathcal{O}(e^{-2\pi t})$$

where $g_1(v, \rho)$ is the genus-one Arakelov Green function, and $g_2(v, \rho) = \int_{\Sigma_1} g_1(v - w)g_1(w) \kappa(w)$.

- These asymptotics can be derived independently from the theta lift representation of φ_{KZ} . Different powers of t come from different orbits [BP'15]. The eigenvalue equation $[\Delta_{Sp(4)} - 5]\varphi_{KZ} = 0$ holds order by order in t .

Asymptotics of higher genus integrands II

- More generally, the w -th term in the Taylor expansion of the $N = 4, h = 2$ amplitude (a homogeneous polynomial of degree $w + 2$ in s_{ij})

$$\mathcal{B}_w(\mathbf{s}_{ij}; \tau) := \int_{\Sigma^4} \frac{|\mathcal{Y}|^2}{(\det \operatorname{Im} \tau)^2} \left(\sum_{1 \leq i < j \leq 4} s_{ij} G(z_i, z_j) \right)^w$$

has an asymptotic expansion of the form [d'H G BP '17]

$$\mathcal{B}_w(\mathbf{s}_{ij}; \tau) = \sum_{k=-w}^w b_w^{(k)}(\mathbf{s}_{ij}; \nu, \rho) (\pi t)^k + \mathcal{O}(e^{-2\pi t})$$

as $t \rightarrow \infty$. Here $b_w^{(k)}(\mathbf{s}_{ij}; \nu, \rho)$ are generalized genus-one modular graph functions.

Asymptotics of higher genus integrands III

- For $w = 2$, corresponding to $\nabla^8 \mathcal{R}^4$ coupling, one finds


$$\begin{aligned} \mathcal{B}_{(2,0)}(\Omega) &= \frac{8\pi^2 t^2}{45} + \frac{2\pi t}{3} g_1 + \left(3E_2 + g_1^2 - \frac{2}{3}F_2 \right) \\ &+ \frac{1}{\pi t} \left(\frac{3}{2}D_3 - \frac{1}{2}D_3^{(1)} - g_3 - g_1 F_2 + \frac{1}{8\pi} \Delta_\nu (F_2^2 + 2F_4) \right) \\ &+ \frac{1}{16\pi^2 t^2} \left[(\Delta_\tau + 8) (F_2^2 + 4F_4) + \mathcal{K}^c \right] + \mathcal{O}(e^{-2\pi t}) \end{aligned}$$

where

$$g_{k+1}(z) = \int_{\Sigma_1} \kappa_1(w) g_1(z-w) g_k(w)$$

$$F_k(z) = \frac{1}{k!} \int_{\Sigma_1} \kappa_1(w) [g_1(w-z) - g_1(w)]^k$$

$$D_k^{(1)}(z) = \int_{\Sigma_1} \kappa_1(w) g_1(z-w) g_1^{k-1}(w)$$

- By iterating non-separating degenerations, one may extract the asymptotics in the maximal degeneration (aka tropical limit) where the string worldsheet is reduced to a graph.
- For $h = 2$, in the limit $V \equiv [\det \text{Im}\tau]^{-1/2} \rightarrow 0$ one obtains 

$$\mathcal{B}_{(0,1)}^{(2)} = \frac{10\pi}{3V} A_{10} + \frac{5\zeta(3)}{\pi^2} V^2 + \mathcal{O}(e^{-V^{-1/2}})$$

$$\begin{aligned} \mathcal{B}_{(2,0)}^{(2)} &= \frac{\pi^2}{V^2} \left[-\frac{26}{189} A_{00} + \frac{20}{189} A_{02} + \frac{20}{99} A_{11} + \frac{64}{45} A_{20} \right] \\ &+ \frac{V\zeta(3)}{\pi} \left[\frac{9}{5} A_{01} + \frac{8}{3} A_{10} \right] + \frac{3\zeta(5)}{2\pi^3} V^3 A_{01} + \beta \frac{\zeta(3)^2}{\pi^4} V^4 + \mathcal{O}(e^{-V^{-1/2}}) \end{aligned}$$

where A_{ij} are modular functions of $\hat{\tau} = \frac{v_2 + i\sqrt{\det \text{Im}\tau}}{\tau_2}$.

- More precisely, in the fundamental domain


$\mathcal{F} = \{0 < \text{Re}\hat{\tau} < 1, |\tau - \frac{1}{2}| > \frac{1}{2}\}$ for the action of $GL(2, \mathbb{Z})$ on $\hat{\tau}$,

$$A_{ij} := D_{-2n}^{(n)} \left[\left(\hat{\tau}^2 (\hat{\tau} - 1)^2 \right)^i \left(\hat{\tau}^2 - \hat{\tau} + 1 \right)^j \right], \quad n = 3i + j$$

where $D_{-2n}^{(n)}$ is the n -th iterated Maass raising operator.

- The leading terms coincides with the two-loop supergravity integrand in Schwinger time representation. How about subleading terms ?

Tropical limit III



Infrared and ultraviolet divergences I

- Unlike supergravity, perturbative string theory has no ultraviolet divergences, since all boundaries of $\mathcal{M}_{h,N}$ correspond to nodal singularities, which are equivalent to long tubes.
- Infrared divergences originate from massless particle exchange and are captured by supergravity, supplemented by an infinite set of higher derivative interactions induced by massive string modes.
- One way to construct the low energy effective action is to separate the moduli space $\mathcal{M}_{h,n}$ into a disjoint union $\mathcal{M}_{h,N}(L) \cup \mathcal{N}_{h,N}(L)$, where $\mathcal{M}_{h,N}(L)$ retains curves with handles of length $t < L$
- *More precisely, $\mathcal{M}_{h,N}(L)$ is such that the length of the shortest geodesic measured w.r.t. the metric of constant curvature is bigger than $\epsilon = \pi/t$.*

Infrared and ultraviolet divergences II

- The integral over $\mathcal{M}_{h,N}(L)$ is manifestly convergent and analytic as $s_{ij} \rightarrow 0$. Its Taylor expansion provides L -dependent local interactions in the **Wilsonian effective action** for the massless modes.
- The integral over $\mathcal{N}_{h,N}(L)$ generates the supergravity amplitude with UV cut-off $\Lambda = 1/\sqrt{\alpha' L}$. Amplitudes are manifestly independent of the sliding scale L .
- It follows that the L/Λ -dependence of the integral over $\mathcal{M}_{h,N}(L)$ must match the UV divergence of the supergravity amplitude. In particular, a term $a \log L$ in the string integral must match a pole a/ϵ in supergravity in dimension $D - 2\epsilon$.

Green Russo Vanhove'10; BP'18


Infrared and ultraviolet divergences III

- For genus-one amplitudes, a power-like term $a_n \tau_2^n \in \mathcal{B}(\tau)$ leads to

$$\mathcal{I}_1 = \int_{\mathcal{F}_1(L)} \mathcal{B}(\tau) Z_{T^d} \mu_1 \sim a_n \frac{L^{d/2+n-1}}{d/2+n-1}$$

or $\mathcal{I}_1 \sim a_n \log L$ in dimension $D = 8 + 2n$.

- The leading integrand $\mathcal{B}_{(0,0)} \sim 1$ reproduces the one-loop \mathcal{R}^4 divergence in $D = 8$ SUGRA
- Subleading integrands $\mathcal{B}_{(1,0)} \ni \zeta(3)/\tau_2$, $\mathcal{B}_{(0,1)} \ni \zeta(5)/\tau_2^2$, $\mathcal{B}_{(2,0)} \ni \zeta(3)^2/\tau_2^2$ reproduce form factor divergences in $D = 6$, $D = 4$.



Infrared and ultraviolet divergences IV

- For genus-two amplitudes, the integral $\mathcal{I}_2 = \int_{\mathcal{F}_2(L)} \mathcal{B}(\tau) Z_{T^d} \mu_2$ has several sources of infrared divergences:
- 1) the minimal **non separating** degeneration $t \rightarrow \infty$: a term $t^n C_n(\rho, \nu)$ in $\mathcal{B}(\tau)$ leads to the one-loop subdivergence

$$\mathcal{I}_2 \sim \frac{L^{d/2+n-2}}{d/2+n-2} \int_{\mathcal{F}_1} \mu_1 \left[\int_{\Sigma_1} \kappa(\nu) C_n(\rho, \nu) \right] Z_{T^d}$$

or a log divergence in $D = 6 + 2n$ if $\int_{\Sigma_1} \kappa(\nu) C_n(\rho, \nu) \neq 0$;

- In $D = 6$ and $D = 4$, this combines with the genus one divergence of the $\nabla^4 \mathcal{R}^4$ and $\nabla^6 \mathcal{R}^4$ form factors as expected
- 2) the minimal **separating** degeneration $\nu \rightarrow 0$ does not lead to any divergence, since the integrand grows like a power of $\log |\nu|$


Infrared and ultraviolet divergences V

- 3) the **maximal separating** degeneration $V \rightarrow 0$ where $V \equiv [\det \text{Im} \tau]^{-1/2}$: a term $V^{-n} \hat{C}_n(\hat{\tau}) \in \mathcal{B}(\tau)$ leads to

$$\mathcal{I}_2 \sim \frac{L^{d+n-3}}{d+n-3} \int_{\mathcal{H}_1/GL(2,\mathbb{Z})} \hat{\mu}_1 \hat{C}_n(\hat{\tau}) \hat{\tau}_2^{3-n-d}$$

hence a log divergence in $D = 7 + n$.

- The leading integrand $\mathcal{B}_{(1,0)} \sim 1$ reproduces the two-loop $\nabla^4 \mathcal{R}^4$ divergence in $D = 7$ SUGRA [Bern et al '98], while the subleading terms $\mathcal{B}_{(0,1)} \sim \zeta(3) V^2$, $\mathcal{B}_{(2,0)} \sim \zeta(3) V + \zeta(5) V^3 + \zeta(3)^2 V^4$ reproduce form factor divergences in $D = 5$, $D = 6, 4, 3$.



- Genus-two string integrands provide a source of real-analytic Siegel modular forms with remarkably simple asymptotics. Are they secretly holomorphic, or theta lifts of holomorphic objects ? Do they satisfy interesting algebraic or differential identities ?
- Can one use information about these integrands to guess non-perturbative expressions for unprotected effective interactions, e.g. $\nabla^8 \mathcal{R}^4$ in type II strings ? Information about the genus three integrand would be very useful.
- More generally, it is desirable to improve our understanding of string perturbation theory at the practical level, either in the RNS or PS framework, and break the $h = 2, N = 4$ barrier.

- Thanks for your attention !