

Non-forward Balitsky-Kovchegov equation and Vector Mesons

Robi Peschanski¹, Cyrille Marquet² and Gregory Soyez³

1- Service de Physique Théorique - CEA/Saclay, 91191 Gif-sur-Yvette Cedex, FRANCE

2- RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973, USA

3- Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA

Considering the Balitsky-Kovchegov QCD evolution equation in full momentum space, we derive the travelling wave solutions expressing the nonlinear saturation constraints on the dipole scattering amplitude at non-zero momentum transfer. A phenomenological application to elastic vector meson production shows the compatibility of data with the QCD prediction: an *enhanced* saturation scale at intermediate momentum transfer.

1 Motivation

The saturation of parton densities at high energy has been mainly studied for the forward dipole-target scattering amplitude $\mathcal{T}(r, q = 0, Y)$, where r, q, Y are, respectively, the dipole size, the momentum transfer and the total rapidity of the process. For instance, the corresponding QCD Balitsky-Kovchegov (BK) equation [2] has been shown to provide a theoretical insight on the “geometric scaling” properties [3] of the related γ^* -proton cross-sections. Indeed, it can be related to the existence of a scaling for $\mathcal{T}(r, q = 0, Y) \sim \mathcal{T}(r^2 Q^2(Y))$ where the saturation scale is $Q^2(Y) \sim \exp cY$ and the constant c can be interpreted as the critical speed of “travelling wave” solutions of the nonlinear BK equation [4]. Our theoretical and phenomenological subjects are the extension of these properties to the non-forward amplitude $\mathcal{T}(r, q \neq 0, Y)$, which is phenomenologically relevant for the elastic production of vector mesons in deep inelastic scattering.

2 BK equation in full momentum space

In order to study the properties of $\mathcal{T}(r, q \neq 0, Y)$, one has first to deal with both conceptual and technical difficulties. It is known that the BK formalism has been originally derived in impact parameter b but then its validity especially at large b is questionable, since it leads to non physical power-law tails. Hence we start with the formulation of the BK equation in momentum q , which is more *local* but has a non-trivial nonlinear form [5]. In fact, despite this problem, the general method of travelling wave solutions can be extended in the non-forward domain [6]. It consists in 3 steps: first, one solves the equation restricted to its linear




Figure 1: q^2 -dependent saturation scale

part which is related to the non-forward Balitsky Fadin Kuraev Lipatov (BFKL) equation [7] for the dipole-dipole amplitude *via* factorisation and whose solution takes the form of a linear superposition of waves. Second, one finds that the nonlinearities act by selecting the travelling wave with *critical* speed c , in a way which, interestingly, is independent of the specific structure of the nonlinear damping terms. Third, one obtains after enough rapidity evolution, a solution which appears independent from initial conditions ($\mathcal{I}_0 \sim r^{2\gamma_0}$), provided these are sharper than the critical travelling wave front profile $\mathcal{T} \sim r^{2\gamma_c}$, with $\gamma_0 > \gamma_c$. Interestingly enough, QCD color transparency satisfies this criterium. Applying these gen-




Figure 2: ρ (H1) and ϕ (ZEUS) differential cross-sections at $W = 75$ GeV

eral results on the non-forward case one finds the following QCD predictions, depending on the relative magnitude of three scales involved in the process, namely q , k_T^{-1} (the target size) and $k_P^{-1} \equiv r$ (the projectile *i.e.* dipole size).

- Near-Forward region $q \ll k_T \ll k_P$: $Q_s^2(Y) \sim k_T^2 \exp cY$
- Intermediate transfer region $k_T \ll q \ll k_P$: $Q_s^2(Y) \sim q^2 \exp cY$
- High transfer region $q \ll k_T \ll k_P$: No saturation.

Our main prediction is thus the validity of the forward travelling wave solution extended in the non-forward intermediate-transfer domain but with an *enhanced* saturation scale by the ratio q^2/k_T^2 , where k_T is a typically small, nonperturbative scale. Hence we are led to predict *geometric scaling* properties with a purely perturbative initial saturation scale given by the transverse momentum. This saturation scale *enhancement* prediction is confirmed by numerical simulations of the BK solutions as shown in Fig.1.

3 QCD Saturation Model for Exclusive VM production

The differential cross-section for exclusive vector meson (VM) production at HERA, see Fig.2, can be theoretically obtained from the non-forward dipole-proton amplitude and from $\Phi_{T,L}^{\gamma^*V}$, the overlap functions between the (longitudinal and transverse) virtual photon and

vector meson wave-functions [8]. For completion, we used two different VM wave-functions of the literature, without noticeable difference in our conclusions. One writes

$$\frac{d\sigma_{T,L}^{\gamma^* p \rightarrow Vp}}{dq^2} = \frac{1}{16\pi} \left| \int d^2r \int_0^1 dz \Phi_{T,L}^{\gamma^* V}(z, r; Q^2, M_V^2) e^{-izq \cdot r} \mathcal{T}(r, q, Y) \right|^2,$$

Following theoretical prescriptions, we consider a forward dipole-proton amplitude \mathcal{N}_{IIM} satisfactorily describing the total DIS cross-sections in a saturation model [9]. We just make the saturation scale varying with q^2 , following the trend shown in Fig.1 and starting from the forward model one $Q_s^2(Y)$, one writes

$$T(r, q; Y) = 2\pi R_p^2 e^{-Bq^2} \mathcal{N}_{IIM}(r^2 Q_s^2(Y, q)) ; Q_s^2(q, Y) = Q_s^2(Y) (1 + c q^2).$$

Cross-sections	q^2 -Sat.	fixed-Sat.
ρ, σ_{el}	1.156	1.732
$\rho, \frac{d\sigma}{dt}$	1.382	1.489
ϕ, σ_{el}	1.322	2.247
$\phi, \frac{d\sigma}{dt}$	1.076	0.931
Total	1.212	1.480

Table 1: Comparison of the χ^2 /points

Table compares the saturation fits for fixed and q^2 -dependent scales, with a favour for the enhanced-scale model in the total. The model gives a comparable fit with a more conventional non-saturation model using a Q^2 -dependent slope $B \propto M_V^2 + Q^2$. Some of our results for the cross-sections are displayed in the figures. In Fig.2, one shows the results of the fit for ρ -production (H1) and ϕ -production (ZEUS) differential cross-sections for a total $\gamma^* - p$ energy $W = 75\text{GeV}$ and different Q^2 values. Let us finally present our predictions for the

The factor $2\pi R_p^2 e^{-Bq^2}$ comes from the non-perturbative proton form factor. For clarity of the analysis, we considered only B and c as free parameters of the non-forward parametrisations, the others being independently fixed by the forward analysis.

In Table 1, one displays the χ^2 /point obtained by a fit of ρ (47 data points) and ϕ (34 points) total elastic production cross-sections and of ρ (50 data points) and ϕ (70 points) differential cross-sections. The Ta-




Figure 3: Predictions for the DVCS measurements. Left plot: cross-section, right plot: differential cross-section.

DVCS cross-section, which is obtained without any free parameter from our analysis. In Fig.3, they are compared with the available data and the agreement is good in the simple chosen parametrisation.

4 Conclusions

Let us summarize our new results

- *Saturation at non-zero transfer*: The Balitsky-Kovchegov QCD evolution equation involving full momentum transfer predicts (besides the known $q = 0$ case) saturation in the *intermediate* transfer range, namely for $Q_0 < q < Q$, where Q_0 (resp. Q) is the target (resp. projectile) typical scale.

- *Characterisation of the universality class*: The universality class of the corresponding travelling-wave solutions is governed by a purely perturbative saturation scale $Q_s(Y) \equiv q^2 \Omega(Y)$, where $\Omega(Y) \sim e^{cY}$ is the same rapidity evolution factor as in the forward case. Consequently the *intermediate transfer* saturation scale gets enhanced by a factor q^2/Q_0^2 .

- *Phenomenology of Vector mesons*: The QCD predictions are applied in the experimentally accessible *intermediate transfer* range of vector meson production. The model uses an interpolation between the forward and non-forward saturation scale together with a parameter-frozen forward saturation model. It fits better the data on ρ (H1) and ϕ (ZEUS) cross-sections than for a non-enhanced saturation.

- *Prospects*: The next phenomenological prospect is to add charm to the discussion, both with the modification of the forward case by including the charm contribution [10] and by also considering the production of Ψ mesons. On a theoretical ground, it would be interesting to go beyond the mean-field approximation of the BK equation.

References

- [1] Slides:
<http://indico.cern.ch/contributionDisplay.py?contribId=75&sessionId=7&confId=9499>
- [2] I. Balitsky, *Nucl. Phys.* **B463** (1996) 99; *Phys. Lett.* **B518** (2001) 235;
Yu.V. Kovchegov, *Phys. Rev.* **D60** (1999) 034008; *Phys. Rev.* **D61** (2000) 074018.
- [3] A.M. Stasto, K. Golec-Biernat and J. Kwiecinski, *Phys. Rev. Lett.* **86** (2001) 596;
C. Marquet and L. Schoeffel, *Phys. Lett.* **B639** (2006) 471.
- [4] S. Munier and R. Peschanski, *Phys. Rev. Lett.* **91** (2003) 232001; *Phys. Rev.* **D69** (2004) 034008; **D70** (2004) 077503.
- [5] C. Marquet and G. Soyez, *Nucl. Phys.* **A760** (2005) 208.
- [6] C. Marquet, R. Peschanski and G. Soyez, *Nucl. Phys.* **A756** (2005) 399.
- [7] L. N. Lipatov, *Sov. Phys. JETP* **63**, 904 (1986) [*Zh. Eksp. Teor. Fiz.* **90**, 1536 (1986)].
- [8] C. Marquet, R. Peschanski and G. Soyez, *Exclusive vector meson production at HERA from QCD with saturation*, hep-ph/0702171.
- [9] E. Iancu, K. Itakura and S. Munier, *Phys. Lett.* **B590** (2004) 199.
- [10] K. Golec-Biernat and S. Sapeta, *Phys. Rev.* **D74** (2006) 054032;
H. Kowalski, L. Motyka and G. Watt, *Phys. Rev.* **D74** (2006) 074016;
G. Soyez, *Saturation QCD predictions with heavy quarks at HERA*, arXiv.0705.3672.