Proceedings of Cheju Island APCTP School and Conference © World Scientific Publishing Company

CONFORMAL BOUNDARY CONDITIONS

JEAN-BERNARD ZUBER

Service de Physique Théorique CEA-Saclay F-91191 Gif-sur-Yvette, France E-mail: zuber@spht.saclay.cea.fr

The question of boundary conditions in conformal field theories is discussed, in the light of recent progress. Solving the consistency condition known as Cardy equation is shown to amount to the algebraic problem of finding integer valued representations of the fusion algebra. Graphs encode these boundary conditions in a natural way, but are also relevant in several aspects of physics "in the bulk". Quantum algebras attached to these graphs contain information on structure constants of the operator algebra, on the Boltzmann weights of the corresponding integrable lattice models etc. Thus Boundary Conformal Field Theory offers a new perspective on several old problems.

0. Introduction

The study of boundary conditions in conformal field theories (CFT) and in the related integrable models has been experiencing a renewal of interest over the last three or four years. This has been caused by its relevance in string and brane theory on the one hand, and in various problems of condensed matter on the other: see [1] for an introduction and references to the first subject, and [2] for the second. As a result, there has been a blossoming of papers studying the possible boundary conditions, the boundary fields and their couplings in the framework of CFT (see [3] for a fairly extensive bibliography as of mid 99); a systematic discussion of boundary conditions preserving the integrability, both in lattice models [4] and in (classical or quantum) field theories [5]; and an investigation of what happens to a critical system in the presence of boundary perturbations, its renormalisation group flows etc [6]. At the same time, new and unexpected connections with "pure" mathematics –operator algebras, quantum symmetries– have also been revealed.

The purpose of these lectures is of course *not* to present exhaustively all these interesting developments, but just to offer a pedagogical (and maybe somewhat biaised) introduction to their simplest aspects and to some of the recent progress. After briefly recalling basic facts on CFT, their chiral constituents and how they are assembled into physically sensible theories, I turn to the discussion of boundary conditions. I show how solving the consistency

condition known as Cardy equation amounts to the algebraic problem of finding non negative integer valued matrix representations of the fusion algebra. These matrices are the adjacency matrices of graphs, which thus encode the boundary conditions in a natural way (sect. 2). The study of the operator algebra of boundary fields exposes new algebraic features attached to these graphs (sect. 3). The latter also contain information on Boltzmann weights of associated lattice integrable models, as I mention briefly in the last section.

1. A lightning review of CFT

This section is devoted to a fast summary of concepts and notations in rational conformal field theories (RCFT).

1.1. Chiral data of RCFT

A rational conformal field theory is defined in terms of a certain number of data. The first set of data specifies the properties of each chiral half, i.e. of the holomorphic or of the antiholomorphic sector of the theory. One is given a certain *chiral algebra*, \mathcal{A} : it may be the Virasoro algebra Vir itself, with its generators L_n , $n \in \mathbb{Z}$, or equivalently the energy-momentum tensor $T(z) = \sum_{n \in \mathbb{Z}} z^{-n-2}L_n$. It may also be one of the extensions of Vir: superconformal algebra, current algebra, W-algebra etc. One is also given a *finite* set \mathcal{I} of irreducible representation spaces (modules) $\{\mathcal{V}_i\}_{i \in \mathcal{I}}$ of \mathcal{A} . Each of these representations of \mathcal{A} is also a representation (reducible or irreducible) of Vir, with a central charge c and with a conformal weight (the lowest eigenvalue of L_0) denoted h_i . Let's recall for future use that c also specifies the coefficient of the anomalous term in the transformation of the energy-momentum tensor T(z) under an analytic change of coordinate $z \mapsto \zeta(z)$

$$\tilde{T}(\zeta) = \left(\frac{\partial z}{\partial \zeta}\right)^2 T(z) + \frac{c}{12} \{z, \zeta\}$$
(1.1)

where $\{z, \zeta\}$ denotes the schwarzian derivative

$$\{z,\zeta\} = \frac{\frac{\partial^3 z}{\partial \zeta^3}}{\frac{\partial z}{\partial \zeta}} - \frac{3}{2} \left(\frac{\frac{\partial^2 z}{\partial \zeta^2}}{\frac{\partial z}{\partial \zeta}}\right)^2 \,. \tag{1.2}$$

By convention, the label i = 1 denotes the identity representation (for which $h_1 = 0$). Finally, we denote by \mathcal{V}_{i^*} the complex conjugate representation of \mathcal{V}_i ; the identity representation is self-conjugate, $1^* = 1$.

Each of these representations is graded for the action of the Virasoro generator L_0 : all the eigenvalues of L_0 differ from the lowest one, h_i , by a non negative integer ¹, and the eigenspace of eigenvalue $h_i + p$ has a certain dimension $\#_p^{(i)}$. It is natural to introduce the "character" of the representation \mathcal{V}_i , which is, up to a prefactor, the generating function of these dimensions

$$\chi_i(q) = \operatorname{tr} q^{L_0 - \frac{c}{24}} = q^{h_i - \frac{c}{24}} \sum_{p=0}^{\infty} \#_p^{(i)} q^p .$$
(1.3)

The simplest example is given by the integrable representations of the affine (current) algebra $\widehat{sl}(2)$. For an integer value of the central charge of the affine algebra (or level) k, the Virasoro central charge is c = 3k/(k+2), and there is a finite set of integrable representations, labelled by an integer $j: 1 \leq j \leq k+1$, whose conformal weights are $h_j = (j^2 - 1)/4(k+2)$. In that case, the conjugation is trivial: $\mathcal{V}_j = \mathcal{V}_{j^*}$. For the representation (j, k), the character reads

$$\chi_j(q) = \frac{1}{\eta^3(q)} \sum_{p=-\infty}^{\infty} (2(k+2)p+j) q^{\frac{(2(k+2)p+j)^2}{4(k+2)}}, \qquad (1.4)$$

where $\eta(q) = q^{\frac{1}{24}} \prod_{1}^{\infty} (1-q^n)$ is the Dedekind function. Such characters are called "specialized characters" since they count states according to their L_0 grading only. Non-specialized characters can be introduced, which are sensitive to the Cartan algebra generator J_0

$$\chi_j(q,u) = \operatorname{tr} q^{L_0 - \frac{c}{24}} e^{2\pi i u J_0} .$$
(1.5)

The expressions of non-specialized characters and/or for higher rank algebras may be found in [7].

Another example is provided by the minimal c < 1 theories. They are parametrized by a pair of coprime integers p and p', and the central charge takes the values $c = 1-6(p-p')^2/pp'$. The irreducible representations of Vir are labelled by a pair of integers (r, s), $1 \le r \le p'-1$, $1 \le s \le p-1$, modulo the identification $(r, s) \equiv (p'-r, p-s)$. Their conformal weights read

$$h_{(r,s)} = h_{(p'-r,p-s)} = \frac{(rp - sp')^2 - (p - p')^2}{4pp'}.$$
(1.6)

Again the conjugation acts trivially. The character of this irreducible representation reads, with the notations $\lambda := (rp - sp')$, $\lambda' := (rp + sp')$

$$\chi_{(r,s)}(q) = \frac{1}{\eta(q)} \sum_{n \in \mathbb{Z}} \left(q^{\frac{(2npp' + \lambda)^2}{4pp'}} - q^{\frac{(2npp' + \lambda')^2}{4pp'}} \right) .$$
(1.7)

¹ This is strictly true only for algebras, whose generators are integrally graded. In a superconformal algebra, or in a parafermionic algebra, the grading would be fractional.

Modular transformations

In the previous expressions, q is a dummy variable. If, however, q is regarded as a complex variable of modulus less than one and written as $q = \exp 2i\pi\tau$, with τ a complex number of positive imaginary part, one may prove that the sum converges and has remarkable modular properties. Under a $PSL(2,\mathbb{Z})$ transformation of τ :

$$\tau \mapsto \frac{a+b\tau}{c+d\tau}$$
, $a, b, c, d \in \mathbb{Z}$, $ad-bc=1$,

the set of functions χ_i transforms linearly, and in fact supports a unitary representation of (the double cover of) $PSL(2,\mathbb{Z})$. In particular there exists a unitary matrix S implementing the transformation $\tau \mapsto -1/\tau$. If $\tilde{q} := \exp{-\frac{2i\pi}{\tau}}$, there exists a unitary $|\mathcal{I}| \times |\mathcal{I}|$ matrix S such that

$$\chi_i(q) = \sum_{j \in \mathcal{I}} S_{ij} \chi_j(\tilde{q}) .$$
(1.8)

Moreover the matrix S satisfies $S^T = S$, $S^{\dagger} = S^{-1}$, $(S_{ij})^* = S_{i^*j} = S_{ij^*}$, $S^2 = C$ = the conjugation matrix defined by $C_{ij} = \delta_{ij^*}$, $S^4 = I$.²

For the c<1 minimal representations,

$$\mathcal{I} = \{ (r,s) \equiv (p'-r, p-s) ; \quad 1 \le r \le p'-1, \ 1 \le s \le p-1 \}$$

and the ${\cal S}$ matrix reads

$$S_{(r,s),(r's')} = \sqrt{\frac{8}{pp'}} (-1)^{(r+s)(r'+s')} \sin \pi r r' \frac{p-p'}{p'} \sin \pi s s' \frac{p-p'}{p}$$
(1.9)

For the $\widehat{sl}(2)$ affine algebra, at level k, for which $\mathcal{I} = \{1, 2, \dots, k+1\}$, one finds

$$S_{jj'} = \sqrt{\frac{2}{k+2}} \sin \frac{\pi j j'}{k+2} , \quad j, j' \in \mathcal{I} .$$
 (1.10)

For non-specialised characters, the transformation reads:

$$\chi_i(q,u) = e^{ik\pi u^2/\tau} \sum_{j\in\mathcal{I}} S_{ij}\chi_j(\tilde{q},u/\tau) .$$
(1.11)

² The fact that $S^2 = C$ rather than $S^2 = I$ as expected from the transformation $\tau \mapsto -1/\tau$ signals that we are dealing with a representation of a double covering of the modular group.

The expression for more general affine algebras may be found in [7].

One notes that the S matrix of the minimal case (1.9) is "almost" the tensor product of two matrices of the form (1.10), at two different levels k = p - 2 and k' = p' - 2. This would be true for |p - p'| = 1 and if one could omit the identification $(r, s) \equiv (p' - r, p - s)$. This is of course not a coincidence but reflects the "coset construction" of c < 1 representations of Vir out of the affine algebra $\widehat{sl}(2)$ [8]. We shall encounter below again this fact that minimal cases are "almost" the tensor products of the $\widehat{sl}(2)$ ones.

Fusion Algebra

Another concept of crucial importance for our discussion is that of fusion algebra. Fusion is an associative and commutative operation among representations of chiral algebras of RCFTs, inherited from the operator product algebra of Quantum Field Theory. It looks similar to the usual tensor product of representations, but contrary to the latter, it is consistent with the finiteness of the set \mathcal{I} and it preserves the central elements (instead of adding them). I shall refer to the literature [7,9] for a systematic discussion of this concept, and just introduce a notation \star to denote it and distinguish it from the tensor product. It is natural to decompose the fusion of two representations of a chiral algebra on the irreducible representations, thus defining multiplicities, or "fusion coefficients"

$$\mathcal{V}_i \star \mathcal{V}_j = \bigoplus_k \mathcal{N}_{ij}{}^k \mathcal{V}_k, \qquad \mathcal{N}_{ij}{}^k \in \mathbb{N} .$$
(1.12)

There is a remarkable formula, due to Verlinde [10], expressing these multiplicities in terms of the unitary matrix S:

$$\mathcal{N}_{ij}{}^{k} = \sum_{\ell \in \mathcal{I}} \frac{S_{i\ell} \, S_{j\ell} \, (S_{k\ell})^{*}}{S_{1\ell}} \,. \tag{1.13}$$

Chiral Vertex Operators

The field theoretic description of chiral halves of CFT (or "chiral cft") makes use of "chiral vertex operators" (CVO): $\phi_{ij}^k(z)$ is a z-dependent intertwiner

from $\mathcal{V}_i \star \mathcal{V}_j$ to \mathcal{V}_k and may be diagrammatically depicted as $\underbrace{k \leftarrow i}_{z,t} j$. In fact, there are as many as $\mathcal{N}_{ij}{}^k$ such independent intertwiners and the notation $\phi_{ij,t}^k(z)$ and the diagram have to be supplemented by a multiplicity label $t \in \{1, 2, \dots, \mathcal{N}_{ij}{}^k\}$. CVO may be composed ("fused") as $\phi_{il}^m(z_1)\phi_{jk}^l(z_2)$ and there is a matrix $F_{lp} \begin{bmatrix} i & j \\ m & k \end{bmatrix}_{t_1 t_2}^{ut}$ expressing that there are two distinct but equivalent

ways of intertwining
$$\mathcal{V}_i \star \mathcal{V}_j \star \mathcal{V}_k$$
 and $\mathcal{V}_m : \frac{m \left| \begin{array}{c} l \\ t_1 z_1 \\ t_2 z_2 \end{array}^j}{k} \right| \xrightarrow{F_{lp}} \frac{m \left| \begin{array}{c} l \\ t_2 u \\ z_2 u \end{array}^j}{k}$. The

matrix F satisfies orthonormality and completeness conditions, and moreover, a quintic ("pentagon") identity expressing the consistency (the associativity) in the fusion of three CVOs. There is also the operation of braiding and the matrix $B_{lp}\begin{bmatrix}i & j\\m & k\end{bmatrix}(\epsilon)$ which relates $\phi_{il}^m(z_1)\phi_{jk}^l(z_2)$ and $\phi_{jp}^m(z_2)\phi_{ik}^p(z_1)$, $\epsilon =$ sign ($\Im m z_{12}$), see [11] for a systematic discussion.

Notice finally that the matrix S admits an extension, denoted S(j), such that S(1) = S. The matrix S(p) gives the modular transformation of 1-point conformal block $\langle \phi_{pi}^i \rangle_{\tau}$ on the torus. See [11] for the explicit expression of S(p) in the simplest cases.

The data c (or k, etc), $\{\mathcal{V}_i, h_i\}_{i \in \mathcal{I}}$, $S(p)_{ij}$, $\mathcal{N}_{ij}{}^k$, $F\begin{bmatrix}i & j\\k & l\end{bmatrix}$ and $B\begin{bmatrix}i & j\\k & l\end{bmatrix}(\pm)$ form what may be called the "chiral data" of the RCFT. They are the basic ingredients in the construction of physical theories. The consistency of the latter imposes however that adequate conditions be satisfied by their spectral data and by the structure constants of their Operator Product Algebra (OPA). This is what we shall consider now, examining in turn the cases of the theory "in the bulk", i.e. in the absence of any boundary, and then in the half-plane.

1.2. Spectral and OPA Data in the Bulk

In the plane punctured at the origin, equiped with the coordinate z, or equivalently on the cylinder of perimeter L with the coordinate w, with the conformal mapping from the latter to the former $z = \exp -2\pi i w/L$, a given RCFT is described by a Hilbert space \mathcal{H}_P . This Hilbert space is decomposable into a *finite* sum of irreducible representations of **two** copies of the chiral algebra (Vir or else), associated with the holomorphic and anti-holomorphic sectors of the theory:

$$\mathcal{H}_P = \oplus Z_{j\bar{j}} \mathcal{V}_j \otimes \mathcal{V}_{\bar{j}} , \qquad (1.14)$$

with (non negative integer) multiplicities $Z_{j\bar{j}}$. By the state-field correspondence, (1.14) also describes the spectrum of primary fields of the theory, i.e. of those fields that transform as heighest weight representations of $\mathcal{A} \otimes \mathcal{A}$. On the cylinder, it is natural to think of the Hamiltonian as the operator of translation along its axis (the imaginary axis in w), or along any helix, defined by its period τL in the w plane, with $\Im m \tau > 0$. If L_{-1}^{cyl} and $\bar{L}_{-1}^{\text{cyl}}$ are the two generators of translation in w and \bar{w} , $H^{\text{cyl},\tau} = (\tau L_{-1}^{\text{cyl}} + \bar{\tau} \bar{L}_{-1}^{\text{cyl}})$. Mapped back in the plane using the transformation law of the energy-momentum tensor (1.1), L_{-1}^{cyl} reads

$$L_{-1}^{\text{cyl}} = -\frac{2\pi i}{L} (L_0 - \frac{c}{24})$$
(1.15)

where the term c/24 comes from the schwarzian derivative of the exponential mapping. The evolution operator of the system, i.e. the exponential of L times the Hamiltonian is thus

$$e^{-H^{\text{cyl},\tau}L} = e^{2\pi i \left(\tau \left(L_0 - \frac{c}{24}\right) - \bar{\tau}(\bar{L}_0 - \frac{c}{24})\right)} .$$
(1.16)

A convenient way to encode the information (1.14) is to look at the partition function of the theory on a torus \mathbb{T} . Up to a global dilatation, irrelevant here, a torus may be defined by its modular parameter τ , $\Im m \tau > 0$, such that its two periods are 1 and τ . Equivalently, it may be regarded as the quotient of the complex plane by the lattice generated by the two numbers 1 and τ :

$$\mathbb{T} = \mathbb{C}/(\mathbb{Z} \oplus \tau \mathbb{Z}) , \qquad (1.17)$$

in the sense that points in the complex plane are identified according to $w \sim w' = w + n + m\tau$, $n, m \in \mathbb{Z}$. There is, however, a redundancy in this description of the torus: the modular parameters τ and $M\tau$ describe the same torus, for any modular transformation $M \in PSL(2,\mathbb{Z})$. The partition function of the theory on this torus is just the trace of the evolution operator (1.16), with the trace taking care of the identification of the two ends of the cylinder into a torus

$$Z = \operatorname{tr}_{\mathcal{H}_P} e^{2\pi i \left[\tau (L_0 - \frac{c}{24}) - \bar{\tau}(\bar{L}_0 - \frac{c}{24})\right]} .$$
(1.18)

Using (1.14) and the definition (1.3) of characters, this trace may be written as

$$Z = \sum Z_{j\bar{j}} \chi_j(q) \chi_{\bar{j}}(\bar{q}) \qquad q = e^{2\pi i \tau} \quad \bar{q} = e^{-2\pi i \bar{\tau}}.$$
 (1.19)

Let's stress that in these expressions, $\bar{\tau}$ is the complex conjugate of τ , and \bar{q} that of q, and therefore, $Z = \sum Z_{j\bar{j}}\chi_j(q) (\chi_{\bar{j}}(q))^*$ is a *sesquilinear* form in the characters. It is a natural physical requirement that this partition function be intrinsically attached to the torus, and thus be invariant under modular transformations. Finally one imposes the extra condition $Z_{11} = 1$ which expresses the unicity of the identity representation (i.e. of the "vacuum").

One is thus led to the problem of finding all possible sesquilinear forms (1.19) with non negative integer coefficients that are modular invariant, and such that $Z_{11} = 1$. As explained in the previous section, the finite set of characters of any RCFT, labelled by \mathcal{I} , supports a unitary representation of the modular group. This implies that any diagonal combination of characters $Z = \sum_{i \in \mathcal{I}} \chi_i(q) \chi_i(\bar{q})$ is modular invariant. In that case, all representations of $\mathcal{A} \otimes \mathcal{A}$ appearing in (1.14) are left-right symmetric, and thus all primary fields

are spinless: $h_j - h_{\bar{j}} = 0$. This situation is referred to as the "diagonal case" or "diagonal theory". Other solutions are, however, known to exist.

The problem has been completely solved only in a few cases: for the RCFTs with an affine algebra, the $\hat{sl}(2)$ [12] and $\hat{sl}(3)$ [13] theories at arbitrary level, plus a host of cases with constraints on the level, e.g. the general $\hat{sl}(N)$ for k = 1 [14]; some of the associated coset theories [8] have also been fully classified, including all the minimal c < 1 theories, N = 2 "minimal" superconformal theories, etc. A good review on the current state of the art is provided by T. Gannon [15]. For a short account of the cases of $\hat{sl}(2)$ and $\hat{sl}(3)$, see [16].

In the case of CFTs with a current algebra, it is in fact better to look at the same problem of modular invariants after replacing in (1.19) all specialized characters by non-specialized ones, v.i.z. $\sum Z_{j\bar{j}}\chi_j(q,\mathbf{u}) (\chi_{\bar{j}}(q,\mathbf{u}))^*$. Because these non-specialized characters are linearly independent, there is no ambiguity in the determination of the multiplicities $Z_{j\bar{j}}$ from Z. This alternative form of the partition function may be seen to result from a modification of the energy-momentum tensor $T(z) \to T(z) - \frac{2\pi i}{L}(\mathbf{u}, \mathbf{J}(z)) - \frac{k}{2}\left(\frac{2\pi}{L}\right)^2(\mathbf{u}, \mathbf{u})$, see [3].

The matrix $Z_{j\bar{j}}$ of (1.14) gives us the spectrum of primary fields of the theory. We have also to determine the couplings of these fields. This may be done by expressing them in terms of the CVO as

$$\Phi_{(i,\bar{i})}(z,\bar{z}) = \sum_{j,\bar{j},k,\bar{k},t,\bar{t}} d^{(k,\bar{k});t,\bar{t}}_{(i,\bar{i})(j,\bar{j})} \phi^k_{i,j;t}(z) \otimes \phi^{\bar{k}}_{\bar{i},\bar{j};\bar{t}}(\bar{z}) .$$
(1.20)

Then matrix elements of $\Phi_{(i,\bar{i})}$ between highest weight states are given in terms of those of CVO, which are supposed to be known

$$\langle k, \bar{k} | \Phi_{(i,\bar{i})}(1,1) | j, \bar{j} \rangle = \sum_{t,\bar{t}} d^{(k,\bar{k});t,\bar{t},\bar{t}}_{(i,\bar{i})(j,\bar{j})} \langle k | \phi^k_{i,j;t}(1) | j \rangle \langle \bar{k} | \phi^{\bar{k}}_{\bar{i},\bar{j};\bar{t}}(1) | \bar{j} \rangle .$$
(1.21)

For $\hat{sl}(2)$ or minimal theories, the latter have been explicitly computed in [17,18]. In the diagonal theories, the *d* coefficients are equal to a product of Kronecker delta symbols $\delta_{i\bar{i}}\delta_{j\bar{j}}\delta_{k\bar{k}}\delta_{t\bar{t}}$ with $t = 1, \dots, \mathcal{N}_{ij}{}^k$ implying that *d* vanishes if $\mathcal{N}_{ij}{}^k = 0$. Thus the expansion coefficients $d_{(i,\bar{i})(j,\bar{j})}^{(k,\bar{k});t,\bar{t}}$ give in general the *relative* OPE coefficients of the non-diagonal model with respect to the diagonal model of same central charge. These numbers are constrained by the requirement of locality of the physical correlators, which makes use of the braiding matrices $B(\pm)$. The resulting set of coupled quadratic equations has been fully solved only in the sl(2) cases (see [17,19] and further references therein).

A curious empirical fact was then observed: one may introduce graphs, whose properties are intimately connected with some features of the modular invariants or of the structure constants. In the simplest case of $\hat{sl}(2)_k$ theories, these graphs are the well-known ADE Dynkin diagrams of Coxeter number h = k+2, and the set of labels j of the diagonal terms appearing in Z in (1.19) is precisely the set of so-called exponents labelling the eigenvalues $2 \cos \pi j/h$ of the adjacency matrix of the diagram (see Table 1). Moreover let's introduce the corresponding eigenvectors ψ_a^j of the adjacency matrix, (a being a vertex of the diagram G) and define the structure constants of the so-called Pasquier algebra [20]

$$M_{ij}{}^{k} = \sum_{a \in G} \frac{\psi_{a}^{i} \psi_{a}^{j} (\psi_{a}^{k})^{*}}{\psi_{a}^{1}} .$$
(1.22)

Then the structure constants $d_{(i,i),(j,j)}^{(k,k)}$ of spinless fields in the theory may be shown to be equal to those of the Pasquier algebra $d_{(i,i),(j,j)}^{(k,k)} = M_{ij}^{k}$. Thus the ADE graph encodes some non trivial information about a subsector of the theory, namely that of spinless fields, their spectrum and OPE.

Table 1: ADE graphs, their Coxeter number and their exponents		
	h	exponents
$A_n \xrightarrow{1 \ 2 \ 3} ^n ^n$	n + 1	$1, 2, \cdots, n$
$D_{n+2} \xrightarrow{1} 2 3 \xrightarrow{n+1} 6$	2(n+1)	$1, 3, \cdots, 2n + 1, n + 1$
$E_6 \xrightarrow{1}{2} \xrightarrow{4}{3} \xrightarrow{7}{7}$	12	1, 4, 5, 7, 8, 11
$E_7 \qquad \stackrel{1}{\bullet} \stackrel{2}{\bullet} \stackrel{4}{\bullet} \stackrel{5}{\bullet} \stackrel{6}{\bullet}$	18	1, 5, 7, 9, 11, 13, 17
$E_8 \qquad \stackrel{1}{\bullet} \stackrel{2}{\bullet} \stackrel{8}{\bullet} \stackrel{4}{\bullet} \stackrel{5}{\bullet} \stackrel{6}{\bullet} \stackrel{7}{\bullet}$	30	1, 7, 11, 13, 17, 19, 23, 29

These empirical facts are known to extend to more complicated theories. In general, we expect that among the pairs (j, \bar{j}) appearing in Z, a special role

will be played by the diagonal subset

$$\mathcal{E} = \{ j | j = \bar{j}, Z_{jj} \neq 0 \} , \qquad (1.23)$$

the elements of which, the "exponents" of the theory, are counted with the multiplicity Z_{jj} . (We assume that \mathcal{E} is stable under conjugation: j and j^* occur with the same multiplicity.) Then the relevant graphs G_i are labelled by the set \mathcal{I} , they have a common set of vertices and their spectrum is described by the set \mathcal{E} in the sense that their eigenvalues are of the form S_{ij}/S_{1j} , $j \in \mathcal{E}$.

The origin and interpretation of these empirical facts had remained elusive until recently. It is one of the virtues of BCFT to have cast a new light on these facts and to have offered a new framework in which they appear more natural and systematic.

2. Boundary Conformal Field Theory

2.1. Spectral Data in the Upper Half Plane

We now turn to the study of RCFT in a half-plane. There are several physical reasons to look at this problem, as mentionned in the Introduction. Here we shall only look at the new information and perspective that this situation gives us on the general structure of RCFT.

In a half-plane, the admissible diffeomorphisms must respect the boundary, taken as the real axis: thus only real analytic changes of coordinates, satisfying $\epsilon(z) = \bar{\epsilon}(\bar{z})$ for $z = \bar{z}$ real, are allowed. The energy momentum itself has this property:

$$T(z) = T(\bar{z})|_{\text{real axis}} , \qquad (2.1)$$

which expresses simply the absence of momentum flow across the boundary and which enables one to extend the definition of T to the lower half-plane by $T(z) := \overline{T}(z)$ for $\Im m z < 0$. There is thus only one copy of the Virasoro algebra $L_n = \overline{L}_n$. This continuity equation (2.1) on T extends to more general chiral algebras and their generators, at the price however of some complication. In general, the continuity equation on generators of the chiral algebra involves some automorphism of that algebra:

$$W(z) = \Omega \overline{W}(\overline{z})|_{\text{real axis}}$$
(2.2)

(see [3] and further references therein).

The half-plane, punctured at the origin, (which introduces a distinction between the two halves of the real axis), may also be conformally mapped on an infinite horizontal strip of width L by $w = \frac{L}{\pi} \log z$. Boundary conditions, loosely specified at this stage by labels a and b, are assigned to fields on the two boundaries z real > 0, < 0 or $\Im w = 0, L$. For given boundary conditions on the generators of the algebra and on the other fields of the theory, i.e. for given automorphisms Ω and given a, b, we may again use a description of the system by a Hilbert space of states \mathcal{H}_{ba} (we drop the dependence on Ω for simplicity). On the half-plane or on the finite-width strip, only **one copy** of the Virasoro algebra, or of the chiral algebra \mathcal{A} under consideration, acts on \mathcal{H}_{ba} , and this space decomposes on representations of Vir or \mathcal{A} according to

$$\mathcal{H}_{ba} = \oplus n_{ib}{}^a \mathcal{V}_i , \qquad (2.3)$$

with a new set of multiplicities $n_{ib}{}^a \in \mathbb{N}$. The natural Hamiltonian on the strip is the translation operator in $\Re e w$, hence, mapped back in the half-plane

$$H_{b|a} = \frac{2\pi}{L} \left(L_0 - \frac{c}{24} \right) .$$
 (2.4)

To recapitulate, in order to fully specify the operator content of the theory in various configurations, we need not only determine the multiplicities "in the bulk" $Z_{j\bar{j}}$ of (1.14), but also the possible boundary conditions a, b on a half-plane and the associated multiplicities $n_{ib}{}^a$. This will be our task in the following, and as we shall see, a surprising fact is that the latter have some bearing on the former.

2.2. Boundary states

In the same way that we found useful to chop a finite segment of the infinite cylinder and identify its ends to make a torus, it is suggested to consider a finite segment of the strip – or a semi-annular domain in the half-plane– and identify its edges, thus making a cylinder. This cylinder can be mapped back into an annular domain in the plane, with open boundaries. More explicitly, consider the segment $0 \leq \Re e w \leq T$ of the strip –i.e. the semi-annular domain in the upper half-plane comprised between the semi-circles of radii 1 and $e^{\pi T/L}$, the latter being identified. It may be conformally mapped into an annulus in the complex plane by $\zeta = \exp(-2i\pi w/T)$, of radii 1 and $e^{2\pi L/T}$, see Fig. 1. By working out the effect of this change of coordinates on the energy-momentum T, using (1.1), one finds that (2.1) implies

$$\zeta^2 T(\zeta) = \overline{\zeta}^2 \overline{T}(\overline{\zeta}) \quad \text{for } |\zeta| = 1, \ e^{2\pi \frac{L}{T}} .$$
(2.5)

 $12 \quad J.\text{-}B. \ Zuber$



Fig. 1: The same domain seen in different coordinates: a semicircular annulus, with the two half-circles identified, a rectangular domain with two opposite sides identified, and a circular annulus.

After radial quantization, this translates into a condition on *boundary states* $|a\rangle$, $|b\rangle \in \mathcal{H}_P$ which describe the system on these two boundaries.

$$(L_n - \bar{L}_{-n})|a\rangle = 0 \tag{2.6}$$

(and likewise for $|b\rangle$).

Exercise : assuming that W transforms as a primary field of conformal weight $(h_W, 0)$, find the corresponding condition on $W(\zeta)$. Then show that the analogue of (2.6) reads $(W_n - (-1)^{h_W} \Omega(\bar{W}_{-n}))|a\rangle = 0.$

We shall now look for a basis of states, solutions of this linear system of boundary conditions. One may seek solutions of these equations in each $\mathcal{V}_j \otimes \mathcal{V}_{\bar{j}} \subset \mathcal{H}_P$, since these spaces are invariant under the action of the two copies of Vir or of the chiral algebra \mathcal{A} . Consider only for simplicity the case of the Virasoro generators.

Lemma There is an independent "Ishibashi state" $|j\rangle$, solution of (2.6), for each $j = \overline{j}$, i.e. $j \in \mathcal{E}$, the set of exponents.

Proof (G. Watts)[21]: Use the identification between states $|a\rangle \in \mathcal{V}_j \otimes \mathcal{V}_{\bar{j}}$ and operators $X_a \in \text{Hom}(\mathcal{V}_{\bar{j}}, \mathcal{V}_j)$, namely $|a\rangle = \sum_{n,\bar{n}} a_{n,\bar{n}} |j,n\rangle \otimes |\bar{j},\bar{n}\rangle \leftrightarrow X_a =$ $\sum_{n,\bar{n}} a_{n,\bar{n}} |j,n\rangle \langle \bar{j},\bar{n}|$. Here we make use of the scalar product in $\mathcal{V}_{\bar{j}}$ for which $\bar{L}_{-n} = \bar{L}_n^{\dagger}$, hence (2.6) means that $L_n X_a = X_a L_n$, i.e. X_a intertwines the action of Vir in the two irreducible representations \mathcal{V}_j and $\mathcal{V}_{\bar{j}}$. By Schur's lemma, this implies that they are equivalent, $\mathcal{V}_j \sim \mathcal{V}_{\bar{j}}$, i.e. that their labels coincide $j = \bar{j}$ and that X_a is proportional to P_j , the projector in \mathcal{V}_j . We shall denote $|j\rangle$ the corresponding state, solution to (2.6).

Since "exponents" $j \in \mathcal{E}$ may have some multiplicity, an extra label should be appended to our notation $|j\rangle\rangle$. We omit it for the sake of simplicity. The previous considerations extend with only notational complications to more general chiral algebras and their possible gluing automorphisms Ω . See [3] for more details and more references on these points. Also, see [22] for an alternative and mathematically more precise discussion of Ishibashi states.

The normalization of this "Ishibashi state" requires some care. One first notices that, for \tilde{q} a real number between 0 and 1,

$$\langle\!\langle j' | \tilde{q}^{\frac{1}{2}(L_0 + L_0 - \frac{c}{12})} | j \rangle\!\rangle = \delta_{jj'} \chi_j(\tilde{q})$$
 (2.7)

up to a constant that we choose equal to 1. It would seem natural to then define the norm of these states by the limit $\tilde{q} \rightarrow 1$ of (2.7). This limit diverges, however, and the adequate definition is rather

$$\langle\!\langle j \| j' \rangle\!\rangle = \delta_{jj'} S_{1j} \tag{2.8}$$

This comes about in the following way: a natural regularization of the above limit is:

$$\langle\!\langle j || j' \rangle\!\rangle = \lim_{\tilde{q} \to 1} q^{c/24} \langle\!\langle j' | \tilde{q}^{\frac{1}{2}(L_0 + L_0 - \frac{c}{12})} | j \rangle\!\rangle \tag{2.9}$$

where q is the modular transform of $\tilde{q} = e^{-2\pi i/\tau}$, $q = e^{2\pi i\tau}$. In a ("unitary") theory in which the identity representation (denoted 1) is the one with the smallest conformal weight, show that in the limit $q \to 0$, the r.h.s. of (2.9) reduces to (2.8). In non unitary theories, this limiting procedure fails, but we keep (2.8) as a definition of the new norm.

At the term of this study, we have found a basis of solutions to the constraint (2.6) on boundary states, and it is thus legitimate to expand the two states attached to the two boundaries of our domain as

$$|a\rangle = \sum_{j \in \mathcal{E}} \frac{\psi_a^j}{\sqrt{S_{1j}}} |j\rangle\rangle \tag{2.10}$$

with coefficients denoted ψ_a^j , and likewise for $|b\rangle$. We define an involution $a \to a^*$ on the boundary states by $\psi_{a^*}^j = \psi_a^{j^*} = (\psi_a^j)^*$, (recall that $j \to j^*$ is an involution in \mathcal{E}). One may show [23] that it is natural to write for the conjugate state

$$\langle b| = \sum_{j \in \mathcal{E}} \langle \langle j| \frac{\psi_{b^*}^j}{\sqrt{S_{1j}}} .$$

$$(2.11)$$

As a consequence

$$\langle b \| a \rangle = \sum_{j \in \mathcal{E}} \frac{\psi_a^j \left(\psi_b^j\right)^*}{S_{1j}} \langle \langle j \| j \rangle = \sum_{j \in \mathcal{E}} \psi_a^j \left(\psi_b^j\right)^*$$
(2.12)

so that the orthonormality of the boundary states is equivalent to that of the ψ 's.

2.3. Cardy equation

Let us return to the annulus $1 \leq |\zeta| \leq e^{2\pi L/T}$ considered in last subsection, or equivalently to the cylinder of length L and perimeter T, with boundary conditions (b.c.) a and b on its two ends. Following Cardy [24], we shall compute its partition function $Z_{b|a}$ in two different ways. If we regard it as resulting from the evolution between the boundary states $|a\rangle$ and $\langle b|$, with $\tilde{q}^{\frac{1}{2}} = e^{-2\pi L/T}$, we find

$$Z_{b|a} = \langle b|(\tilde{q}^{\frac{1}{2}(L_0 + \bar{L}_0 - \frac{c}{12})}|a\rangle = \sum_{j,j' \in \mathcal{E}} \frac{\left(\psi_b^j\right)^* \psi_a^{j'}}{S_{1j}} \langle \langle j|\tilde{q}^{\frac{1}{2}(L_0 + \bar{L}_0 - \frac{c}{12})}|j'\rangle \rangle$$

$$= \sum_{j \in \mathcal{E}} \psi_a^j \left(\psi_b^j\right)^* \frac{\chi_j(\tilde{q})}{S_{1j}} .$$
 (2.13)

On the other hand, if we regard it as resulting from the periodic "time" evolution on the strip with b.c. a and b, using the decomposition (2.3) of the Hilbert space \mathcal{H}_{ba} , and with $q = e^{-\pi T/L}$

$$Z_{b|a}(q) = \sum_{i \in \mathcal{I}} n_{ib}{}^{a} \chi_{i}(q) .$$
 (2.14)

See Fig. 2. Note that string theorists would refer to these two situations as (a): the tree approximation of the propagation of a closed string; (b) the one-loop evolution of an open string. Performing a modular transformation on the



Fig. 2: Two alternative computations of the partition function $Z_{b|a}$: (a) on the cylinder, between the boundary states $|a\rangle$ and $\langle b|$, (b) as a periodic time evolution on the strip, with boundary conditions a and b.

characters $\chi_j(\tilde{q}) = \sum_i S_{ji^*} \chi_i(q)$ in (2.13), and identifying the coefficients of χ_i yields

$$n_{ia}{}^b = \sum_{j \in \mathcal{E}} \frac{S_{ij}}{S_{1j}} \psi_a^j \left(\psi_b^j\right)^* , \qquad (2.15)$$

a fundamental equation for our discussion that we refer to as Cardy equation [24]. In deriving this form of Cardy equation, we have made use of a symmetry property of $n_{ia}{}^{b}$

$$n_{ia}{}^{b} = n_{i^{*}b}{}^{a} \tag{2.16}$$

which follows from the symmetries of S.

(Comment: this identification of coefficients of specialized characters is in general not justified, as the $\chi_i(q)$ are not linearly independent. As in sect. 1, it is better to generalize the previous discussion, in a way which introduces non-specialized –and linearly independent–characters. This has been done in [3] for the case of CFTs with a current algebra. Unfortunately, little is known about other chiral algebras and their non-specialized characters.)

Let us stress that in (2.15), the summation runs over $j \in \mathcal{E}$, i.e. this equation incorporates some information on the spectrum of the theory "in the bulk", i.e. on the modular invariant partition function (1.19).

Cardy equation (2.15) is a non linear constraint relating a priori unknown complex coefficients ψ^{j} to unknown integer multiplicities $n_{ia}{}^{b}$. We need additional assumptions to exploit it.

We shall thus assume that

• we have found an orthonormal set of boundary states $|a\rangle$, i.e. satisfying

$$(n_1)_a{}^b = \sum_{j \in \mathcal{E}} \psi_a{}^j (\psi_b{}^j)^* = \delta_{ab} ; \qquad (2.17)$$

• we have been able to construct a *complete* set of such boundary states $|a\rangle$

$$\sum_{a} \psi_a{}^{j} (\psi_a{}^{j'})^* = \delta_{jj'} .$$
(2.18)

These assumptions imply that

boundary states = # independent Ishibashi states = $|\mathcal{E}|$.

None of these assumptions is innocent, and relaxing one of them forces to reconsider the forthcoming discussion. For example, the constraint of orthogonality has been recently shown to be too strong under certain circumstances: a renormalisation group flow may take us to a boundary state where it is violated [25,26,6]. In the diagonal theories at least, such a state A is a linear superposition with non-negative integer coefficients of boundary states satisfying (2.17): $\psi_A{}^j = \sum_a n_A{}^a \psi_a{}^j$. As for the assumption of completeness, it is not obviously natural to me and would also deserve a critical discussion.

2.4. Representations of the fusion algebra and graphs

Return to Cardy equation (2.15) supplemented by the above assumptions (2.17)-(2.18) and observe that it gives a decomposition of the matrices n_i , defined by $(n_i)_a{}^b = n_{ia}{}^b$, into their orthonormal eigenvectors ψ and their eigenvalues S_{ij}/S_{1j} . Observe also that as a consequence of Verlinde formula (1.13), these eigenvalues form a one-dimensional representation of the fusion algebra

$$\frac{S_{i\ell}}{S_{1\ell}} \frac{S_{j\ell}}{S_{1\ell}} = \sum_{k \in \mathcal{I}} \mathcal{N}_{ij}^{\ k} \frac{S_{k\ell}}{S_{1\ell}}, \qquad \forall i, j, \ell \in \mathcal{I} .$$

$$(2.19)$$

Hence the matrices n_i also form a representation of the fusion algebra

$$n_i n_j = \sum_{k \in \mathcal{I}} \mathcal{N}_{ij}{}^k n_k \tag{2.20}$$

and they thus commute. Moreover, as we have seen above, they satisfy $n_1 = I$, $n_i^T = n_{i^*}$.

Conversely, consider any N-valued matrix representation of the Verlinde fusion algebra n_i , such that $n_i^T = n_{i^*}$. Since the algebra is commutative,

 $[n_i, n_i^T] = [n_i, n_{i^*}] = 0$, the n_i commute with their transpose, (normal matrices), hence they are diagonalizable in a common orthonormal basis. Their eigenvalues are known to be of the form S_{ij}/S_{1j} . They may thus be written as in (2.15). Thus any such N-valued matrix representation of the Verlinde fusion algebra gives a (complete orthonormal) solution to Cardy's equation.

Conclusion:

 $\mathbb N\text{-valued}$ matrix representation of the fusion algebra

 \iff Complete, orthonormal solution of Cardy equation

Moreover, since N-valued matrices are naturally interpreted as graph adjacency matrices, graphs appear naturally and their spectral properties are those anticipated in the last lines of sect. 1.

The relevance of the fusion algebra in the solution of Cardy equation had been pointed out by Cardy himself for diagonal theories [24] and foreseen in general in [27] with no good justification; the importance of the assumption of completeness of boundary conditions was first stressed by Pradisi et al [28].

2.5. The case of $\widehat{sl}(2)$ WZW theories

Problem: Classify all N-valued matrix reps of $\widehat{s\ell}(2)_k$ fusion algebra with k fixed.

The algebra is generated recursively by n_2

$$n_1 = I,$$
 $n_2 n_i = n_{i+1} + n_{i-1},$ $i = 2, \dots, k$ (2.21)

$$S \text{ real } \Rightarrow n_i = n_i^T$$
.

Even though ψ^j and \mathcal{E} are yet unknown, we know from (1.10) that n_2 has eigenvalues of the form

$$\gamma_j = \frac{S_{2j}}{S_{1j}} = 2\cos\frac{\pi j}{k+2}, \quad j \in \mathcal{E} .$$

$$(2.22)$$

We shall discard the case where the matrix n_2 is "reducible", i.e. may be written as a direct sum of two matrices $n_2 = n_2^{(1)} \oplus n_2^{(2)}$, because then all the other matrices n_i have the same property and this corresponds to decoupled boundary conditions. It turns out that irreducible N-valued matrices G with spectrum $|\gamma| < 2$ have been classified [29]. They are the adjacency matrices

either of the A-D-E Dynkin diagrams or of the "tadpoles" $T_n = A_{2n}/\mathbb{Z}_2$. Thus as a consequence of equation (2.15) alone, for a $\hat{sl}(2)$ theory at level k, the possible boundary conditions are in one-to-one correspondence with the vertices of one of these diagrams G, with Coxeter number h = k + 2. If we remember, however, that the set \mathcal{E} must appear in one of the modular invariant torus partition functions, the case $G = T_n$ has to be discarded, and we are left with ADE. (Up to this last step, this looks like the simplest route leading to the ADE classification of $\hat{sl}(2)$ theories.) We thus conclude that for each $\hat{sl}(2)$ theory classified by a Dynkin diagram G of ADE type

$$\begin{split} \mathcal{E} &= \operatorname{Exp}(G), \qquad \dim(n_i) = |\mathcal{E}| = |G| \\ \text{complete orthonormal b. c. } &= a, b, \cdots : \text{ vertices of } G \\ n_2 &= \text{ adjacency matrix of } G \\ n_i &= ``i\text{-th fused adjacency matrix'' of } G \\ \psi^j &= \text{eigenvector of } n_2 \text{ with eigenvalue } \gamma_j \text{ .} \end{split}$$

One checks indeed that the matrices n_i , given by equation (2.15), together with (1.10), have only non negative integer elements. See [3,16] for a review of their remarkable properties and of their ubiquitous rôle in a variety of problems.

2.6. The case of c < 1 minimal models

As recalled above, this case is closely related to the $\widehat{sl}(2)$ models that we just discussed. If $c = 1 - \frac{6(p-p')^2}{pp'}$, as in sect 1.1, the classification of modular invariants is done by pairs of Dynkin diagrams $(A_{p'-1}, G)$, with p equal to the Coxeter number of G. Our problem is then to classify all N-valued matrix representations of the corresponding fusion algebra.

Theorem: [3] The only complete orthonormal solution to Cardy's equation are labelled by pairs (r, a) of nodes of the $A_{p'-1}$ and of the G graphs, with the identification

$$(r,a) \equiv (p'-r,\gamma(a)) \tag{2.24}$$

where γ is the following automorphism of the *G* Dynkin diagram: the natural \mathbb{Z}_2 symmetry for the *A*, D_{odd} and E_6 cases, the identity for the others. The solutions are given explicitly as

$$n_{rs} = N_r \otimes n_s^{(G)} + N_{p'-r} \otimes n_{p-s}^{(G)}$$
(2.25)

or

$$n_{rs;(r_1,a)}{}^{(r_2,b)} = N_{rr_1}{}^{r_2} n_{sa}^{(G)b} + N_{p'-rr_1}{}^{r_2} n_{p-sa}^{(G)b} , \qquad (2.26)$$

with $1 \leq r, r_1, r_2 \leq p'-1$, $1 \leq s \leq p-1$, and a, b are running over the nodes of the Dynkin diagram G; N_r are the fusion matrices of $\widehat{sl}(2)_{p'-2}$, $n_s^{(G)}$ are the representation matrices of the fusion algebra of $\widehat{sl}(2)_{p-2}$ associated with the graph G and introduced in sect 2.5.

Hint: Look at the generators n_{21} and n_{12} of fusion algebra. Both have eigenvalues $|\gamma| < 2$, but now they are no longer irreducible matrices as in sect. 2.5. To show that (2.25) is indeed the only solution requires a detailed discussion, see [3].

Example: The 3-state Potts model. This is the p' = 5, p = 6 (c = 4/5) minimal model, classified by the pair (A_4, D_4) . According to the previous discussion, there are 8 distinct boundary conditions labelled by

$$(r, a) \in (A_4, D_4)/\mathbb{Z}_2$$
 e.g. $r = 1, 2, a \in D_4$

This was first pointed out in [30]: the three boundary conditions (1, a) where a = 1, 3, 4 is one of the three end-points of the D_4 diagram are fixed b.c., corresponding to fixing the value of the Potts "spin" to one of its three values, while in the three b.c. (2, a), the boundary spin may take either value different from a; the b.c. (1, 2), where 2 denotes the middle point of the diagram, is the free boundary condition: the boundary spin may take an arbitrary value; finally the b.c. (2, 2) is more delicate to describe [30]. See also Oshikawa's lectures at this school. The explicit expressions of the different partition functions of type $Z_{(1,1)|(r,a)}$ read

$$\begin{split} &Z_{(1,1)|(1,1)} = \chi_{(1,1)} + \chi_{(1,5)} \\ &Z_{(1,1)|(1,2)} = \chi_{(4,2)} + \chi_{(4,4)} \\ &Z_{(1,1)|(1,a)} = \chi_{(1,3)} \quad \text{if } a = 3,4 \\ &Z_{(1,1)|(2,1)} = \chi_{(3,1)} + \chi_{(3,5)} \\ &Z_{(1,1)|(2,2)} = \chi_{(2,2)} + \chi_{(2,4)} \\ &Z_{(1,1)|(2,a)} = \chi_{(3,3)} \quad \text{if } a = 3,4 \end{split}$$

The others are linear combinations with non-negative integer coefficients of the latter. See also [31,32] for a lattice realization with integrable boundary weights.

2.7. Other cases

It should be clear that the situation that we have described in detail for sl(2) extends to all RCFTs. The matrices n_i solutions to Cardy equation are the adjacency matrices of graphs. In the case of $\widehat{sl}(N)$, it is sufficient to supply the (N-1) fundamental matrices n_p , $p = 1, \dots, N-1$, indexed by the Young \vdots

tableaux of the fundamental representations of sl(N), to determine all of them. The fact that all n_i have non negative integer elements is then non trivial. By Cardy equation again, they satisfy a very restrictive spectral property: their eigenvalues must be of the form S_{ij}/S_{1j} , when j runs over the set \mathcal{E} , i.e. the diagonal part of the modular invariant.

The program of classifying these graphs/boundary conditions has been completed only in a few cases: $\hat{sl}(2)$ as discussed above, $\hat{sl}(3)$ through a combination of some empirical search of relevant graphs [33], of Gannon's classification of the modular invariants [13], and of the recent work of Ocneanu [34], see [3,16] for a discussion; $\hat{sl}(N)_1$ [3], where the results match those obtained in the study of modular invariants [14]: the graphs turn out to be star polygons.

3. Boundary Operator Algebra

According to Cardy [24], changes of boundary conditions can be interpreted as due to the insertion of fields ${}^{b}\Psi_{j,\beta}^{a}(x)$ living on the boundary, $\Im m z = 0$, $x = \Re e z$ of the upper half-plane $z \in \mathbb{H}$. We know the spectrum of these fields from the previous discussion: for a given pair a, b of boundary conditions and a label $i \in \mathcal{I}$, there are $n_{ia}{}^{b}$ independent such fields, which are thus labelled by a multiplicity label $\alpha = 1, \dots, n_{ia}{}^{b}$. Pictorially, we may again use a $i \downarrow$ CVO-like representation for ${}^{b}\Psi_{j,\alpha}^{a}(x) = b_{\alpha x} a$. It is a natural –and physically important– question to determine the correlation functions of these new fields in the possible presence of the "usual" fields "in the bulk". Two quantities of particular importance are

i) the fusion matrix ${}^{(1)}F$ of boundary operators, which plays for the boundary fields the same rôle as the matrix F for the CVO (sect. 1.1),

$$\frac{i}{a_{1}x_{1}} \frac{j}{a_{2}x_{2}} \xrightarrow{(^{1})} F_{cp} \xrightarrow{b} \frac{p}{\beta x_{2}} \frac{j}{x_{12}} , \text{ gives their OPE coefficient} \\
^{b}\Psi_{i,\alpha_{1}}^{c}(x_{1}) {}^{c}\Psi_{j,\alpha_{2}}^{a}(x_{2}) = \sum_{p,\beta,t} {}^{(1)}F_{cp} \begin{bmatrix} i & j \\ b & a \end{bmatrix}_{\alpha_{1}\alpha_{2}}^{\beta} \frac{t}{|x_{12}|^{h_{i}+h_{j}-h_{p}}} {}^{b}\Psi_{p,\beta}^{a}(x_{2}) +$$
(2.1)

(3.1) in which the multiplicity labels run over $\alpha_1 = 1, \dots, n_{ic}{}^b, \alpha_2 = 1, \dots, n_{ja}{}^c, \beta = 1, \dots, n_{pa}{}^b, t = 1, \dots, \mathcal{N}_{ij}{}^p;$

ii) the bulk-boundary coefficients $R_{a,\alpha}^{(i,\bar{i}^*,t)}(p)$ (denoted ${}^{a,\alpha}B_{(i,\bar{i})}^{p,t}$ in [3]) enable one to expand bulk fields Φ in terms of Ψ , close to the boundary, i.e. for small $y = \Im m z$

$$\Phi_{(i,\bar{i})}(z,\bar{z}) = \sum_{a,\alpha,p\in\mathcal{I},t} R^{(i,\bar{i}^*,t)}_{a,\alpha}(p) \frac{1}{(2iy)^{h_i + h_{\bar{i}} - h_p}} {}^a \Psi^a_{p,\alpha}(x) + \dots$$
(3.2)

Here $\alpha = 1, \dots, n_{pa}{}^{a}, t = 1, \dots, \mathcal{N}_{i\bar{i}}{}^{p^*}$. Given these data (and for a chosen normalisation of the Ψ), we may in principle compute all correlation functions of Ψ and Φ [35].

I shall not dwell here on the determination of the fusion matrices ${}^{(1)}F$ and of the bulk-boundary coefficients. They have been the object of much activity lately, in particular on their connection with chiral data and on the algebraic relations that they satisfy. Their explicit determination for the A or D type minimal models has been completed by Runkel [36,37]. I just mention two results:

In the diagonal theories, the matrices ${}^{(1)}F$ coincide with the standard fusion matrices F: this is in accordance with the fact that in that case, the indices a, b, \cdots are of the same nature as i, j, \cdots , i.e. also belong to the set \mathcal{I} [36,3]. Also, the bulk coefficients R(p) are expressible in terms of the matrix S(p)introduced at the end of sect. 1.1, a quite remarkable convergence between seemingly very different objects.

In general, the bulk-boundary coefficient pertaining to the identity, i.e. $R_{a,1}^{(i,i^*,1)}(1)$ is proportional to $\psi_a{}^i/\psi_a{}^1$. It thus satisfies, up to a normalisation, the Pasquier algebra (1.22), [28,23,22,3]

$$\frac{\psi_a{}^i}{\psi_a{}^1}\frac{\psi_a{}^j}{\psi_a{}^1} = \sum_{k\in\mathcal{I}} M_{ij}{}^k\frac{\psi_a{}^k}{\psi_a{}^1}$$

This expresses the OPA of bulk fields of type $\Phi_{(i,i^*)}$ near the boundary (a) and must be compared with the empirical observation made in sect. 1.2 on the rôle of the Pasquier algebra in the OPA of spinless fields.

The determination of the ${}^{(1)}F$ matrix, whose entries are called "cells" by Ocneanu [38], turns out to be an essential step, not only in the study of boundary effects, but also in uncovering hidden algebraic aspects of the theory. For a given set of matrices $\{n_i\}$ –a given "graph" – this set of cells satisfies various non linear relations: orthonormality expressing the fact that it plays the rôle of a change of basis ("3 – *j* coefficients") in tensors products in a certain space, a mixed pentagon identity written symbolically as $F^{(1)}F^{(1)}F = {}^{(1)}F^{(1)}F$, expressing the consistency (associativity) in the fusion of several boundary fields,



Fig. 3: The mixed pentagon identity expressing the associativity of the fusion of boundary fields.

(see Fig. 3), and other identities. See $\left[34,39,40\right]$ for more details and references.

It is thus a non trivial task to determine ${}^{(1)}F$ for a given graph: it may fail, in certain cases, because of some obstruction not revealed by the study of the set $\{n_i\}$ alone. This is what has been announced by Ocneanu [34] for one of the graphs of $\hat{sl}(3)$ listed in [3].

If it exists, this system of cells gives us access not only to the Boundary CFT, as explained above. It turns out to also contain the essential information about integrable lattice models and their Boltzmann weights.

4. Integrable lattice models

It should indeed be mentioned that parallel to the conformal field theoretic discussion sketched in these notes, one may study a class of lattice integrable models, the so-called face, or height, or RSOS, models, which are also described in terms of the same graphs. There the degrees of freedom are attached to sites of a square lattice, and are assigned to take their value in the set of vertices

of the chosen graph. Typically, in the simplest models, neighbouring sites on the lattice are assigned neighbouring vertices on the graph. Boltzmann weights are given for each configuration of four vertices around a square face. They depend on an additional parameter, the spectral parameter u, and must satisfy an integrability condition, the celebrated Yang-Baxter equation. This is realised algebraically through a representation of the Temperley-Lieb algebra, or of some other quotient of the Hecke algebra, constructed on the graph as follows. For a triplet of sites $\mathbf{n} - 1, \mathbf{n}, \mathbf{n} + 1$ along a diagonal zig-zag line on the lattice, and "heights" a, b or d and c assigned to them, the face Boltzmann weight reads

The Us are the generators of the Hecke algebra, i.e. satisfy

$$U_{\mathbf{n}}^{2} = 2 \cos \frac{\pi}{h} U_{\mathbf{n}}$$
$$U_{\mathbf{n}}U_{\mathbf{n}+1}U_{\mathbf{n}} - U_{\mathbf{n}} = U_{\mathbf{n}+1}U_{\mathbf{n}}U_{\mathbf{n}+1} - U_{\mathbf{n}+1}$$
$$U_{\mathbf{n}}U_{\mathbf{m}} = U_{\mathbf{m}}U_{\mathbf{n}} \quad \text{if } |\mathbf{n} - \mathbf{m}| \ge 2 .$$
(4.2)

As a consequence, the face weights satisfy the Yang-Baxter identity

$$X_{\mathbf{n}}(u)X_{\mathbf{n}+1}(u+v)X_{\mathbf{n}}(v) = X_{\mathbf{n}+1}(v)X_{\mathbf{n}}(u+v)X_{\mathbf{n}+1}(u) .$$
(4.3)

The Pasquier models [41] give the simplest and most explicit example: the relevant graphs are (dare I say of course?) the ADE diagrams, and the U matrix element reads

$$(U_{\mathbf{n}})_{bd} = \delta_{ac} \frac{\sqrt{\psi_b^1 \psi_d^1}}{\psi_a^1} .$$

$$(4.4)$$

They are related to the sl(2) algebra. Although generalisations to higher rank are known to exist, there is a lack of such explicit and general formulas. In general, there are additional edge degrees of freedom $\alpha, \gamma, \alpha', \gamma', \alpha = 1, \dots, n_{\Box a}{}^{b}$, etc. The weights for the (generalised) A-type graphs are known from the work of Jimbo et al, and Wenzl [42]. In [27,43] explicit expressions have been given for models associated with some of the graphs of $\hat{sl}(3)$, and a general result has been obtained in [44] for graphs of $\hat{sl}(N)$ corresponding to conformal embeddings. The recent observation by Ocneanu [34] that one may write the above generator of the Hecke algebra as

$$U_{bd}{}^{\gamma'\alpha'}_{\gamma\alpha} = \sum_{\beta} {}^{(1)}F_{b}_{\square} \left[{}^{\square \square}_{c a} \right]^{\beta \ 1}_{\gamma \ \alpha} {}^{(1)}F^{*}_{d\square} \left[{}^{\square \square}_{c a} \right]^{\beta \ 1}_{\gamma' \ \alpha'}.$$
(4.5)

in terms of the ${}^{(1)}F$ matrix previously introduced is thus a significant progress, both practically and conceptually, since it connects problems of apparently different nature.

Finally, in these lattice models, it is legitimate to wonder if boundaries may be introduced without spoiling integrability. This requires a careful determination of the boundary Boltzmann weights, satisfying the so-called Boundary Yang-Baxter Equation [4]. This has now been completed for the unitary minimal models: a large class of boundary weights has been found, which at criticality, match perfectly what we have learnt from BCFT [32]. It remains to connect these boundary weights with quantities defined previously in the context of BCFT to have a fully consistent and unified picture of all questions of boundary conditions in integrable lattice models and conformal theories.

Acknowledgements

It is a pleasure to thank the APCTP (Asia Pacific Center for Theoretical Physics) and the organisers of this meeting, Changrim Ahn, Chaiho Rim and Ryu Sasaki, for their invitation and hospitality in the beautiful island of Cheju and for the stimulating environment of the school and workshop. The work presented here results from several enjoyable collaborations with R. Behrend, P. Pearce and V. Petkova. A large part of the redaction of these notes was carried out at the CERN TH Division, which I am happy to thank for hospitality and support.

References

- C. Schweigert, lectures at the Summer School Nonperturbative Quantum Field Theoretic Methods and their Applications, 19-21 August 2000, Eötvös University, Budapest, Hungary, to appear.
- [2] H. Saleur, Lectures on Non Perturbative Field Theory and Quantum Impurity Problems, I and II, cond-mat 9812110, cond-mat 0007309.
- [3] R.E. Behrend, P.A. Pearce, V.B. Petkova and J.-B. Zuber, Nucl. Phys. B 579 (2000) 707-773, hep-th 9908036.
- [4] R.E. Behrend and P.A. Pearce, J. Phys. A 29 (1996) 7827-7835; Int. J. Mod. Phys. 11 (1997) 2833-2847.
- [5] P. Dorey, lectures at this school.
- [6] G. Watts, lectures at Budapest Summer School, op. cit., to appear;
 K. Graham, I. Runkel and G.M.T. Watts, Renormalisation group flows of boundary theories, to appear in the Proceedings of the TMR network conference Non perturbative Quantum Effects 2000, hep-th 0010082.
- [7] V. Kac, Infinite dimensional algebras, Cambridge University P.; V.G. Kac and D.H. Peterson, Adv. Math. 53 (1984) 125-264;
- J. Fuchs, Affine Lie Algebras and Quantum Groups, Cambridge Univ. Pr. 1992.
- [8] P. Goddard, A. Kent and D. Olive, Comm. Math. Phys. 103 (1986) 105-119.
- [9] P. Di Francesco, P. Mathieu and D. Sénéchal, *Conformal Field Theory*, Springer Verlag 1997.
- [10] E. Verlinde, Nucl. Phys. B300 [FS22] (1988) 360-376.
- [11] G. Moore and N. Seiberg, Nucl. Phys. B313 (1989) 16; Comm. Math. Phys. 123 (1989) 177-254.
- [12] A. Cappelli, C. Itzykson and J.-B. Zuber, Nucl. Phys. B280 [FS18] (1987) 445-465; Comm. Math. Phys. 113 (1987) 1-26;
 A. Kato, Mod. Phys. Lett. A2 (1987) 585-600.
- T. Gannon, Comm. Math. Phys. 161 (1994) 233-263; The Classification of SU(3) Modular Invariants Revisited, hep-th 9404185.
- [14] C. Itzykson, Nucl. Phys. (Proc. Suppl.) 5B (1988) 150-165;
 P. Degiovanni, Comm. Math. Phys. 127 (1990) 71-99.
- [15] T. Gannon, The monstruous moonshine and the classification of CFT, hep-th 9906167.
- [16] J.-B. Zuber, CFT, BCFT, ADE and all that, proceedings of the Bariloche Summer School, Jan 2000, eds R. Coquereaux and R. Trinchero, hep-th 0006151.

- 26 J.-B. Zuber
- [17] Vl.S. Dotsenko and V.A. Fateev, Nucl. Phys. B 251 [FS13] (1985) 691-734.
- [18] A.B. Zamolodchikov and V.A. Fateev, Sov. Phys. JETP 62 (1985) 215-225; Sov. J. Nucl. Phys. 43 (1986) 657-664.
- [19] V.B. Petkova and J.-B. Zuber, Nucl. Phys. B 438 (1995) 347-372, hep-th 9410209.
- [20] V. Pasquier, J. Phys. A20 (1987) 5707-5717.
- [21] G. Watts, private communication.
- [22] J. Fuchs and C. Schweigert, Nucl. Phys. B 530 (1998) 99-136, hep-th 9712257.
- [23] A. Recknagel and V. Schomerus, Nucl. Phys. B 531 (1998) 185-225.
- [24] J.L. Cardy, Nucl. Phys. 324 (1989) 581-596.
- [25] L. Chim, Int. J. Mod. Phys. A11 (1996) 4491-4512, hep-th 9510008;
 I. Affleck, Edge critical behaviour of the 2-dimensional tri-critical Ising model, cond-mat 0005286.
- [26] A. Recknagel, D. Roggenkamp, V. Schomerus, On relevant boundary perturbations of unitary minimal models, hep-th 0003110.
- [27] P. Di Francesco and J.-B. Zuber, Nucl. Phys. B338 (1990) 602-646.
- [28] G. Pradisi, A. Sagnotti and Ya.S. Stanev, Phys. Lett. 381 (1996) 97-104.
- [29] F.M. Goodman, P. de la Harpe and V.F.R. Jones, Coxeter Graphs and Towers of Algebras, Springer-Verlag, Berlin (1989).
- [30] I. Affleck, M. Oshikawa and H. Saleur, J. Phys A 31 (1998) 5827-5842, cond-mat 9804117.
- [31] R.E. Behrend, P.A. Pearce and J.-B. Zuber, J.Phys. A31 (1998) L763-L770, hep-th 9807142.
- [32] R.E. Behrend and P.A. Pearce, Integrable and Conformal Boundary Conditions for $\hat{sl}(2) \ A - D - E$ Lattice Models and Unitary Conformal Field Theories, hep-th 0006094, J. Stat Phys. to appear.
- [33] P. Di Francesco and J.-B. Zuber, in *Recent Developments in Conformal Field Theories*, Trieste Conference 1989, S. Randjbar-Daemi, E. Sezgin and J.-B. Zuber eds., World Scientific 1990;

P. Di Francesco, Int. J. Mod. Phys. A7 (1992) 407-500.

- [34] A. Ocneanu, lectures at the school "Quantum Symmetries in Theoretical Physics and Mathematics", January 2000, Bariloche, R. Coquereaux and R. Trinchero eds, to appear.
- [35] J.L. Cardy and D.C. Lewellen, *Phys. Lett. B* 259 (1991) 274-278;
 D.C. Lewellen, *Nucl. Phys.* B 372 (1992) 654-682.
- [36] I. Runkel, Nucl. Phys. B 549 (1999) 563-578, hep-th 9811178.

- [37] I. Runkel, Nucl. Phys. B 579 (2000) 561-589, hep-th 9908046.
- [38] A. Ocneanu, Paths on Coxeter Diagrams, in Lectures on Operator Theory Fields Institute Monographies, Rajarama Bhat et al edrs, AMS 1999.
- [39] V.B. Petkova and J.-B. Zuber, BCFT: from the boundary to the bulk, to appear in the Proceedings of the TMR network conference Nonperturbative Quantum Effects 2000, hep-th 0009219.
- [40] V.B. Petkova and J.-B. Zuber, *The many faces of Ocneanu triangular cells*, in preparation.
- [41] V. Pasquier, Nucl. Phys. B285 [FS19] (1987) 162-172.
- [42] M. Jimbo, T. Miwa and M. Okado, Lett. Math. Phys. 14 (1987) 123-131; Comm. Math. Phys. 119 (1988) 543-565;
 H. Wenzl, Inv. Math. 92 (1988) 349-383.
- [43] N. Sochen, Nucl. Phys. B360 (1991) 613-640.
- [44] F. Xu, Comm. Math. Phys. 192 (1998) 349-403.