Advanced Statistical Physics

Leticia F. Cugliandolo

Sorbonne Université

Institut Universitaire de France

leticia@lpthe.jussieu.fr
www.lpthe.jussieu.fr/~leticia

Disorder

Randomness

Impurities

No material is perfect and totally free of impurities

(vacancies, substitutions, amorphous structures, etc.)

First distinction

- Weak randomness : phase diagram respected, criticality may change
- Strong randomness : phases modified

Second distinction

- Annealed : fluctuating (easier)
- Quenched : frozen, static (harder)



Quenched disorder

Variables frozen in time-scales over which other variables fluctuate.

Time scales

 $au_0 \ll t_{
m obs} \ll au_{
m eq}^{qd}$

 au_{eq}^{qd} could be the diffusion time-scale for magnetic impurities the magnetic moments of which will be the variables of a **magnetic system**;

or the flipping time of impurities that create random fields acting on other magnetic variables.

Weak disorder (modifies the critical properties but not the phases) *vs.* strong disorder (that modifies both).

e.g. random ferromagnets vs. spin-glasses.

Geometrical problems

Random graphs & Percolation







p = 0.1

p = 0.25

p = 0.5



Neural Networks

Real neural network



Neurons connected by synapsis on a random graph

Figures from AI, Deep Learning, and Neural Networks explained, A. Castrounis

Neural Networks

Sketch & artificial network



The connections in \mathbf{w}^T may have a random component

The state of the neuron up (firing), down (quiescent) is a result of the calculation In the artificial network on chooses the geometry (number of nodes in internal layer, number of hidden layers, connections between layers)

Figures from AI, Deep Learning, and Neural Networks explained, A. Castrounis



Magnetic impurities (spins) randomly placed in an inert host

 $\vec{r_i}$ are random and time-independent since

the impurities do not move during experimental time-scales \Rightarrow

quenched randomness





spins can flip but not move

RKKY potential

$$V(r_{ij}) \propto rac{\cos 2k_F r_{ij}}{r_{ij}^3} s_i s_j$$

very rapid oscillations about 0 positive & negative slow power law decay.



Models on a lattice with random couplings

Ising (or Heisenberg) spins $s_i = \pm 1$ sitting on a lattice

 J_{ij} are random and time-independent since

the impurities do not move during experimental time-scales \Rightarrow

quenched randomness

Magnetic impurities in a metal host



spins can flip but not move

Edwards-Anderson model

$$H_J[\{s_i\}] = -\sum_{\langle ij\rangle} J_{ij} s_i s_j$$

 J_{ij} drawn from a pdf with zero mean & finite variance

Neural networks

Models on graphs with random couplings

The neurons are Ising spins $s_i = \pm 1$ on a graph

 J_{ij} are random and time-independent since

the synapsis do not change during experimental time-scales \Rightarrow

quenched randomness



spins can flip but not move

Hopfield model

$$H_J[\{s_i\}] = -\sum_{\langle ij\rangle} J_{ij}s_is_j$$

memory stored in the synapsis

$$J_{ij} = 1/N_p \sum_{\mu=1}^{N_p} \xi_i^{\mu} \xi_j^{\mu}$$

the patterns ξ_i^{μ}

are drawn from a pdf with

zero mean & finite variance

Constrained satisfaction problems

Problems involving variables which must satisfy some constraints

e.g. equalities, inequalities or both

studied in computer science to

compute their complexity or develop algorithms to most efficiently solve them

Typically, N variables, which have to satisfy M constraints.

e.g. the variables could be the weights of a neural network, and each constraint imposes that the network satisfies the correct input-output relation on one of M training examples (e.g. distinguishing images of cats from dogs).

Statistical physics approach

thermodynamic limit $N
ightarrow \infty$ and $M
ightarrow \infty$ with lpha = M/N finite

K-Satisfiability

The problem is to determine whether the variables of a given Boolean formula F can be assigned in such a way to make the formula evaluate to TRUE (satisfied) Example. Call the variable x

We use x for the evaluation x = TRUE and \overline{x} for the requirement x = FALSE

Take the formula $F = C_1 : x_1 \text{ OR } \overline{x}_2$ made by a single clause C_1

it is satisfiable because one can find the values $x_1 = \text{TRUE}$ (and x_2 free) or $x_2 = \text{FALSE}$ (and x_1 free), which make $C_1 : x_1 \text{ OR } \overline{x}_2 \text{ TRUE}$

This formula is so simple that 3 out of 4 possible configurations of the two variables solve it. This example belongs to the k = 2 class of satisfiability problems since the clause is made by two literals (involving different variables) only. It has M = 1 clauses and N = 2 variables.

K-Satisfiability

Harder to decide formulæ are made of M clauses involving k literals required to take the true value (x) or the false value (\overline{x}) each, these taken from a pool of N variables. An example in k = 3-SAT is

$$F = \begin{cases} C_1 : x_1 \text{ OR } \overline{x}_2 \text{ OR } x_3 \\ C_2 : \overline{x}_5 \text{ OR } \overline{x}_7 \text{ OR } x_9 \\ C_3 : x_1 \text{ OR } \overline{x}_4 \text{ OR } x_7 \\ C_4 : x_2 \text{ OR } \overline{x}_5 \text{ OR } x_8 \end{cases}$$

All clauses have to be satisfied simultaneously so the formula has to be read

${\sf F}:C_1 \; {\sf AND} \; C_2 \; {\sf AND} \; C_3 \; {\sf AND} \; C_4$

When $\alpha \equiv M/N \gg 1$ the problems typically become unsolvable while many solutions exist for $\alpha \ll 1$. A sharp threshold at α_c for $N, M \to \infty$

Random K-Satisfiability

An instance of the problem, i.e. a formula F, is chosen at random with the following procedure :

First one takes k variables out of the N available ones.

Second one decides to require x_i or \overline{x}_i for each of them with probability 1/2

Third one creates a clause taking the OR of these k literals.

Forth one returns the variables to the pool and the outlined three steps are repeated M times.

The M resulting clauses form the final formula.

Change of focus from worse case (most difficult formula) to typical case (just one such constructed formula)

Random K-Satisfiability as a physical model

Boolean variables \Rightarrow Ising spins

 x_i evaluated to TRUE (FALSE) corresponds to $s_i = 1 \ (-1)$

The requirement that a formula be evaluated TRUE by an assignment of variables (i.e. a configuration of spins) \Rightarrow ground state of an adequately chosen energy function = cost function

In the simplest setting, each clause will contribute zero (when satisfied) or one (when unsatisfied) to this cost function.

There are several equivalent ways to reach this goal. The fact that the variables are linked together through the clauses suggests to define k-uplet interactions between them.

Random K-Satisfiability as a physical model

A way to represent a clause in an energy function, for instance,

 C_1 : x_1 OR \overline{x}_2 OR x_3

as an interaction between spins. In this case

 $(1-s_1)(1+s_2)(1-s_3)/8$

This term vanishes if $s_1 = 1$ or $s_2 = -1$ or $s_3 = 1$ and does not contribute to the total energy, that is written as a sum of terms of this kind.

It is then simple to see that the total energy can be rewritten in a way that resembles strongly physical spin models,

$$2^{K}H_{J}[\{s_{i}\}] = M + \sum_{R=1}^{K} (-1)^{R} \sum_{i_{1} < \dots < i_{R}} J_{i_{1} \dots i_{R}} s_{i_{1}} \dots s_{i_{R}}$$
$$J_{i_{1} \dots i_{R}} = \sum_{a=1}^{M} J_{ai_{1}} \dots J_{ai_{R}} \text{ and } J_{ai} = \pm 1 \text{ according to } x_{i} \text{ or } \overline{x}_{i} \text{ in clause } a$$
and zero otherwise

K-Satisfiability & complexity theory

Special interest in computational complexity theory

K-Sat for K ≥ 3 is in the NP complexity class

No algorithm (as yet) has been found that can find an assignment for the variables x_i in polynomial time

one can verify in polynomial time whether an assignment satisfies the given formula

K-SAT is an NP-complete problem

all other problems in the NP complexity class can be formally reduced to the K-SAT problem

Pinning by impurities

Competition between elasticity and quenched randomness

d-dimensional elastic manifold in a transverse N-dimensional **quenched** random potential.



Interface between two phases; vortex line in type-II supercond; stretched polymer.

Distorted Abrikosov lattice



Goa et al. 01

Randomness

Properties

- Spatial inhomogeneity
- Frustration

(spectrum pushed up, degeneracy of ground state)

- Probability distribution of couplings, fields, etc.
- Self-averageness

Frustration

Properties



Frustration enhances the ground-state energy and entropy

One can expect to have **metastable states** too

One cannot satisfy all couplings simultaneously if

$$\prod_{loop} J_{ij} < 0$$

Heterogeneity

Each variable, spin or other, feels a different local field, $h_i = \sum_{j=1}^{z} J_{ij} s_j$,

contrary to what happens in a ferromagnetic sample, for instance.



Each sample is *a priori* different but,

do they all have a different thermodynamic and dynamic behavior?

The disorder-induced free-energy density distribution approaches a Gaussian with vanishing dispersion in the thermodynamic limit :

 $\lim_{N \to \infty} f_N(\beta, J) = f_\infty(\beta)$ independently of disorder

— Experiments : all *typical* samples behave in the same way.

— Theory : one can perform a (hard) average of disorder, [...],

 $-\beta N f_{\infty}(\beta) = \lim_{N \to \infty} [\ln \mathcal{Z}_N(\beta, J)]$

From here, we see that, e.g., the energy density is self-averaging

Replica theory

$$-\beta f_{\infty}(\beta) = \lim_{N \to \infty} \lim_{n \to 0} \frac{\left[\mathcal{Z}_{N}^{n}(\beta, J)\right] - 1}{Nn}$$

The question

Given two samples with different quenched randomness

(e.g. different interaction strengths J_{ij} s or random fields h_i)

but drawn from the same (kind of) distribution

is their behaviour going to be totally different?

Which quantities are expected to be the same and which not?

Observables & distributions

Given a quantity A_J , which depends on the quenched randomness J, it is distributed according to

$$P(A) = \int dJ \ p(J) \ \delta(A - A_J)$$

This pdf is expected to be narrower and narrower (more peaked) as $N \rightarrow \infty$

Therefore, one will observe $A_{typ} = A s.t. \max_A P(A)$

However, it is difficult to calculate A_{typ} , what about calculating $[A] = \int dAP(A)A$?

Warm-up exercise

Exercise 5.1 This exercise provides a useful example of the distinction between *typical* and *average* values of random variables. Consider a random variable z that takes only two values $z_1 = e^{\alpha \sqrt{N}}$ and $z_2 = e^{\beta N}$, with α and β two positive and finite numbers with α unconstrained and $\beta > 1$. The probabilities of the two events are $p_1 = 1 - e^{-N}$ and $p_2 = e^{-N}$. First, confirm that these probabilities are normalised. Second, compute the average $\langle z \rangle$, where the angular brackets indicate average with the probabilities p_1, p_2 , and evaluate it in the limit $N \to \infty$. Third, calculate the most probable value taken by z, that we call z_{typ} , for typical (indeed, if we were to draw the variable we would typically get this value). Compare and conclude. Now, let us study the behaviour of the quantity $\ln z$ that is also a random variable. Compute its average. By which value of z is it determined? Does $\langle \ln z \rangle = (\ln z)_{typ}$ in the large N limit? Is $\langle \ln z \rangle = \ln \langle z \rangle$? The last result demonstrates the difference between what are called *quenched* and *annealed* averages. Which value is larger? Does the comparison comply with Jenssen's inequality? (See App. 5.A for its definition.)

A function is convex function iff $\forall x_1, x_2$ and $t \in [0, 1]$:

$$f(tx_1 + (1-t)x - 2) \le tf(x_1) + (1-t)f(x_2)$$
.

Warm-up exercise



Warm-up exercise



Example : the disordered Ising chain

$$H_J[\{s_i\}] = -\sum_i J_i s_i s_{i+1}$$
 J_i distributed $p(J_i)$ with any pdf

Compute the partition function Z_J by introducing $\sigma_i = s_i s_{i+1}$

$$Z_{J}[\{s_{i}\}] = \sum_{s_{i}=\pm 1} e^{\beta \sum_{i} J_{i} s_{i} s_{i+1}} = \sum_{\sigma_{i}=\pm 1} e^{\beta \sum_{i} J_{i} \sigma_{i}} = \prod_{i=1}^{N} 2 \coth \beta J_{i}$$

(boundary condition effects negligible for $N \to \infty$)

It is a **product** of N random numbers

The free-energy is $-\beta F_J[\{s_i\}] = \sum_{i=1}^N \ln \coth \beta J_i + N \ln 2$

It is a sum of N random numbers

Example : the disordered Ising chain

$$H_J[\{s_i\}] = -\sum_i J_i s_i s_{i+1}$$
 J_i distributed $p(J_i)$ with any pdf

The partition function & the free energy density are different objects

$$Z_J[\{s_i\}] = \prod_{i=1}^N 2 \coth \beta J_i \qquad -\beta f_J[\{s_i\}] = \frac{1}{N} \sum_{i=1}^N \ln \coth \beta J_i + \ln 2$$

Take J_i to be *i.i.d* with zero mean $[J_i] = 0$ & finite variance $[J_i^2] = \sigma^2$ and use the **Central Limit Theorem** :

 $X = \frac{1}{N} \sum_i x_i$ is Gaussian distributed with average $\langle X \rangle = \langle x_i \rangle$ and variance $\langle (X - \langle X \rangle)^2 \rangle = \sigma^2 / N$

Therefore f_J is Gaussian distributed and its variance vanishes for $N \to \infty$ Moreover, $f_J^{\mathrm{typ}} = [f_J]$

Systems with short-range interactions

Divide a, say, cubic system of volume $V = L^d$ in n sub-cubes, of volume $v = \ell^d$ with V = nv



Systems with short-range interactions

For short-range interactions the total free-energy is the sum of two terms, a contribution from the bulk of the subsystems and a contribution from the interfaces between the subsystems :

$$-\beta F_J = \ln Z_J = \ln \sum_{\text{conf}} e^{-\beta H_J(\text{conf})} \approx \ln \sum_{\text{conf}} e^{-\beta H_J(\text{bulk}) - \beta H_J(\text{surf})}$$
$$= \ln \sum_{\text{bulk}} e^{-\beta H_J(\text{bulk})} + \ln \sum_{\text{surf}} e^{-\beta H_J(\text{surf})} = -\beta F_J^{\text{bulk}} - \beta F_J^{\text{surf}}$$

where the \approx indicates that we dropped the contributions of interactions between the bulk and the interfaces (surf)

Systems with short-range interactions

If the interaction extends over a short distance l and the linear size of the boxes is $\ell \gg l$, we also assume that the surface energy is negligible with respect to the bulk one (same for possible entropic contributions) and

$$-\beta F_J \approx -\beta F_J^{\text{bulk}} = \ln \sum_{\text{bulk}} e^{-\beta H_J(\text{bulk})}$$

The disorder dependent free-energy is a sum of $n = (L/\ell)^d$ independent random numbers, each one being the disorder dependent free-energy of the bulk of each subsystem :

$$-\beta F_J \approx \sum_{k=1}^n \ln \sum_{\text{bulk}_k} e^{-\beta H_J(\text{bulk}_k)}$$

In the limit of a very large number of subsystems ($L \gg \ell$ or $n \gg 1$) the CLT \Rightarrow the free-energy density is Gaussian distributed with

 $f_J^{\text{typ}} = [f_J]$

Systems with short-range interactions

The dispersion about the typical value of the total free-energy vanishes in the large n limit, $\sigma_{F_J}/[F_J] \propto \sqrt{n}/n = n^{-1/2} \to 0$

The one of the free-energy density, or intensive free-energy, $f_J=F_J/N$, as well, $\sigma_{f_J}/[f_J]=O(n^{-1/2})$

In a sufficiently large system the typical free-energy density f_J^{typ} is then very close to the averaged $[f_J]$ and one can compute the latter to understand the static properties of typical systems.

Much easier to do analytically. More later.

Failure and quenched vs. Annealed

Go back to the one dimensional disordered Ising chain and show that

the partition function and the spatial correlations

are not self-averaging.

The annealed free-energy is defined as $-\beta F^{\text{annealed}} = \ln[Z_J]$

The quenched free-energy is defined as $-\beta F^{\text{quenched}} = [\ln Z_J]$

Jenssen's inequality applied to the convex function $-\ln y$ implies

 $-\ln[Z_J] \le -[\ln Z_J]$

and for the free-energies one deduces

 $F^{\text{annealed}} = -\beta^{-1} \ln[Z_J] \le -\beta^{-1} [\ln Z_J] = F^{\text{quenched}}$

Methods

disordered systems

Statics

TAP Thouless-Anderson-Palmer

Replica theory

Cavity or Peierls approx.

Bubbles & droplet arguments functional RG¹

fully-connected (complete graph)

Gaussian approx. to field-theories

dilute (random graph)

finite dimensions

Dynamics

Generating functional for classical field theories (MSRJD).

Schwinger-Keldysh closed-time path-integral for quantum dissipative models (the previous is recovered in the $\hbar \rightarrow 0$ limit).

Perturbation theory, renormalization group techniques, self-consistent approx.

Randomness

Properties

— Spatial inhomogeneity

Not all sites behave in the same way

— Frustration

Impossibility to satisfy all conditions imposed by the Hamiltonian

(spectrum pushed up, degeneracy of ground state)

- Annealed vs quenched

Couplings, fields, etc. fluctuate or are frozen

- Quenched disorder : static pdfs of couplings, fields, etc. $f^{\text{annealed}} \leq f^{\text{quenched}}$
- Self-averageness

 $\lim_{N \to \infty} [f^{\text{quenched}}] = \lim_{N \to \infty} f^{\text{typ}}$

- Complex free-energy landscapes

Systems with short-range interactions

Divide a, say, cubic system of volume $V = L^d$ in n sub-cubes, of volume $v = \ell^d$ with V = nv


Randomness

Properties

— Spatial inhomogeneity

Not all sites behave in the same way

- Frustration

Impossibility to satisfy all conditions imposed by the Hamiltonian

(spectrum pushed up, degeneracy of ground state)

- Annealed vs quenched

Couplings, fields, etc. fluctuate or are frozen

- Quenched disorder : static pdfs of couplings, fields, etc. $f^{\text{annealed}} \leq f^{\text{quenched}}$
- Self-averageness

 $\lim_{N \to \infty} [f^{\text{quenched}}] = \lim_{N \to \infty} f^{\text{typ}}$

Complex free-energy landscapes

Phenomenology : homogeneity vs inhomogeneity

In a ferromagnet in equilibrium at temperature $T < T_c$, $\langle s_i \rangle = m(T) \ \forall i$ or $\langle s_i \rangle = -m(T) \ \forall i$ in the two homogeneous, symmetric and degenerate equilibrium states

Phenomenology : homogeneity vs inhomogeneity

In a ferromagnet in equilibrium at temperature $T < T_c$, $\langle s_i \rangle = m(T) \ \forall i$ or $\langle s_i \rangle = -m(T) \ \forall i$ in the two homogeneous, symmetric and degenerate equilibrium states

If one were to follow the time evolution of each spin in one of the two equilibrium states at $T < T_c$, one would see $\overline{s_i}(t) = m(T) + \delta_i(t)$ with $\delta_i(t)$ small time-dependent fluctuation and the overline states for a running time average $\overline{s_i}(t) = \tau^{-1} \int_t^{t+\tau} dt' s_i(t')$



Phenomenology : homogeneity vs inhomogeneity

In a spin-glass in equilibrium at temperature $T < T_c$, one expects $\langle s_i \rangle = m_i(T)$, with a different value for each i, in each inhomogeneous and degenerate equilibrium state.

There may be many different ensembles $\{m_i(T)\}$ that are equilibrium states (degeneracy, similar to what we saw in the frustrated magnets for the ground states but here in the full low T phase)

There is also the up-down symmetry $\{m_i(T)\} \mapsto \{-m_i(T)\}$

Phenomenology : homogeneity vs inhomogeneity

In a spin-glass in equilibrium at temperature $T < T_c$, one expects $\langle s_i \rangle = m_i(T)$, with a different value for each i, in each inhomogeneous and degenerate equilibrium state.

If one were to follow the time evolution of each spin in one of the possibly many equilibrium states at $T < T_c$, one would see $\overline{s_i}(t) = m_i(T) + \delta_i(t)$ with $\delta_i(t)$ small time-dependent fluctuation and the overline states for a running time average $\overline{s_i}(t) = \tau^{-1} \int_t^{t+\tau} dt' s_i(t')$



Randomness

Properties

- Spatial inhomogeneity
 - Not all sites behave in the same way, local order parameters $\{m_i\}$
- Frustration
 - Impossibility to satisfy all conditions imposed by the Hamiltonian

(spectrum pushed up, degeneracy of ground state)

- Annealed vs quenched

Couplings, fields, etc. fluctuate or are frozen

- Quenched disorder : static pdfs of couplings, fields, etc. $f^{\text{annealed}} \leq f^{\text{quenched}}$
- Self-averageness

 $\lim_{N \to \infty} [f^{\text{quenched}}] = \lim_{N \to \infty} f^{\text{typ}}$

- Complex free-energy landscapes

Randomness

Properties

- Spatial inhomogeneity
 - Not all sites behave in the same way, local order parameters $\{m_i\}$
- Frustration
 - Impossibility to satisfy all conditions imposed by the Hamiltonian

(spectrum pushed up, degeneracy of ground state)

- Annealed vs quenched

Couplings, fields, etc. fluctuate or are frozen

- Quenched disorder : static pdfs of couplings, fields, etc. $f^{\text{annealed}} < f^{\text{quenched}}$
- Self-averageness

 $\lim_{N \to \infty} [f^{\text{quenched}}] = \lim_{N \to \infty} f^{\text{typ}}$

- Complex free-energy landscapes

Fully connected Ising models

General model

 $H_J[\{s_i\}] = -\frac{1}{2} \sum_{i \neq j} J_{ij} s_i s_j$ with Ising variables $s_i = \pm 1$

 $\mathcal{O}(1)$ scaling of the local fields \Rightarrow scaling of J_{ij}

What is a local field?

It is the field felt by a selected site

 $h_i = \frac{1}{2} \sum_{ij} J_{ij} s_j$ $j(\neq i)$

and we require it to be $\mathcal{O}(1)$

Fully connected Ising models

General model

 $H_J[\{s_i\}] = -rac{1}{2} \sum_{i
eq j} J_{ij} s_i s_j$ with Ising variables $s_i = \pm 1$

 $\mathcal{O}(1)$ scaling of the local fields \Rightarrow scaling of J_{ij}

In the Curie-Weiss ferromagnetic case

$$J_{ij} = rac{J}{N}$$
 such that $h_i = rac{J}{2N} \sum_{j(\neq i)} s_j = \mathcal{O}(1)$

in the two ferromagnetic $s_i = 1 \; \forall i$ or $s_i = -1 \; \forall i$ phases

Fully connected Ising models

General model

 $H_J[\{s_i\}] = -rac{1}{2}\sum_{i
eq j}J_{ij}s_is_j$ with Ising variables $s_i=\pm 1$

 $\mathcal{O}(1)$ scaling of the local fields \Rightarrow scaling of J_{ij}

In the Curie-Weiss ferromagnetic case

$$J_{ij} = rac{J}{N}$$
 such that $h_i = rac{J}{2N} \sum_{j(\neq i)} s_j = \mathcal{O}(1)$

in the ferromagnetic $s_i = 1 \; \forall i$ or $s_i = -1 \; \forall i$ phases

In the Sherrington-Kirkpatrick disordered case $J_{ij} = \mathcal{O}(\frac{J}{\sqrt{N}}) \text{ such that } h_i \sim \frac{J}{2\sqrt{N}} \sum_{j(\neq i)} s_j = \mathcal{O}(1)$

in the PM or spin-glass phases $s_i=\pm 1 \; orall i$

Fully connected Ising models

General model

 $H_J[\{s_i\}] = -rac{1}{2}\sum_{i
eq j}J_{ij}s_is_j$ with Ising variables $s_i=\pm 1$

 $\mathcal{O}(1)$ scaling of the local fields \Rightarrow scaling of J_{ij}

In the Sherrington-Kirkpatrick disordered case

$$J_{ij} = \mathcal{O}(\frac{J}{\sqrt{N}}) \ \text{ such that } h_i \sim \frac{J}{2\sqrt{N}} \sum_{j(\neq i)} s_j = \mathcal{O}(1)$$

in the PM or spin-glass phases, say, $s_i=\pm 1$ with equal probability

One can use a Gaussian pdf

$$P(J_{ij}) = (2\pi\sigma^2)^{-1/2} \exp[-J_{ij}^2/(2\sigma^2)]$$
 with $\sigma^2 = J^2/N$

Fully connected Ising models

Even more general models (recall the K-sat problem)

 $H_J[\{s_i\}] = -\frac{1}{3!} \sum_{i \neq j \neq k} J_{ijk} s_i s_j s_k$ with Ising variables $s_i = \pm 1$

 $\mathcal{O}(1)$ scaling of the local fields \Rightarrow scaling of J_{ijk}

In the
$$p = 3$$
 Curie-Weiss ferromagnetic case
 $J_{ijk} = \frac{J}{N^{p-1}}$ such that $h_i \sim \frac{J}{2N^{p-1}} \sum_{jk(\neq i)} s_j s_k = \mathcal{O}(1)$
in the two ferromagnetic $s_i = 1$ $\forall i$ or $s_i = -1$ $\forall i$ phases

In the two terromagnetic $s_i = 1$ $\forall i$ or $s_i = -1$ $\forall i$ phases

In the p = 3 disordered case $J_{ijk} = \mathcal{O}(\frac{J}{\sqrt{N^{p-1}}}) \text{ such that } h_i \sim \frac{J}{2\sqrt{N^{p-1}}} \sum_{\substack{i \neq k (\neq i)}} s_j s_k = \mathcal{O}(1)$

in the PM or spin-glass phases $s_i = \pm 1$ with equal probability

Randomness

Properties

- Spatial inhomogeneity
 - Not all sites behave in the same way, local order parameters $\{m_i\}$
- Frustration

Impossibility to satisfy all conditions imposed by the Hamiltonian

(spectrum pushed up, degeneracy of ground state)

Annealed vs quenched

Couplings, fields, etc. fluctuate or are frozen

Quenched disorder : static pdfs of couplings

Gaussian pdf of J_{ij} with $\sigma^2 = J^2/N$

- Self-averageness

 $\lim_{N \to \infty} [f^{\text{quenched}}] = \lim_{N \to \infty} f^{\text{typ}}$

- Complex free-energy landscapes

Randomness

Properties

- Spatial inhomogeneity
 - Not all sites behave in the same way, local order parameters $\{m_i\}$
- Frustration

Impossibility to satisfy all conditions imposed by the Hamiltonian

(spectrum pushed up, degeneracy of ground state)

Annealed vs quenched

Couplings, fields, etc. fluctuate or are frozen

- Quenched disorder : static pdfs of couplings

Gaussian pdf of J_{ij} with $\sigma^2 = J^2/N$

- Self-averageness

 $\lim_{N \to \infty} [f^{\text{quenched}}] = \lim_{N \to \infty} f^{\text{typ}}$

— Complex free-energy landscapes : beyond Ginzburg-Landau

Mean-field theory for PM-FM

Fully connected Curie-Weiss Ising model

Normalize J by the size of the system N to have $\mathcal{O}(1)$ local fields

$$H = -\frac{J}{2N} \sum_{i \neq j} s_i s_j - h \sum_i s_i$$

The partition function reads $\mathcal{Z} = \int_{-1}^{1} du \ e^{-\beta N \mathbf{f}(u)}$ with $Nu = \sum_{i} s_{i}$

$$\mathbf{f}(u) = -\frac{J}{2}u^2 - hu + T\left[\frac{1+u}{2}\ln\frac{1+u}{2} + \frac{1-u}{2}\ln\frac{1-u}{2}\right]$$

Energy terms and entropic contribution stemming from $\mathcal{N}(\{s_i\})$ yielding the same u value.

Use the saddle-point, $\lim_{N\to\infty} f_N(\beta J, \beta h) = \mathbf{f}(u_{sp})$, with

$$u_{sp} = \tanh\left(\beta J u_{sp} + \beta h\right) = \langle u \rangle = m$$

Ginzburg-Landau for PM-FM

Continuous scalar statistical field theory with local aspects

Coarse-grain the spin $\phi(\vec{r}) = V_{\vec{r}}^{-1} \sum_{i \in V_{\vec{r}}} s_i$ Set h = 0



The partition function is $\mathcal{Z} = \int \mathcal{D}\phi \ e^{-\beta V \mathbf{f}(\phi)}$ with V the volume and

$$\mathbf{f}(\phi) = \int d^d r \, \left\{ \frac{1}{2} [\nabla \phi(\vec{r})]^2 + \frac{T-J}{2} \phi^2(\vec{r}) + \frac{\lambda}{4} \phi^4(\vec{r}) \right\}$$

Elastic + potential energy with the latter inspired by the results for the fully-connected model (entropy around $\phi \sim 0$ and symmetry arguments.

Uniform saddle point in the $V \to \infty$ limit : $\phi_{sp}(\vec{r}) = \langle \phi(\vec{r}) \rangle = m$ The free-energy density is $\lim_{V \to \infty} f_V(\beta, J) = \mathbf{f}(\phi_{sp})$

2nd order phase-transition

Continuous scalar statistical field theory

bi-valued equilibrium states related by symmetry



Ginzburg-Landau free-energy

Scalar order parameter

 $g=\beta J$

Features

- Spontaneous symmetry breaking below T_c
- Two equilibrium states related by symmetry $\phi \to -\phi$
- The state is chosen by a pinning field
- If the partition sum is performed over the whole phase space $\langle \phi \rangle = 0$ (a consequence of the symmetry of the action)
- Restricted statistical averages, running over half phase space, yield $\langle \phi \rangle \neq 0$
- Under a magnetic field the free-energy landscape is tilted and one of the minima becomes a metastable state
- The barrier in the free-energy landscape between the two states diverges with the size of the system implying ergodicity breaking

Features

- The function(al)s f(u) ($f(\phi(\vec{r}))$) are large deviation function(al)s determining the probability of finding an equilibrium system with u or $\phi(\vec{r})$
- The system spends $t_{\pm} \simeq e^{N\tau}$ close to each minima and it makes rapid transitions between the two

These results were not fully accepted as realistic at the time

Recall. the discussion on phase transitions & ergodicity breaking

• With p > 2-uplet interactions one finds first order phase transitions (relevant for glasses & K-sat like problems)

Fully connected SG : Sherrington-Kirkpatrick model

$$H = -\frac{1}{2} \sum_{i \neq j} J_{ij} s_i s_j - \sum_i h_i s_i$$

with J_{ij} i.i.d. Gaussian variables, $[J_{ij}] = 0$ and $[J_{ij}^2] = J^2/N = O(1/N)$.

One finds the naive free-energy landscape

$$N\mathbf{f}(\{m_i\}) = -\frac{1}{2} \sum_{i \neq j} J_{ij} m_i m_j + T \sum_{i=1}^{N} \frac{1+m_i}{2} \ln \frac{1+m_i}{2} + \frac{1-m_i}{2} \ln \frac{1-m_i}{2}$$

and the (naive) TAP equations

$$m_{isp} = \tanh(\beta \sum_{j(\neq i)} J_{ij} m_{j_{sp}} + \beta h_i)$$

that determine the restricted averages $m_i = \langle s_i \rangle = m_{isp}$.

Fully connected SG : A simple proof

The more traditional one assumes independence of the spins,

 $P(\{s_i\}) = \prod_i p_i(s_i)$

with $p_i(s_i) = \frac{1+m_i}{2} \delta_{s_i,1} + \frac{1-m_i}{2} \delta_{s_i,-1}$

and uses this form to express $\langle H
angle - T \langle S
angle$ with $S = \ln \mathcal{N}(\{s_i\})$

The energetic contribution is straightforward to evaluate

The entropic contribution is the one we already computed for the Curie-Weiss model, taking care of keeping the indices i

A more powerful proof expresses **f** as the Legendre transform of $-\beta F(h_i)$ with $m_i = N^{-1} \partial [-\beta F(h_i)] / \partial h_i$ and takes care of a "problem" to be solved in the next slides **Georges & Yedidia 91**

Missing : the Onsager reaction term

These equations are not completely correct.

The Onsager reaction term is missing.

This term represents the reaction of the spin i to itself



The magnetisation in *i* produces a field $h'_{j(i)} = J_{ji}m_i = J_{ij}m_i$ on spin *j* This field induces a magnetisation $m'_{j(i)} = \chi_{jj}h'_{j(i)} = \chi_{jj}J_{ij}m_i$ on the spin *j*. This magnetisation produces a field $h'_{i(j)} = J_{ij}m'_{j(i)} = J_{ij}\chi_{jj}J_{ij}m_i$ on site *i*. The equilibrium fluctuation-dissipation relation between susceptibilities and connected correlations implies $\chi_{jj} = \beta \langle (s_j - \langle s_j \rangle)^2 \rangle = \beta(1 - m_j^2)$ and one then has $h'_{i(j)} = \beta(1 - m_j^2)J_{ij}^2m_i$

The Onsager reaction term

The idea of Onsager – or *cavity method* – is that one has to study the ordering of the spin i in the absence of its own effect on the rest of the system.

The total field produced by the sum of $h'_{i(j)} = \beta(1 - m_j^2)J_{ij}^2m_i$ over all the spins j with which it can connect, has to be subtracted from the mean-field created by the other spins in the sample, *i.e. the total local field* should be

$$h_i^{\text{loc}} = \sum_{j(\neq i)} J_{ij} m_j - \beta m_i \sum_{j(\neq i)} J_{ij}^2 (1 - m_j^2)$$

recall that $J_{ij} = \mathcal{O}(1/\sqrt{N})$. Finally, the TAP equations read

$$m_i = \tanh\left\{\sum_{j(\neq i)} \left[\beta J_{ij}m_j - \beta^2 m_i J_{ij}^2 (1 - m_j^2)\right]\right\}$$

Orders of magnitude

The Thouless-Anderson-Palmer (TAP) equations read

$$m_{i} = \tanh \left\{ \sum_{j(\neq i)} [\beta J_{ij} m_{j} - \beta^{2} m_{i} J_{ij}^{2} (1 - m_{j}^{2})] \right\}$$

The first term in the rhs $\sum_{j(\neq i)} J_{ij}m_j \simeq \frac{1}{\sqrt{N}}\sqrt{N} = \mathcal{O}(1)$ because of the central limit theorem.

The second term $\sum_{j(\neq i)} J_{ij}^2(1-m_j^2) \simeq \frac{1}{N} N = \mathcal{O}(1)$ because all terms in the sum are positive definite $(m_j \leq 1 \ \forall j)$

Recall that $m_i = \langle s_i \rangle$

Orders of magnitude

The Thouless-Anderson-Palmer (TAP) equations read

$$m_i = \tanh\left\{\sum_{j(\neq i)} \left[\beta J_{ij}m_j - \beta^2 m_i J_{ij}^2(1-m_j^2)\right]\right\}$$

The first term in the rhs $\sum_{j(\neq i)} J_{ij}m_j \simeq \frac{1}{\sqrt{N}}\sqrt{N} = \mathcal{O}(1)$ because of the central limit theorem.

The second term $\sum_{j(\neq i)} J_{ij}^2 (1 - m_j^2) \simeq \frac{1}{N} N = \mathcal{O}(1)$ because all terms in the sum are positive definite $(m_j \leq 1 \ \forall j)$

Exercise

Check that higher order loops are negligible, since sub-leading in powers of N



Orders of magnitude

The Thouless-Anderson-Palmer (TAP) equations read

$$m_i = \tanh\left\{\sum_{j(\neq i)} \left[\beta J_{ij}m_j - \beta^2 m_i J_{ij}^2(1-m_j^2)\right]\right\}$$

The first term in the rhs $\sum_{j(\neq i)} J_{ij}m_j \simeq \frac{1}{\sqrt{N}}\sqrt{N} = \mathcal{O}(1)$ because of the central limit theorem.

The second term $\sum_{j(\neq i)} J_{ij}^2 (1 - m_j^2) \simeq \frac{1}{N} N = \mathcal{O}(1)$ because all terms in the sum are positive definite $(m_j \leq 1 \forall j)$

Exercise

Check that in the Curie Weiss model $J_{ij} = J/N$ there is no need of Onsager terms



Landscape

Free-energy density at fixed randomness

The TAP equations are the extremization conditions on the TAP free-energy



Summary

Local & global order parameters

 $m_i \equiv \langle s_i \rangle$ = 0 at $T \ge T_c$ $\neq 0$ at $T < T_c$

Magnetization

$$m = rac{1}{N} \sum\limits_{i} m_i = 0$$
 at all temperatures

Edwards-Anderson order parameter

$$\begin{split} q^{\alpha}_{\text{EA}} &\equiv \frac{1}{N} \sum_{i} (m^{\alpha}_{i})^{2} = \frac{1}{N} \sum_{i} \langle s_{i} \rangle^{2}_{\alpha} \\ &= 0 \text{ at } T \geq T_{c} \\ &\neq 0 \text{ at } T < T_{c} \end{split}$$



- How do we know that the TAP equations are correct?
 Phase transition in the SK model
- Back to landscapes
- Statistical averages
- Real replicas
- Replica method



- How do we know that the TAP equations are correct?
 Phase transition in the SK model
- Back to landscapes
- Statistical averages
- Real replicas
- Replica method

Phase transition

For large N one expects $J_{ij}^2 \simeq [J_{ij}^2] = J^2/N$ with $J = \mathcal{O}(1)$

Simplification $m_i = \tanh\left\{\beta\sum_{j(\neq i)}J_{ij}m_j - \beta^2 m_i \frac{J^2}{N}\sum_{j(\neq i)}(1-m_j^2)\right\}$

A 2nd order phase transition \Rightarrow $m_i \simeq 0$ at $T \stackrel{<}{\sim} T_c$ then using $anh y \sim y$

The TAP equations become $m_i \sim \beta \sum_{j(\neq i)} J_{ij} m_j - \beta^2 J^2 m_i$

Diagonalize this eq. going to the basis of eigenvectors of the J_{ij} matrix

The eqs read
$$m_\lambda \sim eta \Big(J_\lambda - eta J^2 \Big) m_\lambda$$

The notation we use is such that

 J_{λ} is an eigenvalue of the J_{ij} matrix associated to the eigenvector \vec{v}_{λ} m_{λ} represents the projection of \vec{m} on the eigenvector \vec{v}_{λ} , $m_{\lambda} = \vec{v}_{\lambda} \cdot \vec{m}$ with \vec{m} the *N*-vector with components m_i , $\vec{m} = (m_1, \dots, m_N)$

Phase transition

If we add a weak external field the eqs read $m_{\lambda} \sim \beta (J_{\lambda} - \beta J^2) m_{\lambda} + \beta h_{\lambda}^{\text{ext}}$ The variation with respect to the field at linear order is

$$\frac{\partial m_{\lambda}}{\partial h_{\lambda}^{\text{ext}}}\Big|_{\vec{h}^{\text{ext}}=\vec{0}} = \beta (J_{\lambda} - \beta J^2) \left. \frac{\partial m_{\lambda}}{\partial h_{\lambda}^{\text{ext}}} \right|_{\vec{h}^{\text{ext}}=\vec{0}} + \beta$$

and the staggered susceptibility (of the projection on \vec{v}_{λ})

$$\chi_{\lambda} \equiv \left. \frac{\partial m_{\lambda}}{\partial h_{\lambda}^{\text{ext}}} \right|_{\vec{h}^{\text{ext}} = \vec{0}} = \beta \left(1 - \beta J_{\lambda} + (\beta J)^2 \right)^{-1}$$

Random matrix theory tells us that the eigenvalues of the random matrix $J_{ij} = \mathcal{O}(1/\sqrt{N})$ are distributed with the Wigner semi-circle law and the largest eigenvalue is $J_{\lambda}^{\max} = 2J$

The staggered susceptibility of staggered magnetization in the direction of the largest eigenvalue diverges at $\beta_c J = 1$ the correct value

Phase transition

If we add a weak external field the eqs read $m_{\lambda} \sim \beta (J_{\lambda} - \beta J^2) m_{\lambda} + \beta h_{\lambda}^{\text{ext}}$ The variation with respect to the field at linear order is

$$\frac{\partial m_{\lambda}}{\partial h_{\lambda}^{\text{ext}}}\Big|_{\vec{h}^{\text{ext}}=\vec{0}} = \beta (J_{\lambda} - \beta J^2) \left. \frac{\partial m_{\lambda}}{\partial h_{\lambda}^{\text{ext}}} \right|_{\vec{h}^{\text{ext}}=\vec{0}} + \beta$$

and the *staggered susceptibility* (of the projection on \vec{v}_{λ})

$$\chi_{\lambda} \equiv \left. \frac{\partial m_{\lambda}}{\partial h_{\lambda}^{\text{ext}}} \right|_{\vec{h}^{\text{ext}} = \vec{0}} = \beta \left(1 - \beta J_{\lambda} + (\beta J)^2 \right)^{-1}$$

Random matrix theory tells us that the eigenvalues of the random matrix J_{ij} are distributed with the Wigner semi-circle law

For
$$J_{ij} = \mathcal{O}(1/\sqrt{N})$$
 the largest eigenvalue is $J_{\lambda}^{\max} = 2J$

The staggered susceptibility for the largest eigenvalue diverges at $\beta_c J = 1$

Without the reaction term the divergence is at the inexact value $T^* = 2T_c$



- How do we know that the TAP equations are correct?
 Phase transition in the SK model
- Back to landscapes
- Statistical averages
- Real replicas
- Replica method



Free-energy density at fixed randomness

The TAP equations are the extremization conditions on the TAP free-energy

 $\frac{\delta F_J^{\rm tap}(\{m_i\})}{\delta m_j}=0$ The stability of the solutions is determined by the Hessian

 $\frac{\delta^2 F_J^{\rm tap}(\{m_i\})}{\delta m_i \delta m_k}$







Features

At fixed randomness

- There are N local order parameters, $m_i, i = 1, \ldots, N$
- The saddle-points are **heterogeneous**: m_i differ from site to site
- At high temperatures only one trivial solution $\{m_i = 0\}$
- At low temperatures the TAP equations have many solutions $\{m_i^{\alpha}\}$, which are extrema of the TAP free-energy landscape, *i.e.* saddles of all types, $\alpha = 1, \ldots, \mathcal{N}_J$
- For each solution $\{m_i^{\alpha}\}$, there is also $\{-m_i^{\alpha}\}$ but apart from this trivial doubling, the remaining solutions are not related by symmetry
- The TAP free-energy can take different values at different $\{m_i^{\alpha}\} \Rightarrow f_{tap}^{\alpha}$
Features

All this is reshuffled for another realization of disorder

- Still N local order parameters, $m_i, i = 1, \dots, N$
- The TAP equations have other solutions $\{m_i^{lpha}\}$, extrema of the TAP freeenergy landscape, $F_J^{
 m tap}$, labelled by $lpha=1,\ldots,\mathcal{N}_J$
- A global order parameter ? The simplest guess $\frac{1}{N}\sum_{i=1}^{N}m_{i}^{\alpha}$ cannot be since it is = 0 One expects as many positive as negative m_{i} s and similarity in

all respects. Another try

$$q_{\rm EA}^{\alpha} = \frac{1}{N} \sum_{i=1}^{N} (m_i^{\alpha})^2$$

• "Typicality expected" (though see below for equilibrium states)

Features

Numbers of metastable states

- N local order parameters, $m_i, i = 1, \ldots, N$
- The TAP equations have many solutions $\{m_i^{lpha}\}$, extrema of the TAP freeenergy landscape, $lpha=1,\ldots,\mathcal{N}_J$
- One can count how many saddles of each kind exist and their **complexity** $\mathcal{N}_J = \prod_{i=1}^N \int_{-1}^1 dm_i \ \delta(m_i - m_i^{\alpha}) \qquad \Sigma = \ln \mathcal{N}$
- how many of these at each level of free-energy density, by inserting a delta-function $\delta(f_J^{ ext{tap}}(\{m_i^lpha\}) f) \Rightarrow \mathcal{N}_J(f)$
- How many with a given stability $\mathcal{N}_J(f,K)$ with K the number of positive eigenvalues of the Hessian, with adequate delta-functions



- How do we know that the TAP equations are correct?
 Phase transition in the SK model
- Back to landscapes
- Statistical averages
- Real replicas
- Replica method

At fixed interactions

The average of a generic observable is

$$\langle O \rangle = \sum_{\alpha} w_{\alpha} \langle O \rangle_{\alpha}$$

In the **FM case**, each state $(\langle \phi \rangle = \pm \phi_0)$ has weight $w_{\pm} = 1/2$ and the sum is $\langle O \rangle = \frac{1}{2} \langle O \rangle_+ + \frac{1}{2} \langle O \rangle_-$ with $\langle O \rangle_{\pm}$ the average in each of the states. For instance, the averaged magnetization vanishes if one sums over the \pm states or it is different from zero if one restricts the sum to only one of them.



FM case

The dashed blue line with two minima $\pm |\phi_0|$ If we have many more?

At fixed interactions

The average of a generic observable is

$$\langle O \rangle = \sum_{\alpha} w_{\alpha} \langle O \rangle_{\alpha}$$

In the FM case, each state ($\langle \phi \rangle = \pm \phi_0$) has weight $w_{\pm} = 1/2$ and the sum is $\langle O \rangle = \frac{1}{2} \langle O \rangle_+ + \frac{1}{2} \langle O \rangle_-$ with $\langle O \rangle_{\pm}$ the average in each of the states. For instance, the averaged magnetization vanishes if one sums over the \pm states or it is different from zero if one restricts the sum to only one of them.



$$\begin{aligned} &\mathsf{FM case} \ f_{+} = f_{-} \\ & w_{\pm} = \frac{e^{-\beta N f_{\pm}}}{e^{-\beta N f_{+}} + e^{-\beta N f_{-}} + e^{-\beta N f_{0}}} \simeq \frac{1}{2} \\ & w_{0} = \frac{e^{-\beta N f_{0}}}{e^{-\beta N f_{+}} + e^{-\beta N f_{-}} + e^{-\beta N f_{0}}} \ll w_{\pm} \end{aligned}$$

At fixed randomness

The average of a generic observable is

$$\langle O \rangle = \sum_{\alpha} w_{\alpha} \langle O \rangle_{\alpha}$$

For systems with quenched randomness

$$w_{\alpha}^{J} = \frac{e^{-\beta N \mathbf{f}_{\alpha}^{J}}}{\sum_{\gamma} e^{-\beta N \mathbf{f}_{\gamma}^{J}}}$$

where we added a super-script to the weight w

 J indicates that the weights depend on the the disorder realization

and α is a label that identifies the TAP solution



One can sum over all saddles irrespectively of their stability. Higher lying ones will be exponentially suppressed or will dominate depending on $\Sigma_J(f, K)$

At fixed randomness

The average of a generic observable is

$$\langle O \rangle = \sum_{\alpha} w_{\alpha} \langle O \rangle_{\alpha}$$

For systems with **quenched randomness**

$$w_{\alpha}^{J} = \frac{e^{-\beta N \mathbf{f}_{\alpha}^{J}}}{\sum_{\gamma} e^{-\beta N \mathbf{f}_{\gamma}^{J}}}$$

The sum over α , in the case in which there are an exponential in N number of TAP solutions, can be replaced by an integral over **f**

$$\langle O \rangle = \mathcal{Z}^{-1}(\beta, J) \int d\mathbf{f} \ e^{-\beta [N\mathbf{f} - T \ln \mathcal{N}_J(\mathbf{f}, \beta)]} \ O(\mathbf{f}, \beta)$$

 \mathcal{N}_J is the number of solutions to the TAP eqs. with free-energy density **f**.

For $N \to \infty$ the integral is dominated by the saddle point

$$\frac{1}{T} = \frac{1}{N} \frac{\partial \ln \mathcal{N}_J(\mathbf{f}, \beta)}{\partial \mathbf{f}} \bigg|_{\mathbf{f}_{sp}} = \frac{1}{N} \frac{\partial \Sigma_J(\mathbf{f}, \beta)}{\partial \mathbf{f}} \bigg|_{\mathbf{f}_{sp}} \quad \text{complexity}$$

Consequences

The equilibrium free-energy f is given by the saddle-point evaluation of the partition sum:

$$f = \mathbf{f}_{sp} - \frac{T}{N} \ln \mathcal{N}_J(\mathbf{f}_{sp}, \beta)$$

The rhs is the Landau free-energy of the problem, with \mathbf{f}_{sp} playing the role of the energy and $N^{-1} \ln \mathcal{N}_J(\mathbf{f}_{sp}, \beta)$ of the entropy

The contribution of the complexity or configurational entropy contribution is negative and in some cases higher lying extrema (metastable states) can dominate the partition sum with respect to lower lying ones if $\ln N_J(\mathbf{f}_{sp}, \beta) \propto N$

This feature is proposed to describe **super-cooled liquids**.

A global observable

Effect of multi-states

What is the expression of the global order parameter once one takes into account the multi-states?

$$q \equiv \frac{1}{N} \sum_{i} \langle s_i \rangle^2 = \frac{1}{N} \sum_{i} (\sum_{\alpha} w_{\alpha}^J m_i^{\alpha})^2 = \frac{1}{N} \sum_{i} \sum_{\alpha} w_{\alpha}^J m_i^{\alpha} \sum_{\beta} w_{\beta}^J m_i^{\beta}$$

note that this is different from $q_{\mathrm{EA}} = rac{1}{N}\sum\limits_{i}(m_{i}^{lpha})^{2}$

Defining now $\left| q_{\alpha\beta} \equiv \frac{1}{N} \sum_{i} m_{i}^{\alpha} m_{i}^{\beta} \right|$

an overlap between different states

and $P_J(q') \equiv \sum_{\alpha\beta} w^J_{\alpha} w^J_{\beta} \,\delta(q' - q_{\alpha\beta})$

we obtain

in
$$q \equiv \frac{1}{N} \sum_{i} \langle s_i \rangle^2 = \int dq' P_J(q') q'$$



- How do we know that the TAP equations are correct?
 Phase transition in the SK model
- Back to landscapes
- Statistical averages
- Real replicas
- Replica method

Overlaps between replicas

Take one sample and run it, with e.g. Monte Carlo, until it reaches equilibrium, measure the spin configuration $\{s_i\}$.

Re-initialize the same sample (same J_{ij}), run it again until it reaches equilibrium, & measure the spin configuration $\{\sigma_i\}$.

Construct the overlap $q_{s\sigma} \equiv N^{-1} \sum_{i=1}^{N} s_i \sigma_i$.

In a PM system the overlap will typically vanish as, say, $N^{-1/2}$



Many repetitions for a system with $N\gg 1$

$$P(q_{s\sigma}) = \delta(q_{s\sigma})$$

Overlaps between replicas

Take one sample and run it, with e.g. Monte Carlo, until it reaches equilibrium, measure the spin configuration $\{s_i\}$.

Re-initialize the same sample (same J_{ij}), run it again until it reaches equilibrium, & measure the spin configuration $\{\sigma_i\}$.

Construct the overlap $q_{s\sigma} \equiv N^{-1} \sum_{i=1}^{N} s_i \sigma_i$.

In a FM system there are four possibilities



Pdf of overlaps between replicas at fixed randomness

Sherrington-Kirkpatrick model with N=4096 at $T=0.4\,T_c$



Finite size corrections in the Sherrington-Kirkpatrick model

Aspelmeier, Billoire, Marinari & Moore (2007)

Ooverlaps between replicas at fixed randomness

Sherrington-Kirkpatrick model with N = 4096 at $T = 0.4 T_c$



Data in each panel for a different realization of the random couplings

Each sample has peaks at $q_{s\sigma} = \pm q_{\rm EA} \simeq \pm 0.75$:

two configurations in the same (or the spin-reversed) state

Overlaps between replicas at fixed randomness

Sherrington-Kirkpatrick model with N = 4096 at $T = 0.4 T_c$



Data in each panel for a different realization of the random couplings

 $0.75 \simeq q_{
m EA} < 1$ and the width of the peaks at $q_{s\sigma} = \pm q_{
m EA}$: due to $0 < T < T_c$ and finite N, respectively

Overlaps between replicas at fixed randomness

Sherrington-Kirkpatrick model with N = 4096 at $T = 0.4 T_c$



Data in each panel for a different realization of the random couplings

Most samples also have peaks at $|q_{s\sigma}| < q_{\rm EA}$: replicas $\{s_i\}$ and $\{\sigma_i\}$ falling in different states

Overlaps between replicas at fixed randomness

SK model with $N \to \infty$ at $T < T_c$



What happens if one averages $P_J(q)$ over disorder

Disordered averaged pdf of overlaps $P(q) = [P_J(q)]$

Parisi 79-82 prescription for the replica symmetry breaking Ansatz yields



Thermodynamic quantities, in particular the equilibrium free-energy density are expressed as functions of P(q).

The equilibrium free-energy density predicted by the replica theory was confirmed by **Guerra & Talagrand 00-04** indepedent mathematical-physics mthods.



- How do we know that the TAP equations are correct?
 Phase transition in the SK model
- Back to landscapes
- Statistical averages
- Real replicas
- Replica method

Typical vs. averaged

TAP vs. Replicas

Precursors

Look at an integer parameter n

and its $n \rightarrow 0$ limit

In 1972 Fortuin and Kasteleyn studied the **Potts model** with n components :

- n=2 Ising
- n = 1 percolation
- n = 0 random resistors

Use the identify $x^n = \exp(n \ln x)$ and expand around n = 0:

$$\lim_{n \to 0} x^n = 1 + n \ln x + \mathcal{O}(n^2)$$

A sketch

$$-\beta[f_J] = \lim_{N \to \infty} \frac{\left[\ln Z_N(\beta, J)\right]}{N} = \lim_{N \to \infty} \lim_{n \to 0} \frac{\left[Z_N^n(\beta, J)\right] - 1}{Nn}$$

 Z_N^n partition function of *n* independent copies of the system : replicas.

Gaussian average over disorder : coupling between replicas

$$\sum_{a} \sum_{i \neq j} J_{ij} s_i^a s_j^a \Rightarrow \sum_{i \neq j} \left(\sum_{a} s_i^a s_j^a \right)^2$$

Quadratic decoupling with the Hubbard-Stratonovich trick

$$Q_{ab}\sum_{i}s_{i}^{a}s_{i}^{b} + \frac{1}{2}Q_{ab}^{2}$$

 Q_{ab} is a 0×0 matrix but it admits an interpretation in terms of **overlaps** The elements of Q_{ab} can evaluated by saddle-point if one exchanges the limits $N \to \infty \ n \to 0$ with $n \to 0 \ N \to \infty$.

In more detail

 Z_N^n partition function of *n* independent copies of the system : replicas.

$$Z_N^n(\beta, J) = \sum_{\substack{\{s_i^{(1)} = \pm 1\} \\ \text{notation } \mathsf{Tr}_{\{s_i^a\}}}} \dots \sum_{\substack{\{e^{-\beta \sum_{a=1}^n \sum_{i \neq j} J_{ij} s_i^a s_j^a \\ e^{-\beta \sum_{a=1}^n \sum_{i \neq j} J_{ij} s_i^a s_j^a }}$$

One can exchange the order of the trace and the average over disorder

$$\left[Z_N^n(\beta,J)\right] = \operatorname{Tr}_{\{s_i^a\}} \int \prod_{i \neq j} dJ_{ij} P(J_{ij}) \ e^{-\beta \sum_{a=1}^n \sum_{i \neq j} J_{ij} s_i^a s_j^a}$$

$$[Z_N^n(\beta, J)] = \mathrm{Tr}_{\{s_i^a\}} \ e^{-\beta H_{\mathrm{eff}}[\{s_i^a\}]}$$

 $H_{\text{eff}}[\{s_i^a\}]$ does not have any randomness but couples the replicas

$$\left|\sum_{a}\sum_{i\neq j}J_{ij}s_{i}^{a}s_{j}^{a}\right| \Rightarrow \sum_{i\neq j}\left(\sum_{a}s_{i}^{a}s_{j}^{a}\right)^{2}$$

In more detail

$$[Z_N^n(\beta,J)] = \mathrm{Tr}_{\{s_i^a\}} \ e^{-\beta H_{\mathrm{eff}}[\{s_i^a\}]}$$

 $H_{\mathrm{eff}}[\{s^a_i\}]$ does not have any randomness but couples the replicas

$$\sum_{i \neq j} \left(\sum_{a} s_i^a s_j^a \right)^2 = \sum_{i \neq j} \sum_{a} \sum_{b} s_i^a s_j^a s_i^b s_j^b \sim \sum_{ab} \sum_{i} s_i^a s_i^b \sum_{j} s_j^a s_j^b$$

One sees Q_{ab} here, introduce their definition via a delta or apply Hubbard-Stratonovich

Once this done, one can exchange the trace (the sum over spin configurations) and the integral over Q_{ab} and end up with

$$[Z_N^n(\beta,J)] \propto \int \prod_{ab} dQ_{ab} e^{-F(Q_{ab})}$$

For the SK model

$$Q_{ab} = q_{ab}$$
 and $p = 2$

$$\begin{split} \beta F(q_{ab}) &= -\frac{N\beta^2 J^2}{2} \left[-\sum_{a\neq b} q_{ab}^p + n \right] - N \ln \zeta(q_{ab}) ,\\ \zeta(q_{ab}) &= \sum_{s_a} e^{-\beta H(q_{ab},s_a)} ,\\ H(q_{ab},s_a) &= -J \sum_{ab} q_{ab} s_a s_b - h \sum_a s_a , \end{split}$$

In more detail

$$[Z_N^n(\beta,J)] = \operatorname{Tr}_{\{s_i^a\}} e^{-\beta H_{\text{eff}}[\{s_i^a\}]} \propto \int \prod_{ab} dQ_{ab} e^{-F(Q_{ab})}$$

 $H_{\mathrm{eff}}[\{s^a_i\}]$ and Q_{ab} do not have any randomness but couple the replicas

The elements of Q_{ab} can be evaluated by saddle-point if one exchanges the limits $N \to \infty \ n \to 0$ with $n \to 0 \ N \to \infty$.

At the saddle-point level one identifies $Q^{sp}_{ab} = N^{-1} \langle \sum_i s^a_i s^b_i \rangle$

The spin glass transition is from the paramagnetic state with $Q_{a\neq b} = 0$ to a spin glass state with $Q_{a\neq b} \neq 0$ as the temperature is decreased.

SK model: replica symmetric Ansatz

Permutation symmetry between replicas \Rightarrow

Insert $Q_{a\neq b} = q$ and $Q_{aa} = 1$ in the effective Hamiltonian

Saddle-point with respect to q and $n \rightarrow 0$

$$q = \int_{-\infty}^{\infty} \frac{dz}{\sqrt{2\pi}} e^{-z^2/2} \tanh\left(\beta J \sqrt{q}z\right)$$

Note the similarity with the equation for m in the Curie-Weiss model

$$q=0$$
 for $T\geq T_c=J$

 $q \neq 0$ for $T < T_c = J$

Problem I Is this solution stable? No

Problem II Does it have a zero-temperature vanishing entropy? No

Problem III Ground state energy density $e = -0.77 \pm 0.01$ while the replica symme-

tric value e = -0.798, is three standard deviations smaller (in units of J)

SK model: one step replica symmetry breaking

Permutation symmetry broken



 $n \times n$ matrix divided in diagonal blocks of size $m \times m$ and the rest

SK model: one step replica symmetry breaking



Problem I Stability : improved but not solved

Problem II Zero-temperature entropy : improved but not solved

Problem III e closer to numerical value

SK model: two step replica symmetry breaking

Permutation symmetry broken



n imes n matrix divided in diagonal blocks of size $m_2 imes m_2$, and the rest in blocks of size $m_1 imes m_1$ and the rest

SK model: two step replica symmetry breaking



Problem I Stability : improved but not solved

Problem II Zero-temperature entropy : improved but not solved

Problem III e closer to numerical value

SK model: full replica symmetry breaking

Blocks of size m_i with parameter q_i

e.g. for replica symmetric case one block a single *q*.

 ∞ number of breaking steps, that is, of blocks

 $m_i \mapsto x$ and the parameter $q_i \mapsto q(x)$

$$[\langle s_i \rangle^2] = \int_0^1 dx \ q(x) = \int \frac{dx}{dq} \ dq \ q(x) = \int dq \ P(q) \ q$$

with

 $P(q) = \frac{dx}{dq}$

Problem I Stability : solved Problem II Zero-temperature entropy : solved S = 0Problem III *e* in agreement with numerical value within numerical accuracy e = -0.7633



- Lectures in English on the blackboard
 - Either simultaneous broadcasting via zoom (see how it goes) or inde-
 - pendent zoom presentation (other time-slot) for absent students.
- Three exercise sessions (Tds).
- Lecture notes will be available at

www.lpthe.jussieu.fr/~leticia/enseignement.html

(the 2019 ones are already here).

Exam modality will be decided according to the sanitary situation & number of students:

Either conventional written exam (see recent years examples) or the study, written description (report) + oral presentation of a subject (paper) ???